

NATIONAL HEALTH AND MORBIDITY SURVEY 2017

ADOLESCENT HEALTH SURVEY 2017

JOHOR



**NATIONAL HEALTH AND MORBIDITY
SURVEY 2017**

(NMRR-16-698-30042)

ADOLESCENT HEALTH SURVEY 2017

JOHOR

Contributors

The following persons had contributed in the interpretation of findings, discussions on implications, conclusions and/or drawing recommendations for this report.

(In alphabetical order)

Abdul Aiman Abd Ghani, Amal Shamsudin, Azli Baharudin @Shaharuddin, Azriman Rosman, Chan Ying Ying, Chandrika A/p Jeevananthan, Cheong Siew Man, Faizah Paiwai, Fazila Haryati Ahmad, Fazly Azry Abdul Aziz, Halizah Mat Rifin, Hamizatul Akmal Abd Hamid, Hasimah Ismail, Jane Ling Miaw Yn, Lim Kuang Kuay, Maisarah Omar, Mohamad Aznuddin Abd Razak, Mohamad Fuad Mohamad Anuar, Mohamad Hasnan Ahmad, Mohd Azahadi Omar, Mohd Hatta Mutalip, Mohd Hazrin Hasim@Hashim, Muhammad Fadhli Mohd Yusoff, Muhammad Suhaimi Mohamad Idrus, Muslimah Yusof, Natifah Che Salleh, Nazirah Alias, Nik Rubiah Nik Abdul Rashid, Nizam Baharom, Noor Aliza Lodz, Noor Ani Ahmad, Noraida Mohamad Kasim, Noraryana Hassan, Norazizah Ibrahim Wong, Norharlina Bahar, Norli Abdul Jabbar, Norlida Zulkafly, Normala Ibrahim, Norzawati Yoep, Nur Azna Mahmud, Nur Hazwani Mohd Hasri, Nur Liana Abdul Majid, Nurashikin Ibrahim, Nurrul Ashikin Abdullah, Rajini Sooryanarayana, Rashidah Ambak, Rasidah Jamaluddin, Rimah Melati Abd. Ghani, Rozanim Kamaruddin, Ruhaya Salleh, Rushidi Ramly, Rusidah Selamat, S Maria Awaluddin, Saidatul Norbaya Buang, Sayan a/l Pan, Shahibul Bariah Mat Ghani, Shubash Shander Ganapathy, Siti Noafika Anwar, Syafinaz Mohd Sallehuddin, Tania Gayle Robert, Tan Lee Ann, Tee Guat Hiong, Thamil Arasu Saminathan, Ting Teck Pei, Wan Salwina Wan Ismail, Wan Shakira Rodzlan Hasani, Yaw Siew Lian.

Editorial Reviewers

Chan Ying Ying, Jane Ling Miaw Yn, Norzawati Yoep, Noor Aliza Lodz, Rajini Sooryanarayana
Tania Gayle A/P Robert, Tan Lee Ann

Editors

S Maria Awaluddin, Muhammad Fadhli Mohd Yusoff, Tahir Aris

© 2017, Institute for Public Health, National Institutes of Health, Ministry of Health Malaysia, Kuala Lumpur.

Perpustakaan Negara Malaysia

Cataloguing-in-Publication Data

National Health and Morbidity Survey 2017 (NMRR-16-698-30042)

Johor Adolescent Health Survey 2017

ISBN: 978-983-2387-58-9

MOH/S/IKU/90.18(RR)

Suggested citation:

Institute for Public Health (IPH) 2017. National Health and Morbidity Survey (NHMS) 2017: Adolescent Health Survey 2017, Johor

Disclaimer:

The views expressed in this report are those of the authors alone and do not necessarily represent the opinions of the other investigators participating in the survey, nor the view or policy of the Ministry of Health.

Produced and Distributed by:

National Health and Morbidity Survey 2017: Adolescent Health Survey 2017

Institute for Public Health

National Institutes of Health

Ministry of Health Malaysia

Jalan Bangsar, 50590 Kuala Lumpur

Federal Territories of Kuala Lumpur, Malaysia

Tel: +603-2297-9400

Fax: +603-2282-3114

Any enquiries or comments on this report should be directed to:

Principal Investigator

National Health and Morbidity Survey 2017: Adolescent Health Survey 2017

Institute for Public Health

National Institutes of Health

Ministry of Health Malaysia

Jalan Bangsar, 50590 Kuala Lumpur

Federal Territories of Kuala Lumpur, Malaysia

Tel: +603-2297-9400

Fax: +603-2282-3114

Published by Institute for Public Health, National Institutes of Health, Ministry of Health Malaysia.

Acknowledgement

The authors would like to thank the Director General of Health Malaysia and the National Health and Morbidity Survey (NHMS) Steering Committee Team for supporting this research project via financial grant and technical support. The research team members would like to express their gratitude to the Director General of Education, Ministry of Education Malaysia, as well as the Director of Family Health Development Division and Director of Nutrition Division, Ministry of Health Malaysia. Our appreciation is also extended towards the various State Education Departments, District Education Offices, School Principals and the selected school officers who assisted in this study successfully. And last but not least, all students from 212 schools whom have participated in this study.

LIST OF ABBREVIATIONS

AHS	Adolescent Health Survey
CDC	Centre Disease Control
GSHS	Global School-based Student Health survey
IAT	Internet Addiction Test
MVIAT	Malay Version Internet Addiction Test
UNAIDS	Joint United Nations Programme on HIV and AIDS
UNICEF	United Nations Children's Fund
UNESCO	United Nations Educational, Scientific and Cultural Organization
WHO	World Health Organization
YRBSS	Youth Risk Behavior Surveillance System

TABLE OF CONTENTS

1.0	INTRODUCTION	1
1.1	Research in Adolescent Health in Malaysian Context.....	1
1.2	Objectives.....	5
1.2.1	General Objectives	5
1.2.2	Specific Objectives.....	5
1.2.3	The NHMS 2017 Organisation Team.....	5
2.0	METHODOLOGY AND SAMPLING DESIGN	9
2.1	Target Population.....	9
2.2	Sampling Frame.....	9
2.3	Sample Size Determination	9
2.4	Sampling Design	11
2.5	Ethical Approval and Consent Forms	11
2.6	Survey Instruments	11
2.7	Data Collection	12
2.8	Data Management.....	12
2.8.1	Data Operation Centre	13
2.8.2	Data Analysis	15
3.0	FINDINGS.....	16
3.1	General Findings.....	16
3.2	Alcohol Consumption	19
3.2.1	Introduction.....	19
3.2.2	Objectives	19
3.2.3	Variable definitions.....	20
3.2.4	Findings.....	20
3.2.5	Discussion/ Conclusion	21
3.2.6	Recommendations.....	21
3.2.7	References	22
3.3.	Dietary Behaviours	26
3.3.1	Introduction.....	26
3.3.2	Objectives	26
3.3.4	Findings.....	27
3.3.5	Discussion/ Conclusion	29
3.3.6	Recommendations.....	30
3.3.7	References	30

3.4 Drug Use	35
3.4.1 Introduction	35
3.4.2 Objectives	36
3.4.3 Variable Definitions	36
3.4.4 Findings.....	37
3.4.5 Discussion/ Conclusion	37
3.4.6 Recommendations.....	38
3.4.7 References	38
3.5 Hygiene (Including Oral Health)	43
3.5.1 Introduction	43
3.5.2 Objectives	43
3.5.3 Variable Definitions	44
3.5.4 Findings.....	44
3.5.5 Discussion/ Conclusion	46
3.5.6 Recommendations.....	47
3.5.7 References	47
3.6 Internet Use and Addiction	58
3.6.1 Introduction	58
3.6.2 Objectives	59
3.6.3 Variable Definitions	59
3.6.4 Findings.....	59
3.6.5 Discussion/ Conclusion	60
3.6.5 Discussion/ Conclusion	60
3.6.6 Recommendations.....	60
3.6.7 References	61
3.7 Mental Health Problems	64
3.7.1 Introduction	64
3.7.2 Objectives	64
3.7.3 Variable definitions.....	64
3.7.4 Findings.....	65
3.7.5 Discussion/ Conclusion	65
3.7.6 Recommendations.....	65
3.7.7 References	66
3.8 Physical Activity.....	70
3.8.1 Introduction	70
3.8.2 Objectives	70

3.8.3	Variable Definitions	70
3.8.4	Findings.....	70
3.8.5	Discussion/ Conclusion	71
3.8.6	Recommendations.....	71
3.8.7	References	72
3.9.	Protective Factors.....	74
3.9.1	Introduction	74
3.9.2	Objectives	74
3.9.3	Variable Definitions	74
3.9.4	Findings.....	75
3.9.5	Discussions/ Conclusion	76
3.9.6	Recommendations.....	76
3.9.7	References	76
3.10	Sexual Behaviours that contribute to HIV Infection, Other STIs and Unintended Pregnancy.....	81
3.10.1	Introduction	81
3.10.2	Objectives	81
3.10.3	Variable Definitions	81
3.10.4	Findings.....	82
3.10.5	Discussion/ Conclusion	82
3.10.6	Recommendations.....	82
3.10.7	References	83
3.11	Tobacco Use	87
3.11.1	Introduction	87
3.11.2	Objectives	87
3.11.3	Variable Definitions	88
3.11.4	Findings.....	88
3.11.5	Discussion/Conclusion	90
3.11.6	Recommendations.....	90
3.11.7	References	90
3.12	Violence and Unintentional Injury	98
3.12.1	Introduction	98
3.12.2	Objectives	98
3.12.3	Variable Definitions	99
3.12.4	Findings.....	100
3.12.5	Discussion / Conclusion	101

3.12.6	Recommendations.....	102
3.12.7	References	103
APPENDICES	107

LIST OF TABLES

METHODOLOGY AND SAMPLING DESIGN

Table 2.1 Distribution of secondary schools sampled by state, NHMS 2017

GENERAL FINDINGS

Table 3.1.1 Response rate at student level by state, NHMS 2017

Table 3.1.2 Comparison between NHMS 2017 sample and total enrolment of students in Malaysia 2017

ALCOHOL CONSUMPTION

Table 3.2.1 Prevalence of ever alcohol drinkers among Form 1 to Form 5 students in Johor, 2017

Table 3.2.2 Prevalence of current alcohol drinkers among Form 1 to Form 5 students Johor, 2017

Table 3.2.3 Percentage of drunkenness among ever alcohol drinkers among Form 1 to Form 5 students in Johor, 2017

Table 3.2.4 Initiation age of alcohol drinking among ever alcohol drinkers among Form 1 to Form 5 students in Johor, 2017

Table 3.2.5 Sources of obtaining alcohol among current alcohol drinkers among Form 1 to Form 5 students in Johor, 2017

Table 3.2.6 Prevalence of social problems (got into trouble with family or friends, missed school, or got into fights) as a result of drinking alcohol among Form 1 to Form 5 students in Johor, 2017

DIETARY BEHAVIOURS

Table 3.3.1 Prevalence of students who had gone hungry most of the time or always because there was not enough food among Form 1 to Form 5 students in Johor, 2017

- Table 3.3.2** Prevalence of fruits intake of at least twice daily in the past 30 days among Form 1 to Form 5 students in Johor, 2017
- Table 3.3.3** Prevalence of vegetables intake of at least three times daily in the past 30 days among Form 1 to Form 5 students in Johor, 2017
- Table 3.3.4** Prevalence of fruits and vegetables intake at least five times daily in the past 30 days among Form 1 to Form 5 students in Johor, 2017
- Table 3.3.5** Prevalence of carbonated soft drinks consumption at least once a day in the past 30 days among Form 1 to Form 5 students in Johor, 2017
- Table 3.3.6** Prevalence of plain water intake 5 times or more per day in the past 30 days among Form 1 to Form 5 students in Johor, 2017
- Table 3.3.7** Prevalence of milk or milk products intake of at least two servings per day in the past 30 days among Form 1 to Form 5 students in Johor, 2017
- Table 3.3.8** Prevalence of consuming food from fast food restaurants of at least three days in the past seven days among Form 1 to Form 5 students in Johor, 2017

DRUG USE

- Table 3.4.1** Prevalence of ever used drugs in a lifetime among Form 1 to Form 5 students in Johor, 2017
- Table 3.4.2** Prevalence of current drug use in the past 30 days among Form 1 to Form 5 students in Johor, 2017
- Table 3.4.3** Prevalence of ever used marijuana in their lifetime among Form 1 to Form 5 students in Johor, 2017
- Table 3.4.4** Prevalence of current marijuana use in the past 30 days among Form 1 to Form 5 students in Johor, 2017
- Table 3.4.5** Prevalence of ever used amphetamines/metaphetamines in their lifetime among Form 1 to Form 5 students in Johor, 2017
- Table 3.4.6** Source of getting drugs in the past 30 days among current drug users, Form 1 to Form 5 students in Johor, 2017
- Table 3.4.7** Age of initiation of first use of drugs amongst ever drug users among Form 1 to Form 5 students in Johor, 2017

HYGIENE (INCLUDING ORAL HEALTH)

- Table 3.5.1** Prevalence of teeth brushing in the past 30 days among Form 1 to Form 5 students in Johor, 2017
- Table 3.5.2** Prevalence of use of fluoridated toothpaste among Form 1 to Form 5 students in Johor, 2017
- Table 3.5.3** Prevalence of use of dental floss among Form 1 to Form 5 students in Johor, 2017
- Table 3.5.4** Timing of last dental visit among Form 1 to Form 5 students in Johor, 2017
- Table 3.5.4.1** Prevalence of last dental visit within the last 24 months among Form 1 to Form 5 students in Johor, 2017
- Table 3.5.5** Prevalence of having missed class due to toothache in the past 12 months among Form 1 to Form 5 students in Johor, 2017
- Table 3.5.6** Prevalence of avoidance of smiling or laughing due to the appearance of their teeth among Form 1 to Form 5 students in Johor, 2017
- Table 3.5.7** Prevalence of use of soap during hand washing in the past 30 days among Form 1 to Form 5 students in Johor, 2017
- Table 3.5.8** Prevalence of handwashing before eating in the past 30 days among Form 1 to Form 5 students in Johor, 2017
- Table 3.5.9** Prevalence of handwashing after using the toilet in the past 30 days among Form 1 to Form 5 students in Johor, 2017

INTERNET USE AND ADDICTION

- Table 3.6.1** Prevalence of internet use in the past 30 days among Form 1 to Form 5 students in Johor, 2017
- Table 3.6.2** Prevalence of positive Malay Version Internet Addiction Test (MVIAT) among Form 1 to Form 5 students in Johor, 2017
- Table 3.6.3** Percentage of reported devices used by internet users among Form 1 to Form 5 students in Johor, 2017
- Table 3.6.4** Percentage of reported devices used by internet addicts among Form 1 to Form 5 students in Johor, 2017

MENTAL HEALTH PROBLEMS

- Table 3.7.1** Prevalence of loneliness "most of the time or always" in the past 12 months among Form 1 to Form 5 students in Johor, 2017
- Table 3.7.2** Prevalence of inability to sleep "most of the time or always" due to worry in the past 12 months among Form 1 to Form 5 students in Johor, 2017
- Table 3.7.3** Prevalence of suicidal ideation in the past 12 months among Form 1 to Form 5 students in Johor, 2017
- Table 3.7.4** Prevalence of suicidal plan in the past 12 months among Form 1 to Form 5 students in Johor, 2017
- Table 3.7.5** Prevalence of suicidal attempt "at least once" in the past 12 months among Form 1 to Form 5 students in Johor, 2017
- Table 3.7.6** Prevalence of not having any close friend among Form 1 to Form 5 students in Johor, 2017

PHYSICAL ACTIVITY

- Table 3.8.1** Prevalence of being physically active (60 minutes daily) for a total of at least 5 days or more among Form 1 to Form 5 students in Johor, 2017
- Table 3.8.2** Prevalence of spending at least 3 hours in sitting activities among Form 1 to Form 5 students in Johor, 2017

PROTECTIVE FACTORS

- Table 3.9.1** Prevalence of truancy in the past 30 days among Form 1 to Form 5 students in Johor, 2017
- Table 3.9.2** Prevalence of having peer support in the past 30 days among Form 1 to Form 5 students in Johor, 2017
- Table 3.9.3** Prevalence of having parental or guardian supervision in the past 30 days among Form 1 to Form 5 students Johor, 2017
- Table 3.9.4** Prevalence of having parental or guardian connectedness in the past 30 days among Form 1 to Form 5 students in Johor, 2017

Table 3.9.5 Prevalence of having parental or guardian bonding in the past 30 days among Form 1 to Form 5 students in Johor, 2017

Table 3.9.6 Prevalence of having parental or guardian respect for privacy in the past 30 days among Form 1 to Form 5 students in Johor, 2017

SEXUAL BEHAVIOURS

Table 3.10.1 Prevalence of ever had sexual intercourse among Form 1 to Form 5 students in Johor, 2017

Table 3.10.2 Prevalence of current sexual intercourse in the past 30 days among Form 1 to Form 5 students in Johor, 2017

Table 3.10.3 Percentage of first sexual intercourse before the age of 14 years among those who ever had sex among Form 1 to Form 5 students in Johor, 2017

Table 3.10.4 Percentage of having at least two sexual partners among students who ever had sex among Form 1 to Form 5 students in Johor, 2017

Table 3.10.5 Percentage of reported condom use during the last sexual intercourse among students who ever had sex among Form 1 to Form 5 students in Johor, 2017

Table 3.10.6 Percentage of reported using "other birth control methods" during the last sexual intercourse among students who ever had sex among Form 1 to Form 5 students in Johor, 2017

TOBACCO USE

Table 3.11.1 Prevalence of current smokers among Form 1 to Form 5 students in Johor, 2017

Table 3.11.2 Prevalence of current cigarette smokers among Form 1 to Form 5 students in Johor, 2017

Table 3.11.3 Prevalence of current any tobacco products use among Form 1 to Form 5 students in Johor, 2017

Table 3.11.4 Prevalence of current E-cigarettes/ Vape use among Form 1 to Form 5 students in Johor, 2017

- Table 3.11.5** Prevalence of current smoke and smokeless tobacco products use by types (except manufactured cigarette and E-cigarette/Vape) among Form 1 to Form 5 students in Johor, 2017
- Table 3.11.6** Prevalence of having tried a cigarette before the age of 14 years among ever smokers among Form 1 to Form 5 students in Johor, 2017
- Table 3.11.7** Prevalence of having tried E-cigarette/ Vape before the age of 14 years among ever E-cigarette/ Vape users among Form 1 to Form 5 students in Johor, 2017
- Table 3.11.8** Prevalence of quit smoking attempts in the past 12 months among those who ever smoked among Form 1 to Form 5 students in Johor, 2017
- Table 3.11.9** Prevalence of quit E-cigarette/ Vape attempts in the past 12 months among those who ever used E-Cigarette/ Vape among Form 1 to Form 5 students in Johor, 2017
- Table 3.11.10** Prevalence of being exposed to people smoking in their presence for at least one day in the past 7 days among Form 1 to Form 5 students in Johor, 2017
- Table 3.11.11** Prevalence of having father or mother or guardian who used any form of tobacco product among Form 1 to Form 5 students in Johor, 2017
- Table 3.11.12** Prevalence of having father or mother or guardian who used E-cigarette/ Vape among Form 1 to Form 5 students in Johor, 2017

VIOLENCE AND UNINTENTIONAL INJURY

- Table 3.12.1** Prevalence of involvement in violence at least once in the past 12 months among Form 1 to Form 5 students Johor, 2017
- Table 3.12.2** Prevalence of being bullied at least once in the past 30 days among Form 1 to Form 5 students in Johor, 2017
- Table 3.12.3** Most common ways of being bullied in the past 30 days among Form 1 to Form 5 students in Johor, 2017
- Table 3.12.4** Prevalence of being abused at least once in the past 30 days among Form 1 to Form 5 students in Johor, 2017
- Table 3.12.5** Prevalence of having had serious injury at least once in the past 12 months among Form 1 to Form 5 students in Johor, 2017

Table 3.12.6 Types of the most serious injury sustained in the past 12 months among students who were injured among Form 1 to Form 5 students in Johor, 2017

Table 3.12.7 Major cause of the most serious injury sustained in the past 12 months among students who were injured among Form 1 to Form 5 students in Johor, 2017

LIST OF APPENDICES

- Appendix 1 Members of Steering Committee NHMS 2015-2018
- Appendix 2 Terms of Reference for NHMS 2017 Steering Committee
- Appendix 3 List of members of Central Coordinating Committee
- Appendix 4 Terms of Reference for NHMS 2017 Central Coordinating Team
- Appendix 5 List of Research Team Members, NHMS 2017
- Appendix 6 List of Liaison Officer and Data Collection Teams
- Appendix 7 Questionnaire & OMR form

Executive Summary

The Adolescent Health Survey 2017 is aimed at assessing the prevalence of health risk behaviours and protective factors amongst secondary school students in Malaysia using self-administered anonymous questionnaires adapted from the Malaysian Global School-based Student Health Survey (GSHS) 2012. This study was conducted among secondary school-going adolescents between 26th March and 3rd May 2017. In order to ensure national representativeness, this study was implemented using a two-stage cluster sampling design. Out of 2738 secondary schools from the Ministry of Education and Ministry of Rural and Regional Development, 212 schools were selected and more than 30,000 students were eligible to participate in this survey. The overall response rate was 89.0% and 27,497 school-going adolescents completed the survey. A total of 14 secondary schools were randomly selected in Johor of which 1,731 students completed the survey out of 1,915 eligible respondents with response rate of 90.4%.

Johor Key Findings

The study observed that the prevalence of current cigarette smokers in Johor school-going adolescents was 9.2%. Amongst those who ever smoked cigarettes, 73.6% had tried cigarettes before the of age 14 years and 74.6% had attempted to quit smoking in the past 12 months. The prevalence of current e-cigarette/vape use in Johor school-going adolescents was 6.6%. Among those who ever smoked e-cigarette/vape, 46.7% had first tried e-cigarette/vape before age 14 years and 68.7% had tried to stop using e-cigarette/vape in the past 12 months. Overall, 10.5% of students reported having parents or guardians who used e-cigarette/vape, 42.8% of students reported having parents or guardians who used any form of tobacco products and 46.5% of students reported having been exposed to people who smoked in their presence in the past seven days. Prevalence of current alcohol drinkers among school-going adolescents in Johor was 7.2%. Among students who had ever consumed alcohol (17.3%), 16.6% reported ever drunkenness in their lifetime. With regards to drug use, 2.6% of students in Johor reported had ever used drugs and the prevalence of current drug users was 1.6%. Prevalence of ever using marijuana in their lifetime was 1.3% and current use of marijuana was 1.2%, while 0.9% of students reported had ever used amphetamines or methamphetamines during their lifetime.

The study found that prevalence of ever having sex among school-going adolescents in Johor was 5.7% and the prevalence of having had sex in the past 30 days was 4.2%. Of those who ever had sex, 27.9% of them had their first sexual experience before age 14 years, and 13.9% had at least two sexual partners. It was found that 5.4% of students reported they, or their partners had used condoms whilst 8.1% used other birth control methods. A total of 26.2% students had been seriously injured in the past 12 months. Among the respondents, 26.1% claimed to have been physically attacked in the past 12 months, while 21.3% of adolescents claimed to have been involved in physical fights. With regards to bullying, 13.9% reported to have ever been bullied in the past 30 days. Physical abuse at home was reported by 8.6% of students while 45.2% of reported verbal abuse at home in the past 30 days.

The overall prevalence of internet use among school-going adolescents in Johor was 93.0% and the prevalence of internet addiction was 33.9%. Smartphones were the most prevalent device used. A total of 8.9% school-going adolescents in Johor reported feeling lonely and 3.1% reported that they had no close friend. A total of 6.5% reported being unable to sleep “most of the time or always” due to worry in the 12 months prior to the survey. Prevalence of suicidal ideation, plan and attempt were 7.5%, 5.7%, and 4.6% respectively. Prevalence of truancy amongst students in the past 30 days was 24.2% while only 49.9% claimed to have peer support. Students who reported having parental or guardian supervision, parental or guardian connectedness, parental or guardian bonding and parental or guardian respect for privacy were 12.3%, 34.9%, 51.0% and 77.6% respectively.

Overall, 88.2% of students reported having cleaned or brushed their teeth at least twice daily while 0.5% of students reported that they had never cleaned or brushed their teeth in the past 30 days. A total of 54.7% students reported use of fluoridated toothpaste, while only 13.7% used dental floss. In the past 30 days, 9.5% never or rarely used soap when washing their hands, 4.4% never or rarely washed their hands before eating and 2.8% reported that they had never or rarely washed their hands after using the toilet. In relation to dietary behaviours, 3.6% reported being hungry most of the time or always in the past 30 days because there was not enough food at home. The consumption of fruits at least twice daily was 38.3% and consumption of vegetables at least thrice daily was 29.2% in the past 30 days. Only 15.8% of students reported fruits and vegetables consumption at least five times daily in the past 30

days. Consumption of carbonated soft drinks of at least once daily in the past 30 days was reported at 23.4% and 7.1% consumed food from fast food restaurants for at least three days in the past seven days. Milk/milk products consumption of at least two servings per day was reported at 27.2% while 59.3% drank plain water five times or more per day in the past 30 days. Prevalence of being physically active for a total of at least 60 minutes daily for five days or more in the past seven days was 25.3% whereas 60.7% of students had spent at least three hours in a typical or usual day in sitting activities.

Malaysia Key Findings

The study observed that the prevalence of current cigarette smokers in school-going adolescents was 13.8%. Amongst those who ever smoked cigarettes, 68.4% had tried cigarettes before the age of 14 years and 71.0% had attempted to quit smoking in the past 12 months. The prevalence of current e-cigarette/vape use in school-going adolescents was 9.8%. Among those who ever smoked e-cigarette/vape, 42.2% had first tried e-cigarette/vape before age 14 years and 60.9% had tried to stop using e-cigarette/vape in the past 12 months. Overall, 13.7% of students reported having parents or guardians who used e-cigarette/vape, 42.2% of students reported having parents or guardians who used any form of tobacco products and 42.0% of students reported having been exposed to people who smoked in their presence in the past 7 days. Prevalence of current alcohol drinkers among school-going adolescents was 10.2% in which alcohol sources were mainly from their own family or bought from stores. Among students who had ever consumed alcohol (19.3%), three quarter had their first alcoholic beverage before age 14 years and 31.9% reported ever drunkenness in their lifetime. With regards to drug use, 4.3% of students reported had ever used drugs and the prevalence of current drug users was 3.4%. Prevalence of ever using marijuana in their lifetime was 2.8% and current use of marijuana was 2.5%, while 2.4% of students reported had ever used amphetamines or methamphetamines during their lifetime.

The study found that prevalence of ever having sex among school-going adolescents was 7.3% and the prevalence of having had sex in the past 30 days was 5.4%. Of those who ever had sex, 31.9% of them had their first sexual experience before age 14 years, and 16.6% had at least two sexual partners. It was found that 12.7% of students reported they, or their partners had used condoms whilst 10.3% used other birth control methods. A total of 29.9% students had been seriously injured in the past 12 months with two most common causes of injury

being falls and motor vehicle accidents. Among respondents, 25.3% claimed to have been physically attacked in the past 12 months, while 24.9% of adolescents claimed to have been involved in physical fights. With regards to bullying, 16.2% reported to have ever been bullied in the past 30 days. The two most common forms of bullying were 'being made fun of because of how their body or face looks' and 'made fun of with sexual jokes, comments or gestures'. Physical abuse at home was reported by 11.8% of students while 43.2% reported verbal abuse at home in the past 30 days.

The overall prevalence of internet use among school-going adolescents was 85.6% and the prevalence of internet addiction was 29.0%. Smartphones were the most prevalent device used. A total of 9.3% secondary school students in Malaysia reported feeling lonely and 3.6% reported that they had no close friend. A total of 7.1% reported being unable to sleep "most of the time or always" due to worry in the 12 months prior to the survey. Prevalence of suicidal ideation, plan and attempt were 10.0%, 7.3%, and 6.9% respectively. Prevalence of truancy amongst students in the past 30 days was 29.4% and only 44.2% claimed to have peer support. Students who reported having parental or guardian supervision, parental or guardian connectedness, parental or guardian bonding and parental or guardian respect for privacy were 13.2%, 32.0%, 42.6%, and 74.4% respectively.

Overall, 87.1% of students reported having cleaned or brushed their teeth at least twice daily while 1.2% of students reported they had never cleaned or brushed their teeth in the past 30 days. A total of 58.3% students reported use of fluoridated toothpaste, while only 19.3% used dental floss. In the past 30 days, 11.6% never or rarely used soap when washing their hands, 6.1% never or rarely washed their hands before eating and 4.8% reported that they had never or rarely washed their hands after using the toilet. In relation to dietary behaviours, 3.9% reported being hungry most of the time or always in the past 30 days because there was not enough food at home. The consumption of fruits at least twice daily was 46.8% and consumption of vegetables at least thrice daily was 36.0% in the past 30 days. Only 23.5% reported fruits and vegetables consumption at least five times daily in the past 30 days. Consumption of carbonated soft drinks of at least once daily in the past 30 days was reported at 36.9% while 11.1% consumed food from fast food restaurants for at least three days in the past seven days. Milk/milk products consumption of at least two servings per day was

reported at 31.0% while 60.4% drank plain water five times or more per day in the past 30 days. Prevalence of being physically active for a total of at least 60 minutes daily for five days or more in the past seven days was 19.8% whereas 50.1% of students had spent at least three hours in a typical or usual day in sitting activities.

Recommendations:

In view of the above findings, the following recommendations are suggested:

- To strengthen awareness, knowledge and practice of positive health-related behaviours through home, school and community settings.
- To develop and disseminate more health education materials on health risk behaviours, its consequences and preventive measures.
- To enhance resilience and coping skills among students through school and community programs and activities such as, Doktor Muda, Minda Sihat, cadets and volunteerism.
- To strengthen protective factors against risky behaviours through intersectoral collaboration with more emphasis on spiritual values and parenting skills.
- Intersectoral collaboration to tackle the social determinants contributing to the adolescent health morbidities and mortalities.
- To evaluate the effectiveness of adolescent health programs provided by various agencies at a regular interval.

1.0 INTRODUCTION

The World Health Organization (WHO) has defined adolescents as a group of population between the ages of 10 to 19 years.¹ Adolescence sub-categories used in Malaysia consists of early adolescence (10-14 years), middle adolescence (15-17 years) and late adolescence (18-19 years).² Total population of adolescents in Malaysia is around 18% out of 31 million population.³ Adolescents are the most valuable asset in the country as they will become future leaders who will continue to sustain our national agendas.

By definition, adolescence is a period of transition from childhood to adulthood where significant changes occur in the form of physical appearance as well as emotional well-being. Rapid biological maturity precedes psychosocial maturity, thus having an impact on health consequences.⁴ Generally, they are perceived as the healthiest population and often overlooked until now. However previous studies had observed multiple morbidities among adolescents resulting from unintentional injuries, risky behaviours such as smoking, use of alcohol and drugs and also involvement in sexual activity.⁵ The current trend of the cyber era, in which more adolescents spend too much time “on line” and have become addicted to the internet, results in detachment from the real world and difficulties adapting with real world communication, which is later associated with mental health problems among adolescents.⁶ Physical inactivity and unhealthy dietary behaviors are associated with obesity and these behaviours may continue until adulthood. The practice of good personal hygiene care and dietary behaviours are equally important aspects that should not be put aside in assessing adolescent health.⁵ Their interaction with the environment also shapes adolescent growth through psychosocial experiences where peer and parental support play an important role. Adolescents are at-risk of premature morbidity and mortality if no preventive measures are taken.⁷ This population should enjoy the highest attainable standards of health with a supportive environment.

1.1 Research in Adolescent Health in Malaysian Context

In Malaysia, the Adolescent Health Unit has been established in 1995 under the Family Health Development Division, Ministry of Health, Malaysia.³ In terms of research activities, a nationwide health risk behavior study among adolescents was conducted in 1996 with four

main scopes namely smoking, alcohol consumption, drug use and sexual activity practices.⁸ Following that, the Global Youth Tobacco Surveys were conducted in 2003 and 2009 to identify tobacco consumption among youth.^{9,10} In 2010, the Institute for Health Behavioural Research had initiated The Youth Behaviour Risk Factor Surveillance (YBRFS), however the respondents only consisted of students from Forms 1, Form 2 and Form 4.¹¹ In realizing that the national data on health risks and behaviours are very much important in developing policy and programs for adolescents, the Ministry of Health, Malaysia took a step forward in collaborating with the World Health Organization (WHO) in conducting the Global School-based Student Health Survey (GSHS) Malaysia in 2012. The survey used a self-administered questionnaire assessing 10 main scopes of adolescent health risk behaviours primarily among students aged 13 to 17 years. The GSHS questionnaire was developed by WHO and the Center for Disease Control and Prevention (CDC) in collaboration with UNICEF, UNESCO, and UNAIDS in 2001. It has been used as a standard tool by more than 100 countries in the world. Local adaption and validation of the questionnaire was done before running the actual nationwide survey.¹²

The purpose of the GSHS was to provide accurate data on health behaviours and protective factors among students to:

- Help countries develop priorities, establish programmes, and advocate for resources for school health and youth health programmes and policies;
- Establish trends in the prevalence of health behaviours and protective factors for use in evaluation of school health and youth health promotion; and
- Allow countries, international agencies, and others to make comparisons across countries and within countries regarding the prevalence of health risk behaviours and protective factors.

The 10 main scopes were:

- Alcohol use
- Dietary behaviours
- Drug use
- Hygiene (including oral health)
- Mental health problems

- Physical activity
- Protective factors
- Sexual behaviours that contribute to HIV infection, other STI, and unintended pregnancy
- Tobacco use
- Violence and unintentional injury

With regards to the importance of serial data in measuring the trends of health risk behaviours among adolescents in Malaysia, the Ministry of Health has listed the Adolescent Health Survey as one of the main scopes in the National Health and Morbidity Survey (NHMS) cycle. The current Adolescent Health Survey used a similar methodology and validated Malaysian GSHS questionnaire in 2012 with the addition of the Malay Version Internet Addiction Test (MVIAT).¹³

References

1. WHO | Adolescent health [Internet]. WHO. [cited 2013 Sep 27]. Available from: http://www.who.int/topics/adolescent_health/en/index.html
2. Ministry of Health Malaysia. Engaging The Adolescent Module using HEADSS Framework. Kuala Lumpur, Malaysia; 2009.
3. Ministry of Health Malaysia. National Adolescent Health Plan of Action [Internet]. Ministry of Health Malaysia; 2015 [cited 2017 Oct 30]. Available from: <http://fh.moh.gov.my/v3/index.php/component/jdownloads/send/21-sektor-kesihatan-remaja/345-nahpoa-2015-2020>
4. WHO | Adolescent development [Internet]. WHO. [cited 2017 Jun 30]. Available from: http://www.who.int/maternal_child_adolescent/topics/adolescence/development/en/
5. WHO | Adolescents: health risks and solutions [Internet]. [cited 2016 Oct 23]. Available from: <http://www.who.int/mediacentre/factsheets/fs345/en/>
6. Gholamian B, Shahnazi H, Hassanzadeh A. The Prevalence of Internet Addiction and its Association with Depression, Anxiety, and Stress, among High-School Students. *Int J Pediatr.* 2017;5(4):4763–4770.
7. Sawyer SM, Afifi RA, Bearinger LH, Blakemore S-J, Dick B, Ezeh AC, et al. Adolescence: a foundation for future health. *The Lancet.* 2012;379(9826):1630–1640.

8. Institute for Public Health. National Health and Morbidity Survey 1996. Vol 2: General Findings. Kuala Lumpur: Ministry of Health Malaysia; 1997.
9. Ministry of Health Malaysia. Report of the Global Tobacco Survey, Malaysia. Kuala Lumpur, Malaysia: Ministry of Health Malaysia; 2009.
10. Manimaran K. Report of the Global Youth Tobacco Survey, Malaysia. Kuala Lumpur, Malaysia: Ministry of Health Malaysia; 2003.
11. Institute For Health Behavioural Research (IHBR). The Youth Behaviour Risk Factor Surveillance (YBRFSS) 2010. Kuala Lumpur: Ministry of Health Malaysia; 2011.
12. Institute for Public Health (IPH). The National Health and Morbidity Survey: Malaysia Global School-based Student Health Survey 2012. Kuala Lumpur: Ministry of Health Malaysia; 2012.
13. Chong Guan N, Isa SM, Hashim AH, Pillai SK, Harbajan Singh MK. Validity of the Malay Version of the Internet Addiction Test: A Study on a Group of Medical Students in Malaysia. *Asia Pac J Public Health*. 2012;27(2):NP2210-NP2219.

1.2 Objectives

1.2.1 General Objectives

To assess the prevalence of health risk behaviours among adolescents in Malaysia in order to review health priorities, program strategies and activities and planning for allocation of resources for adolescent health.

1.2.2 Specific Objectives

1.2.2.1 To identify the prevalence of alcohol use

1.2.2.2 To identify the dietary behaviors

1.2.2.3 To identify the prevalence of drug use

1.2.2.4 To identify the practice of hygiene including oral health

1.2.2.5 To identify the prevalence of internet use and addiction

1.2.2.6 To identify the prevalence of mental health problems

1.2.2.7 To identify the practice of physical activity

1.2.2.8 To identify the prevalence of protective factors

1.2.2.9 To identify the prevalence of sexual behaviours

1.2.2.10 To identify the prevalence of tobacco use

1.2.2.11 To identify the prevalence of violence and unintentional injury

1.2.3 The NHMS 2017 Organisation Team

The organisation of NHMS 2017 was set up at various levels of the Ministry of Health and Ministry of Education in order to conduct this survey.

1.2.3.1 NHMS Steering Committee

The NHMS Steering Committee, chaired by the Director-General of Health was set up at the national level to approve scopes of the NHMS 2015-2018 and to facilitate implementation of the survey.

The members and terms of reference of this committee are shown in **Appendix 1 and 2**.

1.2.3.2 Central Coordinating Team (CCT)

A working committee within the Institute for Public Health was established to coordinate implementation of the survey according to the scheduled Gantt chart. The Operation Centre was set up and led by the CCT team for coordinating and monitoring progress of the survey.

The list of CCT members and terms of reference are shown in **Appendix 3 and 4**. Figure 1 detailed the organisation chart at the Institute for Public Health level. Adolescent Health Survey was part of NHMS 2017 using the sample from secondary schools only.

1.2.3.3 Research Team Members

Research team members for each sub-scope were established and headed by a key-person (among IPH officers) together with the relevant stakeholders. Research team members were responsible for the technical input in development of the questionnaire manual, variable definition, data analysis and writing of the final report.

The list of members for each research teams are shown in **Appendix 5**.

1.2.3.4 State Liason Officers and Data Collection Team.

A State Liaison Officer was appointed in each State to facilitate planning and implementation of data collection within the States.

The list of State Liaison Officers and Data Collection Teams are shown in **Appendix 6**.




Figure 1: Organisation chart for data collection teams NHMS 2017

2.0 METHODOLOGY AND SAMPLING DESIGN

2.1 Target Population

The target population for the Adolescent Health Survey 2017 was school-going adolescents aged between 13 to 17 years in Malaysia. An equal sampling proportion was calculated from 13 States and three Federal Territories to represent adolescents in each State / Federal Territories.

2.2 Sampling Frame

The sampling frame used in this survey was the list of secondary schools from the Ministry of Education and Ministry of Rural and Regional Development. Students' enrolment data of Form 1 to Form 5 from 2,738 secondary schools in 2016 were used.

2.3 Sample Size Determination

Sample size was calculated by using a single proportion formula for estimation of prevalence.

The sample size calculation was based on a few criteria as stated below:

1. Variance of proportion of the variable of interest (Based on Global School-Based Student Health Survey 2012)
2. Margin of error (e) (Between 0.01 to 0.05)
3. Confidence Interval of 95%

To obtain an optimum sample size, a few adjustments were made to the sample size calculation as follows:

1. Adjusted n (srs) for total number of target population (N) (based on the population size for school-going adolescents in 2016)

$$n \geq \frac{n_{SRS}}{1 + \frac{n_{SRS}}{N}}$$

2. Adjusted for design effect (deff) (based on previous survey: GSHS 2012), n (complex) = n * deff
3. Adjusted for n(complex) taking into account expected non-response rate of 25%, n (adj) = n (complex) * (1 + non-response rate)

4. The sample size was adjusted according to the needs of analysis; at the national or state level.

Based on the requirements to achieve the objectives of the survey and above mention considerations, the optimum sample size required was 30, 496 respondents (**Table 2.1**).

Table 2.1: Distribution of Secondary Schools Sampled by State, NHMS 2017

No.	State / Federal Territories	Total Number of Schools	Number of Schools Sampled	Number of students sampled
1	Johor	306	14	1,906
2	Kedah	219	14	1,906
3	Kelantan	193	14	1,906
4	Melaka	87	14	1,906
5	Negeri Sembilan	139	14	1,906
6	Pahang	209	14	1,906
7	Pulau Pinang	141	14	1,906
8	Perak	275	14	1,906
9	Perlis	33	14	1,906
10	Selangor	345	14	1,906
11	Terengganu	162	14	1,906
12	Sabah	246	14	1,906
13	Sarawak	220	14	1,906
14	WP Kuala Lumpur	133	14	1,906
15	WP Labuan	19	8	1,906
16	WP Putrajaya	11	8	1,906
Total		2,738	212	30,496

2.4 Sampling Design

This survey utilised a two-stage stratified cluster sampling design to ensure representativeness of students from Form 1 to Form 5 (13-17 years). Malaysia was stratified into 16 states/ Federal Territories. The first stage of sampling was selection of secondary schools (schools with students from Forms 1 to Form 5). Schools were selected randomly with probability proportionate to school enrolment size. A total of 212 secondary schools were selected to participate in this survey. The second stage of sampling was selection of classes. All classes in each selected school were included in the sampling frame. Systematic random sampling was used to select minimum of four to maximum of 10 classes from each selected school to meet the required sample for each school in 13 states and WP Kuala Lumpur, WP Putrajaya and WP Labuan; 137 and 239 respondents respectively. All students in selected classes were eligible to participate in the survey.

2.5 Ethical Approval and Consent Forms

This study had obtained approvals from the Medical Research and Ethics Committee of the Ministry of Health and Ministry of Education. We obtained permission to carry out the survey from relevant offices of the Ministry of Education at state and district levels as well as from schools selected. Prior to the survey, several meetings with relevant Ministry of Education officers and persons in-charge at selected schools were conducted to ensure readiness in logistic preparation. Teachers were briefed on the parent's consent form, who then distributed it to parents a week prior to the survey. During the actual day of the survey, student's consent was obtained from eligible respondents before survey was conducted. Students who did not receive parental consent or they themselves refused to participate were considered as non-response of eligible participants in this survey.

2.6 Survey Instruments

A validated self-administered bilingual questionnaire adopted from the Malaysian GSHS 2012 with computer-scan-able answer sheets was used. Answer sheets were anonymous to ensure student confidentiality. Majority of the students completed the survey within two teaching periods. The Adolescent Health Questionnaires contained 77 questions addressing the following topics:

- Alcohol use
- Dietary behaviours
- Drug use
- Hygiene (including oral health)
- Internet Use and Addiction
- Mental health
- Physical activity
- Protective factors
- Sexual behaviours that contribute to HIV infection, other STIs, and unintended pregnancy
- Tobacco use
- Violence and unintentional injury

2.7 Data Collection

There were 36 teams to collect data; 4 teams each for Sabah and Sarawak and 2 teams per state in the Peninsular Malaysia and Federal Territories. Each team consisted of a field supervisor, research assistants and a driver. The field supervisor was a permanent staff in the Ministry of Health.

A pilot study was carried out on 6 February 2017. Data collection training was conducted in Kuala Lumpur and Kuching, Sarawak for data collection teams in Peninsular Malaysia, and for Sabah, Sarawak and WP Labuan respectively. The training module comprised of questionnaires and role-plays in handling the survey in classrooms or in a school hall. Data collection was from 26 March to 3 May 2017.

2.8 Data Management

Quality check on data was conducted throughout the survey based on specific identification (ID) numbers; from the state ID until individual student ID (generated for the study). Upon completing the survey, each student placed his / her answer sheet in an envelope. All answer sheets from the same school were wrapped together to make a bundle. These bundles were

collected by assigned drivers for schools in Peninsular Malaysia or using tracked postage for schools in Sabah, Sarawak and WP Labuan.

2.8.1 Data Operation Centre

An operation centre with several stations was set up to receive data “bundles” from the field:

Station 1: Respondent ID checking

Station 2: Scanning

Station 3: Verification

Station 4: Storage




Figure 2 : Work Flow of NHMS 2017

2.8.2 Data Analysis

Analysis was prepared according to objectives of the survey, working definitions and dummy tables. IBM SPSS Statistics for Windows, Version 21.0. (Armonk, NY: IBM Corp.) was used to import raw datasets in Excel form. Data set was checked and cleaned for any inconsistencies. The final data analysis was conducted by using complex sampling design and 95% confidence interval.

A weighting factor was applied to each student record to adjust for non-response and for varying probabilities of selection. Weight estimation was calculated by the following formula:

$$W = W1 \times W2 \times F \times PS$$

Where;

W1 = the inverse of probability of selecting the school

W2 = the inverse of probability of selecting the class within the school

F = the inverse of a school, class and student level non-response adjustment factor

PS = a post stratification adjustment factor calculated by class and gender

The weighted results were used to make important inferences on the health risk behaviours and protective factors of all students from Form 1 to Form 5. Variable definitions used in this survey were derived from the Malaysian GSHS 2012 and was discussed in detail for each scope.

3.0 FINDINGS

3.1 General Findings

A total of 27,497 OMR forms were received from 212 selected schools. The overall response rate was 89.2%. The response rates for schools and classes were 100% each, while the response rate for students was 89.2%. (**Table 3.1.1**).

Table 3.1.1: Response Rate at Student Level by State, NHMS 2017

State	Selected Schools	Eligible students	Completed OMR forms	Response rate (%)
Johor	14	1,915	1,731	90.4
Kedah	14	1,930	1,691	87.6
Kelantan	14	1,900	1,631	85.8
Melaka	14	1,986	1,872	94.3
Negeri Sembilan	14	1,930	1,718	89.0
Pahang	14	1,948	1,784	91.6
Pulau Pinang	14	1,974	1,749	88.6
Perak	14	1,931	1,754	90.8
Perlis	14	1,992	1,667	83.7
Selangor	14	1,840	1,671	90.8
Terengganu	14	1,880	1,669	88.8
Sabah	14	1,965	1,686	85.8
Sarawak	14	1,919	1,779	92.7
WP Kuala Lumpur	14	1,937	1,721	88.8
WP Labuan	8	1,907	1,712	89.8
WP Putrajaya	8	1,869	1,662	88.9
Total	212	30,823	27,497	89.2

Comparison of total estimated population (weighted) with the national secondary school enrolment is shown in **Table 3.1.2** Geographic information system (GIS) on the mapping of selected secondary schools is shown in **Figure 3**. The socio-demographic characteristics of the sample by gender consist of 43.2% (747) males and 56.8% (984) females in Johor . In terms of school locality; 72.0% (1,246) and 28.0% (485) were students studying in schools located in urban areas and rural areas respectively.

Table 3.1.2: Comparison between NHMS 2017 respondents and national enrolment of secondary school students in Malaysia 2017

State	NHMS 2017			National Enrolment 2017	
	Unweighted count	Estimated Enrolment (weighted)	Prevalence (%)	Secondary School Students	Prevalence (%)
Malaysia	27,497	2,146,447	100.0	2,146,509	100.0
State					
Johor	1,731	275,711	12.8	275,700	12.8
Kedah	1,691	154,645	7.2	154,643	7.2
Kelantan	1,631	121,684	5.7	121,683	5.7
Melaka	1,872	67,234	3.1	67,235	3.1
Negeri Sembilan	1,718	88,430	4.1	88,429	4.1
Pahang	1,784	103,630	4.8	103,644	4.8
Pulau Pinang	1,749	112,980	5.3	112,981	5.3
Perak	1,754	181,681	8.5	181,724	8.5
Perlis	1,667	27,012	1.3	27,014	1.3
Selangor	1,671	391,634	18.2	391,623	18.2
Terengganu	1,669	98,667	4.6	98,664	4.6
Sabah	1,686	198,960	9.3	199,006	9.3
Sarawak	1,779	197,888	9.2	197,876	9.2
WP Kuala Lumpur	1,721	112,376	5.2	112,370	5.2
WP Labuan	1,712	5,539	0.3	5,539	0.3
WP Putrajaya	1,662	8,376	0.4	8,378	0.4
Sex					
Male	13,135	1,064,954	49.6	1,064,988	49.6
Female	14,362	1,081,493	50.4	1,081,521	50.4
Form					
Form 1	5,704	451,017	21.0	451,024	21.0
Form 2	5,501	426,924	19.9	426,908	19.9
Form 3	5,837	431,050	20.1	431,043	20.1
Form 4	5,532	414,604	19.3	414,653	19.3
Form 5	4,923	422,852	19.7	422,881	19.7




Figure 3: GIS mapping of the selected secondary schools

3.2 Alcohol Consumption

Contributors: Tania Gayle Robert, Hamizatul Akmal Abd Hamid, Halizah Mat Rifin, Mohd Hatta Mutalip, Norli Abdul Jabbar, Rozanim Kamaruddin, Muhammad Fadhli Mohd Yusoff, Thamil Arasu Saminathan, Jane Ling Miaw Yn, Wan Shakira Rodzlan Hasani, Hasimah Ismail, Nur Liana Abdul Majid

3.2.1 Introduction

Globally, there are about 3.3 million deaths attributable to harmful use of alcohol annually.¹ The South East Asian region contributed to 4.6% of alcohol-attributable deaths and 4.0% burden of disease and injury.¹ People consuming alcohol are prone to more than 200 diseases and injuries as well as social consequences due to its harmful effects.¹ The Centre for Disease Control (CDC) estimates on average in the United States from 2006 to 2010, there were 4,358 alcohol-attributable deaths due to excessive alcohol use for those under 21; 1,580 deaths from motor vehicle crashes, 1,269 as a result of homicides, 492 from suicide, as well as 245 from other injuries such as falls, burns, and drowning.² Alarmingly, about 33% of teens (by age 15) have had at least 1 drink of an alcoholic beverage.³ Evidently, individuals who started drinking before the age of 15 are five times more likely to have alcohol-related problems later in life.⁴ The consequence of early alcohol drinking could lead to memory impairment, thus causing low educational achievement and high absenteeism rates.^{5,6}

3.2.2 Objectives

- i. To determine the prevalence of ever and current drinkers among students from Form 1 to Form 5.
- ii. To describe the socio-demographic characteristics of ever and current drinkers among students from Form 1 to Form 5.
- iii. To identify the age of alcohol drinking initiation among students from Form 1 to Form 5
- iv. To identify the sources of obtaining alcoholic beverages among students from Form 1 to Form 5.
- v. To identify the prevalence of drunkenness among students from Form 1 to Form 5 who consume alcohol.

- vi. To determine the frequency of social problems related to alcohol consumption among students from Form 1 to Form 5.

3.2.3 Variable definitions

- **Drinking alcohol:** A “drink” is a glass of wine, tuak, lihing, bahar, ijuk or toddy; a can of beer, a small glass of liquor’ or mixed drink. Drinking alcohol does not include drinking a few sips of wine for religious purposes.
- **Ever drinkers:** Those who had a history of alcohol consumption in their lifetime.
- **Current drinkers:** Those who had at least a “drink” of alcohol in the past 30 days.
- **Drunkenness:** When someone demonstrates signs such as staggering when walking, not being able to speak right and throwing up after consuming alcohol in a lifetime.
- **Social problems:** Having trouble with family or friends, missed school or got into fights as a result of drinking alcohol in a lifetime.

3.2.4 Findings

The prevalence of ever alcohol drinkers among students from Form 1 to Form 5 in Johor was 17.3% (95% CI: 11.62, 24.85). There was no significant difference noted across school locality, sex and form (**Table 3.2.1**). The prevalence of current alcohol drinkers among students from Form 1 to Form 5 was 7.2% (95% CI: 4.90, 10.30) (**Table 3.2.2**). Among ever alcohol drinkers, 16.6% (95% CI: 11.11, 24.13) reported drunkenness (**Table 3.2.3**) and majority had their first alcoholic beverage at the age of 12 or 13 years old [31.8% (95% CI: 25.74, 38.46)] (**Table 3.2.4**).

Alarmingly, among current alcohol drinkers, more than half (54.1%, 95% CI: 37.33, 70.03) of them obtained their alcoholic beverages from their families (**Table 3.2.5**). The prevalence of students getting into trouble with their family and friends, missed school or got into fights one or more times as a result of drinking alcohol was 2.9% (95% CI: 2.11, 4.04) and there was no significant difference by school locality and sex (**Table 3.2.6**).

3.2.5 Discussion/ Conclusion

The prevalence of current alcohol drinkers among students from Form 1 to Form 5 in Johor was 7.2% (95% CI: 4.90, 10.30) whereas the national prevalence was 10.2% (95% CI: 9.00, 11.60). The prevalence of drunkenness among ever alcohol drinkers in Johor was significantly lower [16.6% (95% CI: (11.11, 24.13))] compared to the national prevalence [31.9% (95% CI: 29.26, 34.70)]. The prevalence of having social problems because of drinking alcohol was also lower [2.9% (95% CI: 2.11, 4.04)] in Johor compared to the national prevalence [4.7% (95% CI: 4.12, 5.24)]. The most common source of obtaining alcoholic beverages among the students was from family; national prevalence [37.7% (95% CI: 33.26, 42.25)]; Johor [54.1% (95% CI: 37.33, 70.03)]. The age of initiation of alcohol drinking among the students was 12 or 13 years old, nationally [33.28% (95% CI: 31.02, 35.62)] and in Johor [31.8% (95% CI: 25.74, 38.46)].

3.2.6 Recommendations

Addressing alcohol drinking among adolescents in Malaysia is vital to prevent social and other harmful effects of alcohol on the health of adolescents. Mapping of localities with high alcohol drinking and drunkenness prevalence is highly recommended, since alcohol use is not rampant and localised in specific populations and locations.

By doing this, interventions can be focused on the identified groups. Among the interventions that can be implemented are:

1. To instill awareness about the danger of underage drinking to the adolescents as early as possible.
2. Family-based prevention programmes: Educating parents on the negative effects of alcohol especially on long-term memory and learning skills, health effects and its social repercussions on adolescent behaviour and subsequent dependence and abuse if not controlled. Parents should play an important role as the firsthand educators on the harmful effects of alcohol to their children and prohibiting children to initiate early alcohol drinking.
3. School-based prevention programmes: Empower counselors and teachers to screen their students for alcohol use and to start intervention for these students as soon as possible.

3.2.7 References

1. World Health Organization, Global status report on alcohol and health 2014. Geneva: World Health Organization
2. National Institute on Alcohol Abuse and Alcoholism (NIAAA), Underage Drinking. February 2017.
https://pubs.niaaa.nih.gov/publications/underagedrinking/Underage_Fact.pdf
3. National Institute on Alcohol Abuse and Alcoholism (NIAAA). Underage Drinking: Why Do Adolescents Drink, What are the Risks, and How Can Underage Drinking Be Prevented?. November 2006. Alcohol Alert. Number 67.
4. World Health Organization, Evidence for the Effectiveness and Cost-Effectiveness of Interventions to Reduce Alcohol-Related Harm, Copenhagen, Regional Office for Europe. 2009
5. Carson A. Alcohol and the developing adolescent brain: evidence review. *J R Coll Physicians Edinb.* 2015 Mar;45(1):12-4. doi: 10.4997/JRCPE.2015.103.
6. Mohd Ramlan Mohd Arshad, Munirah Omar, and Nurul Afiqah Shahdan, "Alcoholism among Youth: A Case Study in Kuala Lumpur, Malaysia," *International Journal of Culture and History* vol. 1, no. 1, pp. 21-28, 2015.

Table 3.2.1: Prevalence of ever alcohol drinkers among Form 1 to Form 5 students in Johor, 2017

Socio-demographic Characteristics	Unweighted Count	Estimated Population	Prevalence (%)	95% CI	
				Lower	Upper
JOHOR	303	47,562	17.3	11.62	24.85
Locality of school					
Urban	256	39,962	20.4	13.53	29.47
Rural	47	7,600	9.6	4.57	18.93
Sex					
Male	114	21,705	15.8	10.66	22.85
Female	189	25,857	18.7	10.87	30.16
Form					
Form 1	41	6,798	11.7	6.69	19.76
Form 2	59	7,766	14.6	8.47	23.93
Form 3	52	8,014	14.8	7.86	26.09
Form 4	78	11,054	20.5	12.65	31.35
Form 5	73	13,930	24.8	12.45	43.30

Table 3.2.2: Prevalence of current drinkers among Form 1 to Form 5 students in Johor, 2017

Socio-demographic Characteristics	Unweighted Count	Estimated Population	Prevalence (%)	95% CI	
				Lower	Upper
JOHOR	124	19,719	7.2	4.90	10.30
Locality of school					
Urban	103	16,315	8.3	5.39	12.62
Rural	21	3,403	4.3	1.97	9.07
Sex					
Male	58	10,543	7.7	4.80	12.09
Female	66	9,176	6.6	4.10	10.53
Form					
Form 1	18	3,081	5.3	2.33	11.67
Form 2	29	3,930	7.4	4.11	12.88
Form 3	23	3,778	7.0	3.53	13.30
Form 4	34	5,015	9.3	5.70	14.75
Form 5	20	3,913	7.0	3.10	14.89

Table 3.2.3: Percentage of drunkenness among ever alcohol drinkers among students Form 1 to Form 5, Johor 2017

Socio-demographic Characteristics	Unweighted Count	Estimated Population	Prevalence (%)	95% CI	
				Lower	Upper
JOHOR	48	7,876	16.6	11.11	24.13
Locality of school					
Urban	40	6,597	16.6	10.42	25.37
Rural	8	1,279	16.8	8.50	30.60
Sex					
Male	20	3,858	17.8	10.64	28.18
Female	28	4,018	15.7	9.30	25.16
Form					
Form 1	8	1,333	20.2	7.27	44.90
Form 2	8	1,125	14.5	6.43	29.49
Form 3	5	687	8.6	3.55	19.29
Form 4	12	1,772	16.0	8.14	29.13
Form 5	15	2,958	21.2	11.16	36.66

Table 3.2.4: Initiation age of alcohol drinking among ever alcohol drinkers among Form 1 to Form 5 students in Johor, 2017

	Unweighted Count	Estimated Population	Prevalence (%)	95% CI	
				Lower	Upper
7 years old or younger	26	4,151	8.7	5.43	13.73
8 or 9 years old	29	4,311	9.1	6.37	12.74
10 or 11 years old	49	7,656	16.1	10.84	23.24
12 or 13 years old	98	15,107	31.8	25.74	38.46
14 or 15 years old	62	9,887	20.8	16.75	25.50
16 years old or older	-	-	-	-	-
Unknown	39	6,450	13.6	7.64	22.94

Table 3.2.5: Sources of getting alcohol among current drinkers among Form 1 to Form 5 students in Johor, 2017

	Unweighted Count	Estimated Population	Prevalence (%)	95% CI	
				Lower	Upper
I bought in a store	21	3,479	21.7	13.76	32.62
I gave someone money to buy	3	558	3.5	1.27	9.19
I got it from my friends	13	2,050	12.8	6.42	23.96
I got it from my family	57	8,656	54.1	37.33	70.03
I stole it	1	173	1.1	0.16	6.84
I got it some other way	6	1,077	6.7	2.55	16.64

Table 3.2.6: Prevalence of social problems (got into trouble with family or friends, missed school, or got into fights) as a result of drinking alcohol among Form 1 to Form 5 students in Johor, 2017

Socio-demographic Characteristics	Unweighted Count	Estimated Population	Prevalence (%)	95% CI	
				Lower	Upper
JOHOR	50	8,055	2.9	2.11	4.04
Strata					
Urban	36	5,861	3.0	1.98	4.49
Rural	14	2,194	2.8	1.72	4.42
Sex					
Male	18	3,552	2.6	1.37	4.83
Female	32	4,503	3.3	2.42	4.35

3.3. Dietary Behaviours

Contributors: Rashidah Ambak, Syafinaz Mohd Sallehuddin, Norlida Zulkafly, Ruhaya Salleh, Cheong Siew Man, Mohamad Hasnan Ahmad, S. Maria Awaluddin, Rusidah Selamat

3.3.1 Introduction

Fruits and vegetables are good sources of complex carbohydrates, vitamins, minerals, and other substances important for adolescent's good health. Dietary patterns that include higher intakes of fruits and vegetables are associated with several health benefits, including a decreased risk for some types of cancer.

3.3.2 Objectives

To describe the prevalence of:

- i. Students who had gone hungry most of the time or always in the past 30 days.
- ii. Fruits intake of at least twice daily in the past 30 days.
- iii. Vegetables intake of at least three times daily in the past 30 days.
- iv. Fruits and vegetables intake of at least five times daily in the past 30 days.
- v. Carbonated drinks consumption of at least once daily in the past 30 days.
- vi. Plain water intake of five times or more daily in the past 30 days.
- vii. Milk or milk products intake of at least two times daily in the past 30 days.
- viii. Food consumption from fast food restaurant of at least three days in the past seven days

3.3.3 Variable Definitions

- **Gone hungry:** students who had gone hungry most of the time or always because there was not enough food at home for the past 30 days or living without financial means to access enough food for active and healthy living.
- **Fruits intake:** fruits intake of at least twice daily in the past 30 days, inclusive of various types of local fruit, seasonal fruit and imported fruit.
- **Vegetables intake:** vegetables intake of at least three times daily in the past 30 days, either eaten raw or cooked.

- **Plain water intake:** Includes mineral water, boiled water, or tap water
- **Carbonated drinks intake:** carbonated drinks consumption of at least once daily in the past 30 days (except mentioned as diet soft drinks).
- **Dairy products intake:** milk or milk products intake at least two times daily in the past 30 days.
- **Fast food intake:** consuming food from fast food restaurants at least three days in the past seven days

3.3.4 Findings

Hunger

The overall prevalence of students who reported being hungry most of the time or always in the past 30 days was 3.6% (95% CI: 2.56, 5.01) with an estimated projection to 9,898 school-going adolescents. In terms of school locality, there was no significant difference reported between students studying in the urban areas, 3.3% (95% CI: 2.06, 5.14) and students from the rural areas, 4.4% (95% CI: 3.04, 6.32). There were 2.8% (95% CI: 1.60, 4.77) of male students and 4.4% (95% CI: 2.82, 6.81) female students who reported being hungry. Form 5 students reported the highest prevalence of being hungry [5.7% (95% CI: 3.00, 10.40)] as compared with other forms (**Table 3.3.1**).

Fruits intake

The overall prevalence of consuming fruits at least twice daily in the past 30 days was 38.2% (95%CI: 34.11, 42.47) with an estimated projection to 105,328 school-going adolescents. In terms of school locality, urban school-going adolescents significantly reported lower prevalence of fruits intake (35.8%; 95% CI: 30.99, 40.90) as compared to the students in the rural areas (44.2%; 95% CI: 42.99, 45.34). There were 39.3% (95% CI: 34.13, 44.79) male students and 37.1% (95% CI: 31.34, 43.22) female students who reported consuming fruits at least twice daily. By form, there was a decreasing trend of consuming fruits at least twice daily in the upper forms (**Table 3.3.2**).

Vegetables intake

The overall prevalence of consuming vegetables at least three times daily in the past 30 days was 29.2% (95%CI: 27.04, 31.55) with an estimated projection of 80,627 school-going adolescents. In terms of school locality, urban school-going adolescents reported a lower

prevalence of vegetables intake (28.3%; 95% CI: 26.07, 30.62) as compared to the students in rural areas (31.6%; 95% CI: 27.51, 35.99). There were 30.2% (95% CI: 26.74, 33.80) male students and 28.3% (95% CI: 25.49, 31.38) female students who reported consuming vegetables at least twice daily. By form, there was no significant difference in consuming vegetables at least twice daily amongst the forms (**Table 3.3.3**).

Fruits and vegetables intake at least five times daily

The overall prevalence of consuming fruits and vegetables at least five times daily in the past 30 days was 15.8% (95% CI: 14.06, 17.81) with an estimated projection of 43,681 school-going adolescents. There was no difference in fruits and vegetables intake at least five times daily by school locality and sex. Form 1 students reported the highest prevalence [21.0% (95% CI: 16.94, 25.72)] of consuming fruits and vegetables at least five times daily as compared to other forms (**Table 3.3.4**).

Carbonated soft drinks intake

The overall prevalence of consuming carbonated drinks at least once daily in the past 30 days was 23.4% (95%CI: 18.04, 29.84) with an estimated projection to 64,528 school-going adolescents. In terms of school locality, the urban school-going adolescents who reported carbonated drinks consumption at least once daily was 22.2% (95% CI: 15.14, 31.24) as compared to the students in rural areas [26.6% (95% CI: 23.60, 29.73)]. The male students reported significantly higher prevalence of carbonated drinks consumption [29.5% (95% CI: 22.99, 37.03)] as compared to the females [17.4% (95% CI: 13.11, 22.67)]. Form 1 students, 28.5% (95% CI: 21.99, 35.98) and Form 2 students, 28.1% (95% CI: 19.43, 38.83) reported the highest prevalence of carbonated drinks consumption as compared to other forms (**Table 3.3.5**).

Plain water intake

The overall prevalence of drinking plain water five times or more daily in the past 30 days was 59.3% (95%CI: 54.09, 64.29) with an estimated projection to 162,589 school-going adolescents. In terms of school locality, the urban school-going adolescents reported higher prevalence of 61.2% (95% CI: 55.02, 66.95) as compared to the students in rural areas [54.7 % (95% CI: 46.32, 62.82)]. Male students reported lower prevalence of 55.3 % (95% CI: 49.42, 60.96) as compared to females [63.3 % (95% CI: 56.53, 69.52)]. Form 2 students reported the

lowest prevalence of drinking plain water five times or more daily [51.9% (95% CI: 42.94, 60.81)] as compared to other forms (**Table 3.3.6**).

Milk and milk products intake

The overall prevalence of milk or milk products intake of at least two times daily in the past 30 days was 27.2% (95%CI: 23.76, 30.83) with an estimated projection to 74,744 school-going adolescents. There was no significant difference in taking milk or milk products by sex and school locality. Form 4 students [20.5% (95%CI: 15.51, 26.60)] showed the lowest prevalence in taking milk or milk products compared to other forms (**Table 3.3.7**).

Fast food intake

The overall prevalence of consuming food from fast food restaurants of at least three days in the past seven days was 7.1% (95%CI: 5.18, 9.73) with an estimated projection to 19,631 school-going adolescents. In terms of school locality, urban school-going adolescents reported lower prevalence of 7.0% (95% CI: 4.46, 10.81) as compared to the students in rural areas at 7.5% (95% CI: 6.11, 9.06). Male students reported higher prevalence of 7.8 % (95% CI: 5.50, 10.91) as compared to the females at 6.5% (95% CI: 4.45, 9.35). By form, there was an increasing trend of consuming food from fast food restaurants of at least three days in the past seven days in upper forms (**Table 3.3.8**).

3.3.5 Discussion/ Conclusion

The prevalence of being hungry most of the time or always during the past 30 days in Johor was 3.6% (95% CI: 2.56, 5.01) which showed no significant difference with the national prevalence [3.9% (95% CI: 3.53, 4.24)] but decreased from the previous Johor GSHS 2012 [5.7% (95% CI: 4.57, 6.99)]¹. Students in Johor reported lower prevalence of consuming fruits at least twice daily in 2017 [38.2% (95% CI: 34.11, 42.47)] compared to Johor GSHS 2012 [41.0% (95% CI: 36.73, 45.36)]¹ and national prevalence [46.8% (95% CI: 45.09, 48.53)] in 2017. For vegetables intake at least three times per day [29.2% (95% CI: 27.04, 31.55)], the prevalence was lower than Johor GSHS 2012 [32.3% (95% CI: 29.17, 35.59)]¹, however it was significantly lower in comparison with the national prevalence [36.0% (95% CI: 34.45, 37.52)]. Students in Johor [15.8% (95% CI: 14.06, 17.81)] reported a significantly lower fruits and vegetables consumption at least five times daily compared to Johor GSHS 2012 [29.8% (95% CI: 26.51, 33.37)], and also significantly lower than the national prevalence [23.5% (95% CI:

22.20, 24.94)]. For consumption of carbonated drink once daily [23.4% (95% CI: 18.04, 29.84)], it has decreased from the prevalence of Johor GSHS 2012 [31.4% (95% CI: 26.09, 37.16)]¹ but lower when compared to the national prevalence [36.9% (95% CI: 35.04, 38.79)]. Consumption of fast food at least three days in the past seven days [7.14% (95% CI: 5.18, 9.73)] was similar to the prevalence of Johor GSHS 2012 [7.3% (95% CI: 5.05, 10.40)]¹ and national prevalence [11.1% (95% CI: 10.24, 12.01)]. Consumption of plain water intake five times or more daily [59.3% (95% CI: 54.09, 64.29)] was similar to the prevalence of Johor GSHS 2012 [56.1% (95% CI: 52.29, 59.86)]¹ and national prevalence [60.4% (95% CI: 59.07, 61.78)].

3.3.6 Recommendations

In view of the findings highlighted above, Malaysian students need more attention towards a healthy eating habit. Parents play an important role to ensure that healthy and nutritious food such as fruits, yogurt, and high fibre snacks are available on shelves and in fridges at home. Appropriate nutrition education program with creative and innovative approaches should be carried out in promoting healthy diet, specifically eating more fruits and vegetables, drinking more plain water, consuming more milk/milk products and reducing fast food intake and carbonated soft drinks intake. Further research should be conducted to identify underlying causes of food insecurity and unhealthy eating habits among students in Malaysia.

3.3.7 References

1. Institute for Public Health (IPH). 2012. National Health and Morbidity Survey, NHMS 2012: Johor Global School-based Student Health Survey 2012. Kuala Lumpur Ministry of Health Malaysia.

Table 3.3.1: Prevalence of students who had gone hungry most of the time or always because there was not enough food among Form 1 to Form 5 students in Johor, 2017

Socio-demographic characteristics	Most of the time or always				
	Unweighted Count	Estimated Population	Prevalence (%)	95% CI	
				Lower	Upper
JOHOR	62	9,898	3.6	2.56	5.01
Locality of school					
Urban	41	6,408	3.3	2.06	5.14
Rural	21	3,490	4.4	3.04	6.32
Sex					
Male	21	3,798	2.8	1.60	4.77
Female	41	6,100	4.4	2.82	6.81
School level					
Lower secondary	33	5,056	3.1	1.76	5.25
Upper secondary	29	4,842	4.4	2.77	6.91
Form					
Form 1	8	1,272	2.2	0.93	5.11
Form 2	13	1,715	3.2	1.35	7.47
Form 3	12	2,069	3.8	1.67	8.50
Form 4	12	1,662	3.1	1.51	6.19
Form 5	17	3,179	5.7	3.00	10.40

Table 3.3.2: Prevalence of fruits intake of at least twice daily in the past 30 days among Form 1 to Form 5 students in Johor, 2017

Socio-demographic characteristics	Yes (at least twice daily)				
	Unweighted Count	Estimated Population	Prevalence (%)	95% CI	
				Lower	Upper
JOHOR	659	105,328	38.2	34.11	42.47
Locality of school					
Urban	446	70,233	35.8	30.99	40.90
Rural	213	35,094	44.2	42.99	45.34
Sex					
Male	299	53,942	39.3	34.13	44.79
Female	360	51,385	37.1	31.34	43.22
School level					
Lower secondary	462	71,925	43.5	37.70	49.41
Upper secondary	197	33,402	30.3	25.60	35.45
Form					
Form 1	168	28,586	49.3	42.27	56.39
Form 2	171	23,344	43.8	33.59	54.56
Form 3	123	19,996	36.9	30.21	44.10
Form 4	111	16,971	31.4	24.32	39.48
Form 5	86	16,431	29.2	20.80	39.39

Table 3.3.3: Prevalence of vegetables intake of at least three times daily in the past 30 days among Form 1 to Form 5 students in Johor, 2017

Socio-demographic characteristics	Yes (at least three times daily)				
	Unweighted Count	Estimated Population	Prevalence (%)	95% CI	
				Lower	Upper
JOHOR	502	80,627	29.2	27.04	31.55
Locality of school					
Urban	348	55,517	28.3	26.07	30.62
Rural	154	25,110	31.6	27.51	35.99
Sex					
Male	225	41,352	30.2	26.74	33.80
Female	277	39,275	28.3	25.49	31.38
School level					
Lower secondary	336	51,570	31.2	28.29	34.19
Upper secondary	166	29,057	26.4	22.54	30.57
Form					
Form 1	114	19,236	33.2	28.97	37.69
Form 2	124	16,613	31.2	25.39	37.61
Form 3	98	15,722	29.0	24.60	33.83
Form 4	87	13,154	24.3	19.66	29.73
Form 5	79	15,902	28.3	23.09	34.15

Table 3.3.4: Prevalence of fruits and vegetables intake at least five times daily in the past 30 days among Form 1 to Form 5 students in Johor, 2017

Socio-demographic characteristics	meet 2 serving of fruits and 3 serving of vegetables				
	Unweighted Count	Estimated Population	Prevalence (%)	95% CI	
				Lower	Upper
JOHOR	276	43,681	15.8	14.06	17.81
Locality of school					
Urban	187	29,313	14.9	13.00	17.07
Rural	89	14,367	18.1	15.52	20.96
Sex					
Male	121	21,657	15.8	13.39	18.53
Female	155	22,023	15.9	13.70	18.36
School level					
Lower secondary	203	31,274	18.9	15.92	22.29
Upper secondary	73	12,406	11.3	8.97	14.03
Form					
Form 1	72	12,169	21.0	16.94	25.72
Form 2	74	9,961	18.7	13.98	24.53
Form 3	57	9,144	16.9	13.20	21.30
Form 4	40	6,139	11.4	8.12	15.68
Form 5	33	6,268	11.2	7.25	16.78

Table 3.3.5: Prevalence of carbonated soft drinks consumption at least once a day in the past 30 days among Form 1 to Form 5 students in Johor, 2017

Socio-demographic characteristics	Yes (at least once a day)				
	Unweighted Count	Estimated Population	Prevalence (%)	95% CI Lower Upper	
JOHOR	388	64,528	23.4	18.04	29.84
Locality of school					
Urban	263	43,427	22.2	15.14	31.24
Rural	125	21,102	26.6	23.60	29.73
Sex					
Male	219	40,501	29.5	22.99	37.03
Female	169	24,027	17.4	13.11	22.67
School level					
Lower secondary	261	41,024	24.8	19.34	31.29
Upper secondary	127	23,504	21.3	14.72	29.85
Form					
Form 1	94	16,423	28.5	21.99	35.98
Form 2	110	14,989	28.1	19.43	38.83
Form 3	57	9,612	17.7	13.13	23.51
Form 4	78	12,654	23.4	16.03	32.88
Form 5	49	10,851	19.3	11.51	30.55

Table 3.3.6: Prevalence of plain water intake 5 times or more per day in the past 30 days among Form 1 to Form 5 students in Johor, 2017

Socio-demographic characteristics	Yes (at least 5 times daily)				
	Unweighted Count	Estimated Population	Prevalence (%)	95% CI Lower Upper	
JOHOR	1027	162,589	59.3	54.09	64.29
Locality of school					
Urban	760	119,212	61.2	55.02	66.95
Rural	267	43,378	54.7	46.32	62.82
Sex					
Male	406	75,364	55.3	49.42	60.96
Female	621	87,225	63.3	56.53	69.52
School level					
Lower secondary	613	93,409	56.8	50.65	62.81
Upper secondary	414	69,180	63.0	56.15	69.29
Form					
Form 1	197	32,936	57.5	50.75	63.92
Form 2	209	27,619	51.9	42.94	60.81
Form 3	207	32,854	61.0	51.25	69.92
Form 4	220	31,945	59.2	51.18	66.83
Form 5	194	37,235	66.5	57.68	74.38

Table 3.3.7 : Prevalence of milk or milk products intake at least two servings per day in the past 30 days among Form 1 to Form 5 students in Johor, 2017

Socio-demographic characteristics	Unweighted Count	Estimated Population	Prevalence %	95% CI	
				Lower	Upper
JOHOR	466	74,744	27.2	23.76	30.83
Locality of school					
Urban	319	50,530	25.8	21.94	30.10
Rural	147	24,214	30.5	25.51	35.92
Sex					
Male	208	37,630	27.5	23.20	32.31
Female	258	37,114	26.8	22.59	31.44
School level					
Lower secondary	330	50,919	30.8	26.90	35.00
Upper secondary	136	23,825	21.7	17.00	27.18
Form					
Form 1	125	21,194	36.7	33.74	39.72
Form 2	123	16,450	30.9	24.18	38.47
Form 3	82	13,275	24.5	18.89	31.10
Form 4	71	11,080	20.5	15.51	26.60
Form 5	65	12,745	22.8	16.93	29.93

Table 3.3.8: Prevalence of consuming food from fast food restaurants of at least three days in the past seven days among Form 1 to Form 5 students in Johor, 2017

Socio-demographic characteristics	Yes (at least three days)				
	Unweighted Count	Estimated Population	Prevalence (%)	95% CI	
				Lower	Upper
JOHOR	119	19,631	7.1	5.18	9.73
Locality of school					
Urban	84	13,709	7.0	4.46	10.81
Rural	35	5,922	7.5	6.11	9.06
Sex					
Male	57	10,651	7.8	5.50	10.91
Female	62	8,980	6.5	4.45	9.35
School level					
Lower secondary	63	9,802	5.9	3.90	8.93
Upper secondary	56	9,829	8.9	5.99	13.07
Form					
Form 1	14	2,585	4.5	2.11	9.23
Form 2	23	3,190	6.0	2.85	12.21
Form 3	26	4,027	7.4	4.75	11.44
Form 4	31	4,780	8.8	5.91	13.03
Form 5	25	5,049	9.0	5.25	14.96

3.4 Drug Use

Contributors: Muhammad Fadhli Mohd Yusoff, Rushidi Ramly, Wan Shakira Rodzlan Hasani, Hasimah Ismail, Hamizatul Akmal Abd Hamid, Norli Abdul Jabbar, Jane Ling Miaw Yn, Halizah Mat Rifin, Tania Gayle Robert, Thamil Arasu Saminathan, Nur Liana Ab. Majid.

3.4.1 Introduction

According to the World Drug Report 2017, an estimated quarter of a billion people, or around five per cent of the global adult population, used drugs at least once in 2015. Even more worrisome is the fact that about 29.5 million of those drug users, or 0.6 percent of the global adult population, suffer from drug use disorders.¹ This means that their drug use is harmful to the point that they may experience drug dependence and require treatment.

Marijuana or cannabis remains the world's most widely used drug, with an annual prevalence of 3.8 per cent of the adult population, or an estimated 183 million people (range 128 million to 238 million), having used marijuana in the past year.² Amphetamines remain the second most commonly used drug worldwide, with an estimated 35 million past-year users (range 13 million to 58 million), and the use of amphetamines, particularly methamphetamine, is perceived to be increasing in many regions, including most parts of Asia.

This drug menace contributes to various social and medical ill health implications particularly among adolescents. The broad range of problems reported by young people include deteriorating family relations, truancy, poorer performance in school, mental disorders such as depression and anxiety, drug induced psychosis particularly with the New Psychoactive Substances (NPS), unwanted and unprotected sexual activity, accidents, violence and increased risk of blood-borne viral diseases such as HIV and Hepatitis C for those injecting drug users (IDU). Some youth engaging in heavy substance use will continue to do so into adulthood and will experience various longer-term health and social problems.

3.4.2 Objectives

- i. To determine the prevalence of ever and current drug use among Form 1 to Form 5 students
- ii. To describe the socio-demographic characteristics of ever and current drug use among Form 1 to Form 5 students
- iii. To determine the prevalence of ever and current marijuana use among Form 1 to Form 5 students
- iv. To describe the socio-demographic characteristics of ever and current marijuana use among Form 1 to Form 5 students
- v. To determine the prevalence and socio-demographic characteristics of ever amphetamines or methamphetamines use among Form 1 to Form 5 students
- vi. To identify the age of initiation and the sources of obtaining drugs among Form 1 to Form 5 students

3.4.3 Variable Definitions

- **Drug use:** taking of heroin, morphine, glue, amphetamine or methamphetamines (ecstasy, syabu, ice), marijuana/*ganja* (except prescribed medicine).
- **Ever drug use:** students who had history of drug use in their lifetime
- **Current drug use:** students who used drugs in the past 30 days
- **Ever Marijuana use:** students who had history of marijuana use in their lifetime
- **Current Marijuana use:** students who used marijuana in the past 30 days
- **Ever Amphetamine or Methamphetamines use:** students who had history of amphetamine or methamphetamines use in their lifetime.

3.4.4 Findings

Overall, the prevalence of ever drug use among students from Form 1 to Form 5 in Johor was 2.6% (95% CI: 1.61, 4.12) with an estimated population of 7,124 students. The prevalence of ever drug use was significantly higher among males [(4.7%, 95% CI: 3.06, 7.01)] as compared to females [0.5% (95%CI: 0.18-1.62)]. Those who studied in rural areas [3.2% (95% CI: 1.83, 5.67)] and Form 2 students [3.7% (95% CI: 1.06, 12.34)] had the highest prevalence of ever used drugs. **(Table 3.4.1)**. The prevalence of current drug use in the past 30 days among Form 1 to Form 5 students in Johor was 1.6% (95% CI: 0.79, 3.37). The current drug users were highest among those who were from urban schools [1.9% (95% CI: 0.81, 4.31)], males (2.9%, 95% CI: 1.51, 5.41) and Form 2 students (3.2%, 95% CI: 0.74, 13.09) **(Table 3.4.2)**. The prevalence of ever used marijuana in lifetime was 1.3% (95% CI: 0.71, 2.52) among Form 1 to Form 5 students in Johor **(Table 3.4.3)** and 1.2% (95% CI: 0.51, 2.65) were current marijuana users in the past 30 days **(Table 3.4.4)**. The prevalence was significantly higher in males [2.5% (95% CI: 1.32, 4.64)] compared to females among ever marijuana users. Meanwhile, about 0.9% (95% CI: 0.38, 2.14) students reported that they had ever used amphetamines or methamphetamines during lifetime **(Table 3.4.5)**. The prevalence was highest among males and Form 2 students with the prevalence of 1.6% (95% CI: 0.76, 3.39) and 2.1% (95% CI: 0.38, 10.56) respectively.

Among current drug users, most of the students obtained their drugs by buying them from someone [31.5% (95% CI: 17.19, 50.50)]. Other than that, they also obtained the drugs from their friends [4.6% (95%CI: 0.41, 36.05)], gave someone else money to buy the drugs [3.0% (95%CI: 0.31, 23.09)] or got the drugs from other ways [6.7% (95% CI: 1.28, 28.59)] **(Table 3.4.6)**. Among ever drug users, about 16.1% (95% CI: 6.52, 34.59) had initiated drug use at the age of 7 years or younger. The percentage by school locality was 0.5% (95% CI: 0.17, 1.48) in urban areas and 0.2% (95% CI: 0.03, 1.39) in rural areas **(Table 3.4.7)**.

3.4.5 Discussion/ Conclusion

The prevalence of ever drug use and current drug use in Johor did not differ with the national prevalence. However, the prevalence of current drug use was significantly higher in male students. As for ever used marijuana and ever used amphetamines or methamphetamine,

the prevalence was also significantly higher among males. Among those who ever used drugs, majority of students admitted that they had been using drugs at the age of seven years or younger. Most of those who were current drug users had obtained their drugs by buying from someone.

3.4.6 Recommendations

Prevention and control measures of drug use among adolescents in Malaysia needs to be strengthened and started early from the primary school level. New strategies and approaches must be developed to address this issue. Health education and the dangers of drug abuse should be emphasized. Early detection and intervention also needs to be given priority.

3.4.7 References

1. Word Drug Report 2017: Executive Summary Conclusions and Policy Implications. (ISBN:978-92-1-148291-1, eISBN: 978-92-1-060623-3, United Nations publication, Sales No. E.17.XI.6).
2. United Nations Office on Drugs and Crime, *World Drug Report 2017: Global Overview*. (ISBN: 978-92-1-148291-1, eISBN: 978-92-1-060623-3, United Nations publication, Sales No. E.17.XI.6).

Table 3.4.1: Prevalence of ever used drugs in lifetime among Form 1 to Form 5 students in Johor, 2017

Socio-demographic Characteristics	Unweighted Count	Estimated Population	Prevalence (%)	95% CI	
				Lower	Upper
JOHOR	40	7,124	2.6	1.61	4.12
Locality of school					
Urban	26	4,552	2.3	1.21	4.41
Rural	14	2,571	3.2	1.83	5.67
Sex					
Male	35	6,382	4.7	3.06	7.01
Female	5	742	0.5	0.18	1.62
Form					
Form 1	6	1,140	2.0	0.63	5.95
Form 2	13	1,996	3.7	1.06	12.34
Form 3	7	1,239	2.3	1.03	4.99
Form 4	9	1,584	2.9	1.24	6.78
Form 5	5	1,164	2.1	0.62	6.73

Table 3.4.2 : Prevalence of current drug use in the past 30 days among Form 1 to Form 5 students in Johor, 2017

Socio-demographic Characteristics	Unweighted Count	Estimated Population	Prevalence (%)	95% CI	
				Lower	Upper
JOHOR	26	4,529	1.6	0.79	3.37
Locality of school					
Urban	21	3,682	1.9	0.81	4.31
Rural	5	846	1.1	0.39	2.85
Sex					
Male	22	3,941	2.9	1.51	5.41
Female	4	587	0.4	0.11	1.61
Form					
Form 1	2	355	0.6	0.08	4.51
Form 2	11	1,729	3.2	0.74	13.09
Form 3	5	859	2.3	0.54	4.56
Form 4	5	856	1.6	0.48	5.12
Form 5	3	730	1.3	0.26	6.30

Table 3.4.3 : Prevalence of ever used marijuana in lifetime among students Form 1 to Form 5, Johor 2017

Socio-demographic Characteristics	Unweighted Count	Estimated Population	Prevalence (%)	95% CI	
				Lower	Upper
JOHOR	20	3704	1.3	0.71	2.52
Locality of the school					
Urban	14	2448	1.2	0.55	2.81
Rural	6	1256	1.6	0.61	4.06
Sex					
Male	18	3412	2.5	1.32	4.64
Female	2	292	0.2	0.05	0.97
Form					
Form 1	-	-	-	-	-
Form 2	5	718	1.3	0.39	4.60
Form 3	5	936	2.3	0.60	4.83
Form 4	6	1041	1.9	0.75	4.86
Form 5	4	1010	1.8	0.50	6.21

Table 3.4.4 : Prevalence of current marijuana use in the past 30 days among Form 1 to Form 5 students in Johor, 2017

Socio-demographic Characteristics	Unweighted Count	Estimated Population	Prevalence (%)	95% CI	
				Lower	Upper
JOHOR	18	3,214	1.2	0.51	2.65
Locality of school					
Urban	15	2,703	1.4	0.54	3.45
Rural	3	510	0.6	0.19	2.19
Sex					
Male	15	2,787	2.0	0.94	4.35
Female	3	427	0.3	0.09	1.04
Form					
Form 1	1	178	0.3	0.04	2.29
Form 2	6	981	1.8	0.28	11.02
Form 3	5	859	2.3	0.54	4.56
Form 4	4	694	1.3	0.38	4.27
Form 5	2	502	0.9	0.10	7.73

Table 3.4.5: Prevalence of ever amphetamines/metaphetamines use in a lifetime among Form 1 to Form 5 students in Johor, 2017

Socio-demographic Characteristics	Unweighted Count	Estimated Population	Prevalence (%)	95% CI	
				Lower	Upper
JOHOR	14	2,507	0.9	0.38	2.14
Locality of school					
Urban	10	1,743	0.9	0.27	2.88
Rural	4	764	1.0	0.43	2.15
Sex					
Male	12	2,215	1.6	0.76	3.39
Female	2	292	0.2	0.05	0.97
Form					
Form 1	3	577	1.0	0.24	4.06
Form 2	7	1,109	2.1	0.38	10.56
Form 3	-	-	2	-	-
Form 4	2	319	0.6	0.13	2.70
Form 5	2	502	0.9	0.10	7.73

Table 3.4.6 : Source of getting drugs in the past 30 days among current drug users, Form 1 to Form 5 students in Johor, 2017

	Unweighted Count	Estimated Population	Prevalence (%)	95% CI	
				Lower	Upper
I bought them from someone	7	1,348	31.5	17.19	50.50
I gave someone else money to buy it for me	1	128	3.0	0.31	23.09
I stole it or got it without permission	-	-	-	-	-
I got it from my friend	1	197	4.6	0.41	36.05
I got it from my family	-	-	-	-	-
I got it some other ways	2	287	6.7	1.28	28.59

Table 3.4.7 : Age of initiation of first use of drugs amongst ever drug users among Form 1 to Form 5 students in Johor, 2017

	Total					Urban				
	Unweighted Count	Estimated Population	Prevalence (%)	95% CI		Unweighted Count	Estimated Population	Prevalence (%)	95% CI	
				Lower	Upper				Lower	Upper
7 years old or younger	7	1,148	16.1	6.52	34.59	6	989	0.5	0.17	1.48
8 or 9 years old	1	178	2.5	0.29	18.38	-	-	-	-	-
10 or 11 years old	1	140	2.0	0.20	16.97	-	-	-	-	-
12 or 13 years old	6	1,058	14.9	8.33	25.11	4	719	0.4	0.15	0.90
14 or 15 years old	3	558	7.8	1.51	31.93	1	173	0.1	0.01	0.71
16 years old or older	2	286	4.0	0.96	15.32	2	286	0.1	0.04	0.60

Table 3.4.7 : Age of initiation of first use of drugs amongst ever drug users among Form 1 to Form 5 students in Johor, 2017 (cont.)

	Rural				
	Unweighted Count	Estimated Population	Prevalence (%)	95% CI	
				Lower	Upper
7 years old or younger	1	158	0.2	0.03	1.39
8 or 9 years old	1	178	0.2	0.03	1.56
10 or 11 years old	1	140	0.2	0.03	1.12
12 or 13 years old	2	339	0.4	0.13	1.42
14 or 15 years old	2	385	0.5	0.06	3.55
16 years old or older	-	-	-	-	-

3.5 Hygiene (Including Oral Health)

Contributors: Yaw Siew Lian, Natifah Che Salleh, Nurrul Ashikin Abdullah, Ting Teck Pei, Mohamad Fuad Mohamad Anuar, Sayan a/I Pan, Norazizah Ibrahim Wong.

3.5.1 Introduction

Dental caries has substantial impact to children, families, and health systems, including dental and medical consequences, loss of working time on the part of caregivers and increased expenditures. It is a multi-factorial infectious disease that leads to pain and infection, consequently affecting the quality of life, general health, productivity and educational performance of the child.¹ Daily tooth brushing with a fluoridated toothpaste and regular dental check-up at least once a year can help prevent dental caries and periodontal disease.

Clean hands can stop spread of parasites from one person to another. These parasites consume nutrients from children they infect, causing abdominal pain and can impair learning by slowing cognitive development. Hand washing is one of the most effective ways to prevent the spread of parasites, especially during these key times: before, during and after preparing food, before eating and after using the toilet.^{2,3}

3.5.2 Objectives

To describe the prevalence of:

- i. Tooth brushing frequency in the past 30 days
- ii. Fluoridated toothpaste usage
- iii. Dental floss usage
- iv. Timing of last visit to a dentist or dental nurse
- v. Having missed class due to toothache in the past 12 months
- vi. Avoidance of smiling or laughing due to the appearance of their teeth
- vii. Using soap during hand washing in the past 30 days
- viii. Hand washing before eating in the past 30 days
- ix. Hand washing after using the toilet in the past 30 days

3.5.3 Variable Definitions

- **Clean or brush teeth:** Regular tooth brushing using toothbrush and toothpaste to keep the mouth, teeth and gums clean and healthy
- **Last saw a dentist or dental nurse:** Seen a dentist or dental nurse for a check-up, scaling or other dental treatment

3.5.4 Findings

Cleaning or brushing teeth during the past 30 days

Overall, only 0.5% (95%CI: 0.24, 0.91) of students reported they had never cleaned or brushed their teeth during the past 30 days. About 1.4% (95%CI: 0.88, 2.28) reported they had cleaned or brushed their teeth less than once daily, 9.9% (95%CI: 8.26, 11.77) once daily and 88.2% (95%CI: 86.15, 90.05) at least twice daily (**Table 3.5.1**).

Use of fluoridated toothpaste

Overall, only 54.7% (95%CI: 47.79, 61.42) of students reported use of fluoridated toothpaste and 7.4% (95%CI: 5.56, 9.86) claimed that they were not using fluoridated toothpaste. About 37.9% (95%CI: 31.81, 44.36)] of students reported not knowing whether their toothpaste contained fluoride (**Table 3.5.2**).

Use of dental floss for cleaning teeth

Overall, only 13.7% (95%CI: 11.95, 15.60) of students reported that they use dental floss for cleaning their teeth and 86.3% (95%CI: 84.40, 88.05) claimed that they do not use dental floss to clean their teeth (**Table 3.5.3**).

Last dental visit

Overall, only 45.7% (95%CI: 42.31, 49.08) of students reported that they had their last dental visit (inclusive of both dental treatment and dental check-up) in the past 12 months. The prevalence of students who had their last dental visit between 12-24 months was 11.5% (95%CI: 9.00, 14.54) while those with last dental visit more than 24 months was 6.1% (95%CI: 4.67, 7.87). About 9.8% (95%CI: 6.89, 13.75) of students had never ever had a dental visit

while a substantial proportion [27.0% (95%CI: 22.79, 31.62)] did not know when was their last dental visit (**Table 3.5.4**).

Missing class or school due to toothache

Overall, 4.3% (95% CI: 3.05, 6.16) of students had missed class or school due to toothache in the past 12 months (**Table 3.5.5**).

Avoid smiling or laughing due to the appearance of their teeth

Overall, 15.4% (95%CI: 13.45, 17.55) of students reported that they avoided smiling or laughing due to the appearance of their teeth (**Table 3.5.6**). By school grade, the prevalence of students in Form 1 [12.6% (95%CI: 9.91, 15.86)] and Form 2 [14.0% (95%CI: 12.31, 15.86)] who had avoided smiling or laughing due to the appearance of their teeth was significantly lower compared to Form 4 [21.8% (95%CI: 15.93, 20.09)].

Use of soap when washing hands during the past 30 days

Overall in the past 30 days, about 9.5% (95%CI: 7.79, 11.59) of students reported that they never or rarely used soap when washing their hands, 25.4% (95%CI: 22.66, 28.29) reported they had only used soap sometimes when washing their hands while a fairly high proportion of 65.1% (95%CI: 61.35, 68.68) had always used soap when washing their hands (**Table 3.5.7**).

Washing hand before eating during the past 30 days

Overall in the past 30 days, about 4.4% (95%CI: 2.82, 6.80) of students never or rarely washed their hands before eating, 10.0% (95%CI: 7.87, 12.72) reported they had sometimes washed their hands before eating while a fairly high proportion of students [85.6% (95%CI: 81.29, 89.00)] had always washed their hands before eating (**Table 3.5.8**).

Washing hand after using the toilet or latrine during the past 30 days

Overall in the last 30 days, about 2.8% (95%CI: 2.10, 3.72) of students reported they had never or rarely washed their hands after using the toilet, 7.8% (95%CI: 6.26, 9.63) reported they had only washed their hands sometimes after using the toilet while 89.4% (95%CI: 87.10, 91.36) of students claimed that they had always or most of the times washed their hands after using the toilet (**Table 3.5.9**).

3.5.5 Discussion/ Conclusion

The above findings related to the hand hygiene and oral hygiene practises of students in the age group of 13-17 years in Johor. Overall, the prevalence of students who had cleaned/brushed their teeth less than one time per day during the past 30 days (1.4%) was lower than the findings for Johor in GSHS 2012 (2.6%)⁴ and the national prevalence in 2017 (2.4%). This survey also found that almost a similar proportion of students (54.7%) use fluoridated toothpaste in comparison with the findings for Johor in GSHS 2012 (54.1%)⁴ and lower compared to national prevalence in 2017 (58.3%). The prevalence of students who did not know whether their toothpaste contained fluoride (37.9%) was lower than the national prevalence (31.8%) and higher than findings for Johor in GSHS 2012 (29.3%).⁴ The survey findings also showed that a significantly lower proportion of students (13.7%) reported using dental floss to clean their teeth as compared to the national prevalence (19.3%). The use of dental floss was not studied in the first national GSHS in 2012.

Our findings on the proportion of students who never or rarely used soap when washing their hands during the past 30 days in this study (9.5%) was significantly lower than the findings for Johor in GSHS 2012 (15.2%)⁴ and also lower than the national prevalence (11.6%). This study also found that the prevalence of students who never or rarely washed their hands before eating (4.4%) was lower compared to the findings for Johor in GSHS 2012 (6.8%)⁴ and the national prevalence (6.1%). The prevalence of students who never or rarely washed their hands after going to the toilet in this study (2.8%) was significantly lower compared to both the findings for Johor in GSHS 2012 (6.3%)⁴ and the national prevalence (4.8%).

Overall, the findings in relation to hand hygiene and oral hygiene practises of students in the age group of 13-17 years showed that there is still room for further improvement. In terms of oral hygiene, the prevalence of students who had cleaned/brushed their teeth less than one time per day (1 in 100) during the past 30 days has improved compared to previous study (almost 3 in 100).⁴ Other areas of concern were the low proportion of students (about 1 in 2 students) reported using a fluoridated toothpaste and the low proportion (1 in 10) of students who reported use of dental floss to clean their teeth. In terms of hand hygiene, about 1 in 10 students reported that they had never or rarely used soap when washing their hands, and 4

in 100 students reported they had never or rarely washed their hands before eating. In addition, the proportion of students who never or rarely washed their hands after going to the toilet was about 3 in 100. Overall, these hygiene practises did not significantly differ by school locality, gender and school grade.

3.5.6 Recommendations

Taking cognizance of these findings, there is a need to emphasis on promoting hygiene among school-going adolescents through attitude and behavioural changes in maintenance of personal hygiene. Thus, the following recommendations are made:

1. More effort is needed to strengthen health education on oral and hand hygiene habits in schools and at home.
2. There is a need to further promote toothpaste literacy in our oral health promotion efforts.
3. Surveillance of hygiene habits and practices among adolescents need to be continued at regular intervals.
4. Further studies are recommended to study the factors contributing to poor hygiene habits among adolescents.

3.5.7 References

1. Mathur V.P. & Dhillon J.K. Dental Caries: A Disease Which Needs Attention. *Indian J Pediatr* (2017). <https://doi.org/10.1007/s12098-017-2381-6>. DOI<https://doi.org/10.1007/s12098-017-2381-6>. Publisher: Springer India
2. When & How to Wash Your Hands. Centers for Disease Control and Prevention. <https://www.cdc.gov/handwashing/when-how-handwashing.html>. Retrieve 7 August 2017
3. Borchgrevink CP, Cha JM, Kim SH. Hand washing practices in a College Town Environment. *J of Environmental Health*, Volume 75, No. 8, April 2013.
4. Institute for Public Health (IPH) 2012. The National Health and Morbidity Survey: Johor Global School-based Student Health Survey 2012. Kuala Lumpur: Ministry of Health Malaysia.

Table 3.5.1: Prevalence of brushing teeth in the past 30 days among Form 1 to Form 5 students in Johor, 2017

Socio-demographic characteristics	Never clean or brush teeth					Less than 1 day				
	Unweighted Count	Estimated Population	Prevalence (%)	95% CI		Unweighted Count	Estimated Population	Prevalence (%)	95% CI	
				Lower	Upper				Lower	Upper
JOHOR	7	1,289	0.5	0.24	0.91	22	3,907	1.42	0.88	2.28
Locality of school										
Urban	5	883	0.4	0.19	1.06	18	3,264	1.66	1.00	2.76
Rural	2	406	0.5	0.19	1.38	4	643	0.81	0.35	1.85
Sex										
Male	7	1,289	0.9	0.49	1.80	17	3,174	2.31	1.52	3.51
Female						5	733	0.53	0.25	1.14
Form										
Form 1	1	193	0.3	0.05	2.34	5	1,057	1.82	0.69	4.70
Form 2	4	597	1.1	0.48	2.60	5	728	1.37	0.67	2.78
Form 3	-	-	-	-	-	4	648	1.20	0.51	2.77
Form 4	-	-	-	-	-	5	801	1.48	0.56	3.86
Form 5	2	499	0.9	0.22	3.48	3	673	1.20	0.16	8.56

Table 3.5.1: Prevalence of brushing teeth in the past 30 days among Form 1 to Form 5 students in Johor, 2017 (cont.)

Socio-demographic characteristics	Once per day					2 times or more per day				
	Unweighted Count	Estimated Population	Prevalence (%)	95% CI		Unweighted Count	Estimated Population	Prevalence (%)	95% CI	
				Lower	Upper				Lower	Upper
JOHOR	160	27,205	9.9	8.26	11.77	1,541	243,133	88.2	86.15	90.05
Locality of school										
Urban	120	20,364	10.4	8.25	12.98	1,103	171,726	87.5	84.75	89.83
Rural	40	6,841	8.6	8.00	9.30	438	71,407	90.1	88.94	91.06
Sex										
Male	106	19,686	14.4	11.11	18.34	617	112,998	82.4	78.66	85.59
Female	54	7,519	5.4	4.15	7.09	924	130,135	94.0	92	95
Form										
Form 1	32	5,808	10.0	8.07	12.39	303	50,902	87.8	84.48	90.53
Form 2	33	4,631	8.7	6.07	12.29	357	47,342	88.8	83.99	92.34
Form 3	33	5,518	10.2	6.79	14.99	301	48,048	88.6	83.69	92.21
Form 4	40	6,489	12.0	7.54	18.59	321	46,745	86.5	80.77	90.73
Form 5	22	4,759	8.5	5.63	12.61	259	50,095	89.4	83.77	93.26

Table 3.5.2: Prevalence of use of fluoridated toothpaste among Form 1 to Form 5 students in Johor, 2017.

Socio-demographic characteristics	Yes					No				
	Unweighted Count	Estimated Population	Prevalence (%)	95% CI		Unweighted Count	Estimated Population	Prevalence (%)	95% CI	
				Lower	Upper				Lower	Upper
JOHOR	933	150,538	54.7	47.79	61.42	126	20,442	7.4	5.56	9.86
Locality of school										
Urban	693	110,459	56.4	48.29	64.23	88	13,997	7.1	4.84	10.45
Rural	240	40,079	50.4	38.78	62.03	38	6,445	8.1	5.78	11.27
Sex										
Male	396	74,088	54.1	45.43	62.51	71	12,823	9.4	6.95	12.50
Female	537	76,449	55.3	47.86	62.49	55	7,618	5.5	4.06	7.45
Form										
Form 1	185	30,828	53.4	43.54	62.92	30	5,460	9.5	5.81	15.00
Form 2	176	23,186	43.6	31.74	56.24	46	6,604	12.4	5.98	24.01
Form 3	187	30,439	56.1	45.14	66.58	18	2,904	5.4	3.45	8.22
Form 4	214	32,048	59.3	54.12	64.30	19	2,851	5.28	3.53	7.82
Form 5	171	34,036	60.7	50.74	69.92	13	2,622	4.68	2.21	9.63

Table 3.5.2: Prevalence of use of fluoridated toothpaste among Form 1 to Form 5 students in Johor, 2017 (cont.)

Socio-demographic characteristics	Don't Know				
	Unweighted Count	Estimated Population	Prevalence (%)	95% CI	
				Lower	Upper
JOHOR	669	104,259	37.9	31.81	44.36
Locality of school					
Urban	462	71,308	36.4	29.23	44.28
Rural	207	32,951	41.5	31.67	51.98
Sex					
Male	279	50,056	36.5	29.38	44.37
Female	390	54,203	39.2	32.15	46.73
Form					
Form 1	125	21,492	37.2	29.92	45.10
Form 2	176	23,385	44.0	32.89	55.70
Form 3	133	20,871	38.5	27.62	50.67
Form 4	133	19,136	35.4	30.06	41.16
Form 5	102	19,375	34.6	26.87	43.19

Table 3.5.3: Prevalence of use of dental floss among Form 1 to Form 5 students in Johor, 2017.

Socio-demographic characteristics	Yes					No				
	Unweighted Count	Estimated Population	Prevalence (%)	95% CI		Unweighted Count	Estimated Population	Prevalence (%)	95% CI	
				Lower	Upper				Lower	Upper
JOHOR	232	37,676	13.7	11.95	15.60	1,498	237,905	86.3	84.40	88.05
Locality of school										
Urban	176	28,565	14.6	12.68	16.68	1,069	167,541	85.4	83.32	87.32
Rural	56	9,111	11.5	8.79	14.83	429	70,364	88.5	85.17	91.21
Sex										
Male	100	19,119	13.9	11.66	16.58	647	118,028	86.1	83.42	88.34
Female	132	18,557	13.4	11.70	15.31	851	119,876	86.6	84.69	88.30
Form										
Form 1	54	8,994	15.6	12.35	19.40	286	48,835	84.4	80.60	87.65
Form 2	46	6,183	11.6	7.88	16.76	353	47,115	88.4	83.24	92.12
Form 3	44	7,153	13.2	9.90	17.38	294	47,061	86.8	82.62	90.10
Form 4	44	6,314	11.7	8.23	16.33	322	47,721	88.3	83.67	91.77
Form 5	44	9,032	16.1	11.19	22.54	243	47,171	83.9	77.46	88.81

Table 3.5.4: Timing of last dental visit among Form 1 to Form 5 students in Johor, 2017

Socio-demographic characteristics	During the past 12 months					Between 12-24 months ago				
	Unweighted Count	Estimated Population	Prevalence (%)	95% CI		Unweighted Count	Estimated Population	Prevalence (%)	95% CI	
				Lower	Upper				Lower	Upper
JOHOR	790	125,808	45.7	42.31	49.08	205	31,635	11.5	9.00	14.54
Locality of school										
Urban	576	91,419	46.6	42.32	51.03	160	24,057	12.3	9.16	16.26
Rural	214	34,388	43.3	39.81	46.80	45	7,578	9.5	6.65	13.50
Sex										
Male	315	58,394	42.6	38.06	47.31	77	14,116	10.3	7.68	13.69
Female	475	67,414	48.7	44.86	52.55	128	17,519	12.7	10.04	15.84
Form										
Form 1	144	24,415	42.1	33.76	50.97	39	6,493	11.2	7.91	15.64
Form 2	154	20,577	38.8	32.03	46.02	50	6,374	12.0	7.87	17.93
Form 3	159	25,264	46.6	40.04	53.28	35	5,618	10.4	7.46	14.23
Form 4	173	24,913	46.1	37.64	54.80	49	6,955	12.9	8.35	19.32
Form 5	160	30,638	54.5	44.01	64.63	32	6,196	11.0	7.68	15.58

Table 3.5.4: Timing of last dental visit among Form 1 to Form 5 students in Johor, 2017 (cont.)

Socio-demographic characteristics	More than 24 months ago					Never				
	Unweighted Count	Estimated Population	Prevalence (%)	95% CI		Unweighted Count	Estimated Population	Prevalence (%)	95% CI	
				Lower	Upper				Lower	Upper
JOHOR	99	16,727	6.1	4.67	7.87	166	26,978	9.8	6.89	13.75
Locality of school										
Urban	75	12,627	6.4	4.78	8.64	107	17,051	8.7	5.48	13.55
Rural	24	4,099	5.2	3.07	8.55	59	9,927	12.5	7.77	19.47
Sex										
Male	59	11,255	8.2	6.20	10.81	89	15,678	11.4	8.24	15.67
Female	40	5,472	4.0	2.82	5.51	77	11,300	8.2	5.17	12.66
Form										
Form 1	19	3,650	6.3	3.68	10.57	25	4,443	7.7	4.60	12.50
Form 2	26	3,486	6.6	4.17	10.22	60	8,406	15.8	10.09	24.00
Form 3	17	2,894	5.3	3.29	8.55	36	5,921	10.9	7.36	15.92
Form 4	22	3,376	6.2	3.67	10.43	29	4,767	8.8	5.17	14.65
Form 5	15	3,321	5.9	3.26	10.49	16	3,441	6.1	3.41	10.75

Table 3.5.4: Timing of last dental visit among Form 1 to Form 5 students in Johor, 2017 (cont.)

Socio-demographic characteristics	Don't know				
	Unweighted Count	Estimated Population	Prevalence (%)	95% CI	
				Lower	Upper
JOHOR	469	74,309	27.0	22.79	31.62
Locality of school					
Urban	326	50,827	25.9	20.53	32.19
Rural	143	23,482	29.5	25.43	34.03
Sex					
Male	206	37,578	27.4	22.95	32.40
Female	263	36,731	26.5	21.61	32.12
Form					
Form 1	114	18,960	32.7	28.75	36.94
Form 2	107	14,200	26.8	22.25	31.84
Form 3	91	14,517	26.8	20.01	34.84
Form 4	93	14,025	26.0	17.35	36.92
Form 5	64	12,607	22.4	14.96	32.21

Table 3.5.4.1: Prevalence of last dental visit within the last 24 months among Form 1 to Form 5 students in Johor, 2017.

Socio-demographic characteristics	Unweighted Count	Estimated Population	Prevalence (%)	95% CI	
				Lower	Upper
JOHOR	995	157,443	57.2	52.56	61.64
Locality of school					
Urban	736	115,476	58.9	52.88	64.71
Rural	259	41,966	52.8	50.10	55.49
Sex					
Male	392	72,509	52.9	47.81	57.97
Female	603	84,933	61.4	56.30	66.17
Form					
Form 1	183	30,908	53.3	46.81	59.73
Form 2	204	26,951	50.8	43.09	58.49
Form 3	194	30,882	57.0	49.42	64.20
Form 4	222	31,868	59.0	47.77	69.32
Form 5	192	36,834	65.5	54.27	75.29

Table 3.5.5: Prevalence of having missed class due to toothache in the past 12 months among Form 1 to Form 5 students in Johor, 2017.

Socio-demographic characteristics	Yes					No				
	Unweighted Count	Estimated Population	Prevalence (%)	95% CI		Unweighted Count	Estimated Population	Prevalence (%)	95% CI	
				Lower	Upper				Lower	Upper
JOHOR	76	11,992	4.3	3.05	6.16	1,655	263,720	95.7	93.84	96.95
Locality of school										
Urban	58	9,181	4.7	3.00	7.23	1,188	187,056	95.3	92.77	97.00
Rural	18	2,811	3.5	2.69	4.63	467	76,664	96.5	95.37	97.31
Sex										
Male	34	5,978	4.4	2.46	7.61	713	131,169	95.6	92.39	97.54
Female	42	6,014	4.3	3.19	5.87	942	132,550	95.7	94.13	96.81
Form										
Form 1	12	2,094	3.6	2.34	5.55	329	55,867	96.4	94.45	97.66
Form 2	27	3,906	7.3	3.48	14.77	372	49,392	92.7	85.23	96.52
Form 3	13	2,099	3.9	2.90	5.15	325	52,115	96.1	94.85	97.10
Form 4	17	2,563	4.7	2.09	10.42	349	51,472	95.3	89.58	97.91
Form 5	7	1,329	2.4	0.92	5.97	280	54,874	97.6	94.03	99.08

Table 3.5.6: Prevalence of avoidance of smiling or laughing due to the appearance of their teeth among Form 1 to Form 5 students in Johor, 2017.

Socio-demographic characteristics	Yes					No				
	Unweighted Count	Estimated Population	Prevalence (%)	95% CI		Unweighted Count	Estimated Population	Prevalence (%)	95% CI	
				Lower	Upper				Lower	Upper
JOHOR	268	42,320	15.4	13.45	17.55	1,459	232,722	84.6	82.45	86.55
Locality of school										
Urban	197	31,188	15.9	13.48	18.77	1,045	164,379	84.1	81.23	86.52
Rural	71	11,132	14.0	11.91	16.40	414	68,342	86.0	83.60	88.09
Sex										
Male	107	19,662	14.4	11.48	17.86	638	117,093	85.6	82.14	88.52
Female	161	22,658	16.4	14.70	18.22	821	115,628	83.6	81.78	85.30
Form										
Form 1	42	7,246	12.6	9.91	15.86	297	50,322	87.4	84.14	90.09
Form 2	55	7,441	14.0	12.31	15.86	343	45,739	86.0	84.14	87.69
Form 3	51	7,903	14.6	10.90	19.23	287	46,311	85.4	80.77	89.10
Form 4	79	11,782	21.8	15.93	29.09	287	42,253	78.2	70.91	84.07
Form 5	41	7,948	14.2	11.14	17.89	245	48,096	85.8	82.11	88.86

Table 3.5.7: Prevalence of use of soap during hand washing in the past 30 days among Form 1 to Form 5 in Johor, 2017

Socio-demographic characteristics	Never or rarely use soap during hand washing					Sometimes				
	Unweighted Count	Estimated Population	Prevalence (%)	95% CI		Unweighted Count	Estimated Population	Prevalence (%)	95% CI	
				Lower	Upper				Lower	Upper
JOHOR	165	26,241	9.5	7.79	11.59	434	69,931	25.4	22.66	28.29
Locality of school										
Urban	129	20,440	10.4	8.35	12.94	305	48,386	24.67	21.27	28.42
Rural	36	5,801	7.3	5.67	9.34	129	21,545	27.11	23.60	30.93
Sex										
Male	71	13,080	9.5	7.08	12.73	196	36,292	26.46	23.40	29.77
Female	94	13,160	9.5	7.99	11.28	238	33,640	24.30	20.92	28.03
Form										
Form 1	27	4,647	8.0	5.20	12.17	86	14,854	25.63	19.53	32.85
Form 2	45	6,155	11.5	8.31	15.82	104	13,661	25.63	21.03	30.84
Form 3	34	5,319	9.8	7.10	13.40	83	13,437	24.78	19.79	30.56
Form 4	29	4,380	8.1	4.36	14.64	83	12,704	23.56	18.93	28.92
Form 5	30	5,741	10.2	6.86	14.95	78	15,275	27.18	21.26	34.04

Table 3.5.7: Prevalence of use of soap during hand washing in the past 30 days among Form 1 to Form 5 in Johor, 2017 (cont.)

Socio-demographic characteristics	Most of the times or Always				
	Unweighted Count	Estimated Population	Prevalence (%)	95% CI	
				Lower	Upper
JOHOR	1,131	179,430	65.1	61.35	68.68
Locality of school					
Urban	811	127,302	64.9	60.13	69.40
Rural	320	52,128	65.6	59.95	70.83
Sex					
Male	480	87,775	64.0	59.81	67.99
Female	651	91,655	66.2	61.94	70.21
Form					
Form 1	228	38,460	66.4	57.69	74.05
Form 2	250	33,482	62.8	57.27	68.06
Form 3	221	35,459	65.4	59.34	71.01
Form 4	253	36,842	68.3	60.16	75.49
Form 5	179	35,187	62.6	55.36	69.33

Table 3.5.8: Prevalence of handwashing before eating in the past 30 days among Form 1 to Form 5 students in Johor, 2017

Socio-demographic characteristics	Never or rarely wash hands before eating					Sometimes				
	Unweighted Count	Estimated Population	Prevalence (%)	95% CI		Unweighted Count	Estimated Population	Prevalence (%)	95% CI	
				Lower	Upper				Lower	Upper
JOHOR	75	12,105	4.4	2.82	6.80	179	27,640	10.0	7.87	12.72
Locality of school										
Urban	62	9,975	5.1	3.05	8.37	144	21,924	11.2	8.55	14.49
Rural	13	2,130	2.7	1.97	3.66	35	5,715	7.2	5.03	10.22
Sex										
Male	43	7,719	5.6	3.32	9.40	73	13,286	9.7	7.79	12.02
Female	32	4,386	3.2	1.91	5.21	106	14,354	10.4	7.08	14.95
Form										
Form 1	16	2,887	5.0	2.65	9.19	39	6,754	11.7	7.92	16.83
Form 2	23	3,028	5.7	2.41	12.79	50	6,504	12.2	8.35	17.50
Form 3	14	2,133	4.0	2.23	6.93	26	4,102	7.6	4.68	12.16
Form 4	10	1,846	3.4	1.45	7.84	41	5,799	10.7	6.91	16.30
Form 5	12	2,210	3.9	2.06	7.37	23	4,480	8.0	5.73	10.98

Table 3.5.8: Prevalence of handwashing before eating in the past 30 days among Form 1 to Form 5 students in Johor, 2017 (cont.)

Socio-demographic characteristics	Most of the times and Always				
	Unweighted Count	Estimated Population	Prevalence (%)	95% CI	
				Lower	Upper
JOHOR	1,475	235,643	85.6	81.29	89.00
Locality of school					
Urban	1,039	164,170	83.7	78.33	87.99
Rural	436	71,473	90.1	86.70	92.72
Sex					
Male	630	115,975	84.7	79.29	88.84
Female	845	119,667	86.5	80.67	90.71
Form					
Form 1	286	48,319	83.4	76.72	88.40
Form 2	326	43,766	82.1	71.87	89.19
Form 3	296	47,655	88.4	82.84	92.37
Form 4	315	46,389	85.9	80.31	90.03
Form 5	252	49,513	88.1	83.58	91.49

Table 3.5.9: Prevalence of handwashing after using the toilet in the past 30 days among Form 1 to Form 5 students in Johor, 2017

Socio-demographic characteristics	Never or rarely wash hands after using the toilet					Sometimes				
	Unweighted Count	Estimated Population	Prevalence (%)	95% CI		Unweighted Count	Estimated Population	Prevalence (%)	95% CI	
				Lower	Upper				Lower	Upper
JOHOR	50	7,714	2.8	2.10	3.72	133	21,449	7.8	6.26	9.63
Locality of school										
Urban	33	5,048	2.6	1.82	3.62	88	14,399	7.3	5.46	9.81
Rural	17	2,666	3.4	2.10	5.33	45	7,050	8.9	7.11	11.01
Sex										
Male	27	4,610	3.4	2.47	4.55	51	9,712	7.1	4.87	10.18
Female	23	3,104	2.2	1.55	3.23	82	11,736	8.5	6.77	10.57
Form										
Form 1	13	2,205	3.8	1.77	8.00	35	5,959	10.3	6.60	15.66
Form 2	19	2,588	4.9	3.34	7.04	30	3,953	7.4	5.11	10.70
Form 3	6	966	1.8	0.81	3.87	21	3,383	6.2	3.47	10.96
Form 4	10	1,478	2.7	1.53	4.84	27	4,124	7.6	4.76	12.01
Form 5	2	477	0.8	0.22	3.24	20	4,029	7.2	4.51	11.21

Table 3.5.9: Prevalence of handwashing after using the toilet in the past 30 days among Form 1 to Form 5 students in Johor, 2017 (cont.)

Socio-demographic characteristics	Most of the time and Always				
	Unweighted Count	Estimated Population	Prevalence (%)	95% CI Lower	95% CI Upper
JOHOR	1,547	246,439	89.4	87.10	91.36
Locality of school					
Urban	1,124	176,681	90.1	87.25	92.35
Rural	423	69,758	87.8	83.98	90.77
Sex					
Male	669	122,825	89.6	85.97	92.31
Female	878	123,614	89.3	86.78	91.36
Form					
Form 1	293	49,796	85.9	79.30	90.66
Form 2	349	46,648	87.7	83.81	90.76
Form 3	311	49,865	92.0	87.75	94.83
Form 4	329	48,432	89.6	84.50	93.20
Form 5	265	51,698	92.0	88.11	94.67

3.6 Internet Use and Addiction

Contributors: S. Maria Awaluddin, Normala Ibrahim, Chan Ying Ying, Rimah Melati Abd. Ghani, Amal Shamsudin, Wan Salwina Wan Ismail, Norharlina Bahar, Saidatul Norbaya Buang, Nik Rubiah Nik Abd. Rashid.

3.6.1 Introduction

Internet use has grown rapidly in Malaysia parallel with an increasing number of households having computers and mobile gadgets that are linked to internet access. It has been reported that the increase in internet use is more pronounced in urban than in rural areas. Major online activities include information seeking for educational, vocational and socializing purposes. Internet usage affects both adult and adolescent age groups equally compared to older age group.¹ Internet usage could be harmful to adolescents when it is not monitored or supervised by an adult. Uncontrolled use of internet has been linked to psychosocial problems such as addiction to online gaming, pornographic websites and social networking sites, not to mention exposing the adolescents to cyber-bullying environments. Excessive internet use and cyber bullying may in turn further worsen the mental health condition of these adolescents and predisposed them to depression, anxiety and stress.²

The terminology of internet addiction to describe excessive internet use has been debated and various tools have been developed to measure the internet addiction. The most widely used tool was the Internet Addiction Test (IAT) developed by Kimberly S. Young.³ The questionnaire consists of 20 items measured using a 5-point Likert scale (1=rarely, 2=occasionally, 3=frequently, 4=often and 5=always) with the highest possible score of 100. Internet addiction is defined as a person scoring 43 points or above. This questionnaire was validated in Bahasa Malaysia by Chong Guan et al in 2012 and was named Malay Version Internet Addiction Test (MVIAT).⁴ The (MVIAT) showed a good internal consistency (Cronbach's $\alpha = 0.91$, $P < 0.001$), parallel reliability (intra-class coefficient correlation=0.88, $P < 0.001$) and concurrent validity with Compulsive Internet Use Scale (Pearson's correlation= 0.84, $P < 0.00$).

3.6.2 Objectives

- i. To identify the prevalence of internet use among adolescents in the past 30 days
- ii. To identify the prevalence of internet addiction among adolescents
- iii. To identify the type of devices used by internet users and addicts

3.6.3 Variable Definitions

- **Internet Use:** Internet access using any internet connecting devices for the past 30 days.
- **Internet Addiction:** Using a self-administered 20-item Malay Version Internet Addiction Test (MVIAT) questionnaire, respondents with a total score of 43 or above (all items answered) were defined as having internet addiction.

3.6.4 Findings

There were 1731 secondary school students in Johor who responded to this module, of which 120 students did not surf the internet for the past 30 days and 1576 students completed the MVIAT. The overall prevalence of internet use was 93.0% (95% CI: 90.05, 95.13) with an estimated projection to 256,415 school-going adolescents. In terms of school locality, there was no significant difference in internet usage between students studying in urban areas (92.8%; 95% CI: 88.68, 95.52) and students from rural areas (93.5%; 95% CI: 90.26, 95.66). There were 91.5% (95% CI: 87.43, 94.39) male students and 94.4% (95% CI: 91.69, 96.33) female students who used internet in the past 30 days in Johor. Students from higher forms were more likely to use internet in the past 30 days (**Table 3.6.1**).

The prevalence of internet addiction among school-going adolescents in Johor was 33.9% (95% CI: 30.56, 37.34). In terms of school locality, there was no significant difference in internet addiction between students studying in urban areas (35.3%; 95% CI: 31.99, 38.70) and students from rural areas (30.4%; 95% CI: 24.20, 37.38). There were 36.6% (95% CI: 32.42, 40.96) male students and 31.2% (95% CI: 27.55, 35.05) female students with positive MVIAT. Form 4 students were observed as having the highest prevalence of internet addiction in Johor (**Table 3.6.2**). Smartphone was the most prevalent device used by the internet users and addicts followed by the computer, laptop or notebook group and tablet or iPad group (**Table 3.6.3 & Table 3.6.4**).

3.6.5 Discussion/ Conclusion

The prevalence of internet usage and internet addiction in Johor was significantly higher than the national prevalence as Johor was considered among the urbanised states in Malaysia. Students in higher forms were observed as having a higher prevalence of internet usage, however there were no significant differences in terms of school locality and sex of the respondents. The prevalence of internet addiction was significantly higher in Form 4 students compared to Form 1 and Form 2 students. However, no significant difference was observed in prevalence of internet addiction by school locality and sex of the respondents. Majority of the students used smartphones to surf the internet.

3.6.5 Discussion/ Conclusion

The prevalence of internet usage and internet addiction in Pahang does not differ significantly compared to the national prevalence. Students in higher forms were observed as having a higher prevalence of internet usage and internet addiction. Most of the students used smartphones to surf the internet.

3.6.6 Recommendations

1. The internet addiction problems among Malaysian students is considered as new health risk behaviour that needs further in-depth understanding on internet accessibility, usage, difference in type of content viewed by age, gender and ethnicity, psychosocial factors and role of significant adult supervision, peers and environment to assist in designing specific strategies in prevention program.
2. As internet use and addiction increase with age, it is important that adolescents develop self-awareness on sign of internet addiction and develop self-control against excessive internet usage. It is proposed that the component of self-awareness and control against excessive internet usage is to be integrated into the 'Click Wisely Program' which was introduced by the Malaysian Communication Multimedia Commission (MCMC).
3. Parents should be made aware on dangers of excessive internet usage by secondary students and its detrimental effects on students' health and social development. It is

proposed that development of health-related information on the internet addiction should be designed specifically for students and parents. To ensure wide circulation of the messages, it is recommended that the Ministry of Health to collaborate with MCMC in disseminating the information through MCMC social network and Malaysian ICT volunteer (MIV) programs.

4. Smart partnerships with various agencies (governments, NGOs and private sectors) need to be enhanced to disseminate greater awareness on dangers of internet addiction and safe usage of internet to children, adolescents, parents, teachers and the community at large to assist in promotive, preventive and early intervention of internet addiction.

3.6.7 References

1. Malaysian Communications and Multimedia Commission. Internet Users Survey 2016 [Internet]. 2016 [cited 2017 Dec 8]. Available from: <https://www.mcmc.gov.my/resources/statistics/internet-users-survey>
2. Gholamian B, Shahnazi H, Hassanzadeh A. The Prevalence of Internet Addiction and its Association with Depression, Anxiety, and Stress, among High-School Students. *Int J Pediatr.* 2017;5(4):4763–4770.
3. Young KS. Internet addiction: The emergence of a new clinical disorder. *Cyberpsychol Behav.* 1998;1(3):237–44.
4. Chong Guan N, Isa SM, Hashim AH, Pillai SK, Harbajan Singh MK. Validity of the Malay Version of the Internet Addiction Test: A Study on a Group of Medical Students in Malaysia. *Asia Pac J Public Health.* 2012;27(2):NP2210-NP2219.

Table 3.6.1 : Prevalence of internet use in the past 30 days among Form 1 to Form 5 students in Johor, 2017

Socio-demographic characteristics	Internet Users				
	Unweighted Count	Estimated Population	Prevalence (%)	95% CI	
				Lower	Upper
JOHOR	1,611	256,415	93.0	90.05	95.13
Locality of school					
Urban	1,159	182,136	92.8	88.68	95.52
Rural	452	74,278	93.5	90.26	95.66
Sex					
Male	681	125,546	91.5	87.43	94.39
Female	930	130,869	94.4	91.69	96.33
Form					
Form 1	301	50,523	87.2	80.88	91.60
Form 2	346	45,990	86.3	73.90	93.33
Form 3	326	52,094	96.1	90.77	98.40
Form 4	356	52,562	97.3	93.37	98.91
Form 5	282	55,246	98.3	95.41	99.38

Table 3.6.2: Prevalence of positive Malay Version Internet Addiction Test (MVIAT) among Form 1 to Form 5 students in Johor, 2017

Socio-demographic characteristics	Positive MVIAT				
	Unweighted Count	Estimated Population	Prevalence (%)	95% CI	
				Lower	Upper
JOHOR	580	93372	33.9	30.56	37.34
Locality of school					
Urban	432	69220	35.3	31.99	38.70
Rural	148	24152	30.4	24.20	37.38
Sex					
Male	270	50176	36.6	32.42	40.96
Female	310	43196	31.2	27.55	35.05
Form					
Form 1	71	12652	21.8	14.52	31.46
Form 2	120	15621	29.3	22.27	37.50
Form 3	122	19940	36.8	28.19	46.29
Form 4	160	23630	43.7	38.47	49.14
Form 5	107	21529	38.3	30.85	46.36

Table 3.6.3: Percentage of reported devices used by internet users among Form 1 to Form 5 students in Johor, 2017

Type of Devices	Unweighted Count	Percentage (%)	95% CI	
			Lower	Upper
Smartphone	1,531	95.67	94.60	96.54
Computer, Laptop, Notebook	959	60.14	54.86	65.19
Tablet, Ipad use	412	25.41	21.74	29.48

Table 3.6.4: Percentage of reported devices used by internet addicts among Form 1 to Form 5 students in Johor, 2017

Type of Devices	Unweighted Count	Percentage (%)	95% CI	
			Lower	Upper
Smartphone	562	96.8	94.72	98.11
Computer, Laptop, Notebook	370	64.3	60.00	68.29
Tablet, Ipad use	181	31.7	25.69	38.45

3.7 Mental Health Problems

Contributors: Noor Ani Ahmad, Muslimah Yusof, Mohamad Aznuddin Abd Razak, Noraida Mohamad Kasim, Fazly Azry Abdul Aziz, Rasidah Jamaluddin, Nurashikin Ibrahim.

3.7.1 Introduction

The World Health Organisation (WHO) estimates that one in five adolescents and children suffer from mental health problem and half of all mental illnesses begin by the age of 14.¹ In Malaysia, the National Health and Morbidity Survey (NHMS) 2015 reported prevalence of mental health problems among 16 to 19 years old of 34.7%, while among 10-15 years the prevalence was 11.4%.² In particular, WHO reported that suicide is the second leading cause of death among those youth.³ In 2011, NHMS observed that 2.4% of youth 16-24 years reported suicidal ideation.⁴

3.7.2 Objectives

- i. To identify the prevalence of loneliness in the past 12 months
- ii. To identify the prevalence of inability to sleep due to worry in the past 12 months
- iii. To identify the prevalence of suicidal ideation in the past 12 months
- iv. To identify the prevalence of suicidal plan in the past 12 months
- v. To identify the prevalence of suicidal attempt in the past 12 months
- vi. To identify the prevalence of not having close friend

3.7.3 Variable definitions

- **Lonely “most of the time or always”**: Responded either “most of the time” or “always” for felt lonely during past 12 months prior to the survey.
- **Unable to sleep “most of the time or always” due to worry**: Responded either “most of the time” or “always” for being worried about something that he/she could not sleep at night during past 12 months prior to the survey.
- **Suicidal ideation**: ever seriously considered attempting suicide in the past 12 months prior to the survey.
- **Suicidal plan**: made a plan of attempted suicide in the past 12 months prior to the survey.
- **Suicidal attempt**: attempted suicide at least once in the past 12 months prior to the survey.

- **No close friend:** Do not have any close friend

3.7.4 Findings

Overall, 8.9% (95%CI: 7.06, 11.07) of secondary school students in Johor reported that they felt lonely “most of the time or always” (**Table 3.7.1**). A total of 6.5% (95%CI: 4.71, 8.80) reported being unable to sleep “most of the time or always” due to worry (**Table 3.7.2**). In the past 12 months prior to the survey, suicidal ideation, plan and attempt, were reported by 7.5% (95%CI: 5.87, 9.55), 5.7% (95%CI: 4.24, 7.61), and 4.6% (95%CI: 3.00, 7.02) of students, respectively (**Table 3.7.3, Table 3.7.4 & Table 3.7.5**). **Table 3.7.6** showed that 3.1% (95%CI: 2.20, 4.41) of the students had no close friend. There is no significant difference in the prevalence by locality, sex or form for all these conditions.

3.7.5 Discussion/ Conclusion

The prevalence of being lonely among secondary school students in Johor increased from 7.1% in the Johor GSHS 2012⁵ to 8.9% in this current survey. There was no difference in the prevalence of “inability to sleep due to worry” in the two surveys (6.4% vs 6.5%). Significant improvement in the prevalence of suicidal behaviours was observed among the secondary school students in Johor as compared to the prevalence in 2012. Suicidal ideation, plan and attempts decreased from 11.3% to 7.5%, 8.1% to 5.7%, and 9.7% to 4.6%, respectively.⁵ The 2017 figures were also lower as compared to the national figures. In addition to these, less students reported having no close friend as compared to previous the Johor GSHS 2012 (3.1% vs 3.9%).⁵

3.7.6 Recommendations

1. Screening of at-risk students by School Health Teams and referral for further management
2. Holistic intervention programmes targeted to students at risk of suicide
3. Strengthen students coping skills and resilience through interactive health promotion activities

3.7.7 References

1. Mental Health. Child and Adolescent mental health.
http://www.who.int/mental_health/maternal-child/child_adolescent/en/
2. Institute for Public Health (IPH). 2015. National Health and Morbidity Survey, NHMS 2015. Ministry of Health Malaysia
3. WHO Fact Sheet. Suicide. www.who.int/mediacentre/factsheets/fs398/en/. Updated on March 2017.
4. Institute for Public Health (IPH). 2011. National Health and Morbidity Survey, NHMS 2011. Ministry of Health Malaysia
5. Institute for Public Health (IPH). 2012. National Health and Morbidity Survey, NHMS 2012: Johor Global School-based Student Health Survey 2012. Kuala Lumpur Ministry of Health Malaysia.

Table 3.7.1: Prevalence of loneliness "most of the time or always" in the past 12 months among Form 1 to Form 5 students in Johor, 2017

Socio-demographic characteristics	Unweighted count	Estimated population	Prevalence (%)	95% CI	
				Lower	Upper
JOHOR	158	24,420	8.9	7.06	11.07
Locality of school					
Urban	110	16,845	8.6	6.70	10.95
Rural	48	7,575	9.5	5.92	15.00
Sex					
Male	58	10,535	7.7	5.30	11.03
Female	100	13,885	10.0	8.18	12.22
Form					
Form 1	18	2,933	5.1	2.92	8.64
Form 2	38	5,040	9.5	6.36	13.84
Form 3	40	6,262	11.6	7.30	17.81
Form 4	38	5,572	10.3	6.81	15.40
Form 5	24	4,612	8.2	4.92	13.39

Table 3.7.2: Prevalence of inability to sleep "most of time or always" due to worry in the past 12 months among Form 1 to Form 5 students in Johor, 2017

Socio-demographic characteristics	Unweighted count	Estimated population	Prevalence (%)	95% CI	
				Lower	Upper
JOHOR	111	17,813	6.5	4.71	8.80
Locality of school					
Urban	79	12,589	6.4	4.33	9.40
Rural	32	5,223	6.6	3.90	10.87
Sex					
Male	44	8,138	5.9	3.56	9.73
Female	67	9,674	7.0	5.43	8.94
Form					
Form 1	20	3,426	5.9	3.35	10.22
Form 2	18	2,534	4.8	2.35	9.39
Form 3	27	4,306	7.9	4.80	12.87
Form 4	30	4,438	8.2	5.24	12.65
Form 5	16	3,109	5.5	2.99	10.03

Table 3.7.3: Prevalence of suicidal ideation in the past 12 months among Form 1 to Form 5 students in Johor, 2017

Socio-demographic characteristics	Unweighted count	Estimated population	Prevalence (%)	95% CI	
				Lower	Upper
JOHOR	135	20,659	7.5	5.87	9.55
Locality of school					
Urban	102	15,339	7.8	5.85	10.40
Rural	33	5,320	6.7	4.36	10.18
Sex					
Male	50	8,813	6.4	4.93	8.33
Female	85	11,846	8.6	6.46	11.30
Form					
Form 1	33	5,635	9.7	6.03	15.30
Form 2	34	4,453	8.4	5.48	12.61
Form 3	26	4,102	7.6	4.39	12.79
Form 4	31	4,501	8.4	6.25	11.08
Form 5	11	1,969	3.5	1.16	10.13

Table 3.7.4: Prevalence of suicidal plan in the past 12 months among Form 1 to Form 5 students in Johor, 2017

Socio-demographic characteristics	Unweighted count	Estimated population	Prevalence (%)	95% CI	
				Lower	Upper
JOHOR	101	15,681	5.7	4.24	7.61
Locality of school					
Urban	78	12,106	6.2	4.38	8.62
Rural	23	3,575	4.5	2.80	7.21
Sex					
Male	40	7,091	5.2	3.43	7.74
Female	61	8,591	6.2	4.74	8.07
Form					
Form 1	21	3,525	6.1	4.10	8.93
Form 2	33	4,534	8.5	4.65	15.05
Form 3	14	2,144	4.0	2.31	6.69
Form 4	21	3,159	5.8	3.24	10.32
Form 5	12	2,320	4.1	1.79	9.32

Table 3.7.5: Prevalence of suicidal attempt "at least once" in the past 12 months among Form 1 to Form 5 students in Johor, 2017

Socio-demographic characteristics	Unweighted count	Estimated population	Prevalence (%)	95% CI	
				Lower	Upper
JOHOR	81	12,699	4.6	3.00	7.02
Locality of school					
Urban	57	8,861	4.5	2.56	7.84
Rural	24	3,837	4.8	2.77	8.35
Sex					
Male	31	5,484	4.0	2.33	6.80
Female	50	7,215	5.2	3.44	7.82
Form/ Class					
Form 1	23	3,831	6.6	3.82	11.19
Form 2	25	3,507	6.6	3.33	12.60
Form 3	12	1,884	3.5	1.98	6.04
Form 4	12	1,827	3.4	2.10	5.40
Form 5	9	1,650	2.9	1.37	6.23

Table 3.7.6: Prevalence of not having any close friend among Form 1 to Form 5 students Johor, 2017

Socio-demographic characteristics	Unweighted count	Estimated population	Prevalence (%)	95% CI	
				Lower	Upper
JOHOR	53	8,569	3.1	2.20	4.41
Locality of school					
Urban	42	6,571	3.4	2.45	4.59
Rural	11	1,998	2.5	0.87	7.09
Sex					
Male	29	5,361	3.9	2.64	5.78
Female	24	3,208	2.3	1.30	4.10
Form/ Class					
Form 1	12	2,197	3.8	2.52	5.69
Form 2	10	1,336	2.5	1.14	5.51
Form 3	9	1,513	2.8	1.23	6.20
Form 4	13	1,786	3.3	1.51	7.09
Form 5	9	1,737	3.1	0.95	9.69

3.8 Physical Activity

Contributors: Lim Kuang Kuay, Mohd Azahadi Omar, Chan Ying Ying, Chandrika A/p Jeevananthan, Azli Baharudin @Shaharuddin, Nazirah Alias

3.8.1 Introduction

Physical inactivity is a public health problem worldwide and is the fourth leading cause of death worldwide.¹ It is an important contributor to major non-communicable diseases.^{2,3} Sufficient physical activity has substantial health benefits for children and adolescents in terms of improving cardiovascular health⁴, mental health⁵ and academic performance⁶. Despite these established benefits, a substantial proportion of young people fail to meet physical activity guidelines.

3.8.2 Objectives

1. To identify the prevalence of being physically active for a total of at least 60 minutes daily for five days or more in the past seven days,
2. To identify the prevalence of spending at least three hours in sitting activities in a typical or usual day

3.8.3 Variable Definitions

- **Physical activity:** any activity that increases the heart rate and makes one breathe hard. Examples of physical activities include sports, playing with friends, walking to school, running, fast walking, biking or dancing.
- **Physically active:** physically active for at least 60 minutes per day, for a minimum of five days per week (sum of all the time spent in any kind of physical activity each day).

3.8.4 Findings

There were 1,727 secondary school students who responded to this module. The prevalence of being physically active for a total of at least 60 minutes daily for five days or more in the past seven days was 25.3% (95% CI: 23.30, 27.50), with an estimated projection to 69,680 school-going adolescents. In terms of school locality, it was higher in urban areas [25.5% (95% CI: 23.14, 28.12)] compared to rural areas [24.8% (95% CI: 21.15, 28.94)]. By sex, the prevalence was significantly

higher in males [32.8% (95% CI: 29.47, 36.32)] than in females [18.0% (95% CI: 15.66, 20.55)]. By form, Form 4 had the highest prevalence [28.2% (95% CI: 23.82, 33.09)], followed by Form 1 [25.8% (95% CI: 21.35, 30.73)] and Form 3 [25.3% (95% CI: 20.97, 30.28)].

Regarding sitting activities, the prevalence of spending at least three hours in a typical or usual day in sitting activities was 60.7% (95% CI: 58.36, 62.93). In terms of school locality, it was significantly higher in urban areas [64.1% (95% CI: 61.41, 66.74)] compared to rural areas [52.2% (95% CI: 47.73, 56.57)]. By sex, the prevalence among females [64.3% (95% CI: 61.31, 67.28)] was significantly higher than males [57.0% (95% CI: 53.46, 60.40)]. By form, Form 5 had the highest prevalence [69.2% (95% CI: 63.57, 74.34)], followed by Form 4 [67.4% (95% CI: 62.35, 72.04)] and Form 3 [67.0% (95% CI: 61.82, 71.85)].

3.8.5 Discussion/ Conclusion

The prevalence of school-going adolescents being physically active for a total of at least 60 minutes daily for five days or more in the past 7 days in Johor [25.3% (95% CI: 23.30, 27.50)] was significantly higher than the national prevalence [19.8% (95% CI: 19.21, 20.40)] and the previous Johor GSHS 2012⁷ [19.9% (95% CI: 16.64, 23.70)]. Male students were more active than female students. However, there were no significant differences in terms of locality and form. The prevalence of students who spent at least three hours in sitting activities in a typical or usual day [60.7% (95% CI: 58.36, 62.93)] was significantly higher than the national prevalence [50.1% (95% CI: 49.38, 50.85)] and the previous Johor GSHS 2012⁷ [49.2% (95% CI: 46.39, 52.04)]. It was significantly higher in Form 5 compared to Form 4 and Form 3 students. However, there was no significant difference in terms of sex of the students.

3.8.6 Recommendations

Schools represent an important setting for promoting physical activity as the adolescents spend approximately half of their waking day at school. Development of creative and innovative “school-based interventions” such as engaging social media to promote healthy lifestyles particularly on increasing physical activity need to be explored.

3.8.7 References

1. Cora LC, Estelle VL, Shigeru I, Jasem RA, Grit L, Sonja K. The pandemic of physical inactivity: global action for public health. *The Lancet*. 2012; 380(9838): 294-305.
2. Farid AWG, Mohamed RA, Ayesha AR, Mohd I. The Prevalence of Physical Activity and Sedentary Lifestyle among Adolescents in Palestine. *Can. J. Basic Appl. Sci.* 2013;03 (01):135-144.
3. Sharif R, Chong KH, Zakaria NH, Ong ML, Reilly JJ, Wong JE, Saad HA, Poh BK. Results From Malaysia's 2016 Report Card on Physical Activity for Children and Adolescents. *J Phys Act Health*. 2016; 13(11): S201-S205.
4. Ekelund U, Luan J, Sherar LB, Esliger DW, Griew P, Cooper A. Moderate to vigorous physical activity and sedentary time and cardiometabolic risk factors in children and adolescents. *JAMA*. 2012;307: 704–712.
5. Biddle SJH, Asare M. Physical activity and mental health in children and adolescents: a review of reviews. *Br J Sports Med*. 2011; 45: 886–895.
6. Singh A, Uijtdewilligen L, Twisk JWR, van Mechelen W, Chinapaw MJM. Physical activity and performance at school: a systematic review of the literature including a methodological quality assessment. *Arch Pediatr Adolesc Med*. 2012; 166: 49–55.
7. Institute for Public Health (IPH). *The National Health and Morbidity Survey: Malaysia Global School-based Student Health Survey 2012*. 2012. Kuala Lumpur: Ministry of Health Malaysia.

Table 3.8.1 : Prevalence of being physically active (60 minutes daily) for a total of at least 5 days or more among Form 1 to Form 5 students in Johor, 2017

Socio-demographic characteristics	Physically Active for at least 5 days or more				
	Unweighted Count	Estimated Population	Prevalence (%)	95% CI	
				Lower	Upper
JOHOR	417	69,680	25.3	23.30	27.50
Locality of school					
Urban	298	50,038	25.5	23.14	28.12
Rural	119	19,643	24.8	21.15	28.94
Sex					
Male	242	44,834	32.8	29.47	36.32
Female	175	24,846	18.0	15.66	20.55
Form					
Form 1	85	14,873	25.8	21.35	30.73
Form 2	87	11,782	22.2	18.32	26.54
Form 3	83	13,740	25.3	20.97	30.28
Form 4	95	15,252	28.2	23.82	33.09
Form 5	67	14,034	25.2	20.31	30.74

Table 3.8.2: Prevalence of spending at least 3 hours in sitting activities Form 1 to Form 5 students in Johor, 2017

Socio-demographic characteristics	spending at least 3 hours on sedentary activities				
	Unweighted Count	Estimated Population	Prevalence (%)	95% CI	
				Lower	Upper
JOHOR	1,057	166,990	60.7	58.36	62.93
Locality of school					
Urban	803	125,529	64.1	61.41	66.74
Rural	254	41,461	52.2	47.73	56.57
Sex					
Male	421	78,123	57.0	53.46	60.40
Female	636	88,867	64.3	61.31	67.28
Form					
Form 1	161	26,986	46.7	41.46	51.96
Form 2	218	28,351	53.5	48.56	58.40
Form 3	230	36,338	67.0	61.82	71.85
Form 4	248	36,411	67.4	62.35	72.04
Form 5	200	38,903	69.2	63.57	74.34

3.9. Protective Factors

Contributors: Faizah Paiwai, S Maria Awaluddin, Nur Azna Mahmud, Noor Aliza Lodz, Norzawati Yoep

3.9.1 Introduction

According to Centre for Disease Control and Prevention (CDC), protective factors are individual or environmental characteristics, conditions, or behaviours that reduce the effects of stressful life events.¹ Important protective factors such as parental attachment and peer support especially for adolescence could influence their attitude and behaviour. Research has shown that adolescent attachment and relationships with parents and peers as well as schools they attend is a predictor of their psychological well-being.² Protective factor could help avoid unhealthy behaviours such as violence, sexual risk behaviours, tobacco use and drug abuse.³ In order to measure the level of protective factors among school students, truancy is seen as an indicator, which is monitored by lower prevalence, as truancy often acts as the initial behaviour that can lead to other risky behaviour.⁴

3.9.2 Objectives

To identify prevalence of:

- i. Truancy in the past 30 days
- ii. Peer support in the past 30 days
- iii. Parental or guardian supervision in the past 30 days
- iv. Parental or guardian connectedness in the past 30 days
- v. Parental or guardian bonding in the past 30 days,
- vi. Parental or guardian respect for privacy in the past 30 days,

3.9.3 Variable Definitions

- **Truancy:** missed class or school without permission for at least one day in the past 30 days. (This variable is monitored with lower prevalence to define as protective factors)
- **Peer support:** students in their school were kind and helpful most of the time or always during the past 30 days.

- **Parental or guardian supervision:** parents or guardians had always or most of the time, checked to see if their homework was done in the past 30 days.
- **Parental or guardian connectedness:** parents or guardians had always or most of the time, understood their problems and worries in the past 30 days.
- **Parental or guardian bonding:** parents or guardians had always or most of the time, really knew what they were doing with their free time in the past 30 days.
- **Parental or guardian respect for privacy:** parents or guardians had never or rarely gone through their things without their approval in the past 30 days.

3.9.4 Findings

In Johor, the overall prevalence of truancy among students in the past 30 days was 24.2% (95% CI: 18.41, 31.19) of which males were 29.1% (95% CI: 21.81, 37.74) and females were 19.4% (95% CI: 14.06, 26.06). Prevalence of truancy according to form was 15.7% (95% CI: 11.38, 21.37), 23.9% (95% CI: 17.65, 31.51), 17.4% (95% CI: 10.43, 27.49), 32.4% (95% CI: 23.76, 42.33) and 32.1% (95% CI: 21.15, 45.52) from Form 1 to Form 5 respectively. In terms of school locality, prevalence of truancy in rural areas was 22.2% (95% CI: 17.71, 27.48) and urban areas was 25.0% (95% CI: 17.39, 34.67) (**Table 3.9.1**).

Overall prevalence of having peer support among students in Johor was 49.9% (95% CI: 44.43, 55.28); of which 43.4% (95% CI: 37.04, 49.96) in males, 56.2% (95% CI: 51.35, 61.02) in females (**Table 3.9.2**). Total prevalence of having parental or guardian supervision was 12.3% (95% CI: 10.53, 14.37) which was 15.0% (95% CI: 12.56, 17.90) in males, 9.6% (95% CI: 7.76, 11.90) in females (**Table 3.9.3**). The overall prevalence of having parental or guardian connectedness was 34.9% (95% CI: 31.05, 38.88) of which 35.7% (95% CI: 30.85, 40.83) in males and 34.0% (95% CI: 30.07, 38.27) in females. In terms of school locality, having parental or guardian connectedness was 34.1% (95% CI: 29.41, 39.03) in urban areas and 36.9% (95% CI: 31.19, 42.93) in rural areas (**Table 3.9.4**). Overall prevalence of having parental bonding was 51.0% (95% CI: 46.88, 55.13) which was 49.1% (95% CI: 43.73, 54.44) in males and 52.9% (95% CI: 48.36, 57.47) in females (**Table 3.9.5**). A total 77.6% (95% CI: 74.58, 80.42) of students reported having parental respect for privacy, 72.2% (95% CI: 69.94, 74.37) in males and 83.0% (95% CI: 78.35, 86.84) in females (**Table 3.9.6**).

3.9.5 Discussions/ Conclusion

Prevalence of truancy and protective factors (peer support, parental supervision, parental connectedness, parental bonding and parental respect) among students in Johor within the past 30 days showed no significant difference from the national prevalence. There were also no significant difference in the prevalence of truancy and protective factors reported between the national AHS 2017 and Johor GSHS 2012.⁵

3.9.6 Recommendations

1. Monitoring attendance closely by participation of schools, parent and local organizations through enforcement of mandatory attendance law allows identification of at risk and truancy behaviour among school students.
2. Positive reinforcement which focuses on the positive points of behavior will encourage children to improve peer support, self-control and respect for others.
3. Establishment of school programs that need parent's supervision will help in improving the parenting skills especially in parental attachment.
4. Development of interventions that strengthen the protective factors among school students is important and more effective in reducing risk in order to improve the outcomes experienced by the students

3.9.7 References

1. Centers for Disease Control and Prevention. School Connectedness: Strategies for Increasing Protective Factors Among Youth. Atlanta, GA: U.S. Department of Health & Human Services; 2009. <https://www.cdc.gov/healthyyouth/protective/>. Accessed on 9 August 2017
2. Oldfield J, Humphrey N, Hebron J. The role of parental and peer attachment relationships and school connectedness in predicting adolescent mental health outcomes. *Child and Adolescent Mental Health*. 2016 Feb 1;21(1):21-9.
3. Ferreira M, Gaspar de Matos M, Diniz JA. The Influenced of Risk and Protective Factors on Tobacco, Alcohol and Drugs in Adolescence. *Risk*. 2013; 6.

4. Rocque M, Jennings WG, Piquero AR, Ozkan T, Farrington DP. The importance of school attendance: Findings from the Cambridge Study in Delinquent Development on the Life-Course Effects of Truancy. *Crime & Delinquency*. 2017 May; 63(5):592-612.
5. Institute for Public Health (IPH) 2012. The National Health and Morbidity Survey: Malaysia Global School-based Student Health Survey 2012. Kuala Lumpur: Ministry of Health Malaysia.

Table 3.9.1 : Prevalence of truancy in the past 30 days among Form 1 to Form 5 students in Johor, 2017

Socio-demographic Characteristics	Truancy				
	Unweighted Count	Estimated Population	Prevalence (%)	95% CI	
				Lower	Upper
JOHOR	401	66,742	24.2	18.41	31.19
Locality of school					
Urban	297	49,088	25.0	17.39	34.67
Rural	104	17,654	22.2	17.71	27.48
Sex					
Male	211	39,965	29.1	21.81	37.74
Female	190	26,777	19.4	14.06	26.06
Form					
Form 1	51	9,125	15.7	11.38	21.37
Form 2	94	12,707	23.9	17.65	31.51
Form 3	57	9,413	17.4	10.43	27.49
Form 4	115	17,440	32.4	23.76	42.33
Form 5	84	18,057	32.1	21.15	45.52

Table 3.9.2 : Prevalence of peer support in the past 30 days among Form 1 to Form 5 students in Johor, 2017.

Socio-demographic Characteristics	Most of the time or always				
	Unweighted Count	Estimated Population	Prevalence (%)	95% CI	
				Lower	Upper
JOHOR	872	137,358	49.9	44.43	55.28
Locality of school					
Urban	633	98,254	50.1	43.48	56.66
Rural	239	39,104	49.3	39.96	58.74
Sex					
Male	321	59,420	43.4	37.04	49.96
Female	551	77,938	56.2	51.35	61.02
Form					
Form 1	189	31,288	54.2	47.23	60.95
Form 2	169	22,242	41.7	33.81	50.10
Form 3	156	24,130	44.5	35.53	53.86
Form 4	196	29,005	53.7	44.14	62.95
Form 5	162	30,693	54.6	41.32	67.27

Table 3.9.3 : Prevalence of parental or guardian supervision in the past 30 days among Form 1 to Form 5 students in Johor, 2017

Socio-demographic Characteristics	Most of the time or always				
	Unweighted Count	Estimated Population	Prevalence (%)	95% CI	
				Lower	Upper
JOHOR	209	33,967	12.3	10.53	14.37
Locality of school					
Urban	150	24,466	12.5	10.38	14.91
Rural	59	9,501	12.0	8.79	16.06
Sex					
Male	114	20,619	15.0	12.56	17.90
Female	95	13,347	9.6	7.76	11.90
Form					
Form 1	71	12,481	21.5	17.61	26.06
Form 2	55	7,516	14.1	10.43	18.79
Form 3	36	5,894	10.9	7.02	16.46
Form 4	31	4,723	8.7	5.82	12.92
Form 5	16	3,354	6.0	3.97	8.88

Table 3.9.4: Prevalence of parental or guardian connectedness in the past 30 days among Form 1 to Form 5 students in Johor, 2017

Socio-demographic Characteristics	Most of the time or always				
	Unweighted Count	Estimated Population	Prevalence (%)	95% CI	
				Lower	Upper
JOHOR	601	96,067	34.9	31.05	38.88
Locality of school					
Urban	425	66,826	34.1	29.41	39.03
Rural	176	29,241	36.9	31.19	42.93
Sex					
Male	264	48,887	35.7	30.85	40.83
Female	337	47,180	34.0	30.07	38.27
Form					
Form 1	133	22,665	39.1	34.49	43.93
Form 2	135	18,024	33.9	28.01	40.38
Form 3	121	19,320	35.6	28.79	43.12
Form 4	118	17,761	32.9	28.54	37.51
Form 5	94	18,297	32.6	24.00	42.46

Table 3.9.5 : Prevalence of parental or guardian bonding in the past 30 days among Form 1 to Form 5 students in Johor, 2017

Socio-demographic Characteristics	Most of the time or always				
	Unweighted Count	Estimated Population	Prevalence (%)	95% CI	
				Lower	Upper
JOHOR	889	140,444	51.0	46.88	55.13
Locality of school					
Urban	640	100,004	51.1	45.32	56.78
Rural	249	40,440	50.9	48.88	52.89
Sex					
Male	368	67,300	49.1	43.73	54.44
Female	521	73,144	52.9	48.36	57.47
Form					
Form 1	177	29,922	51.6	44.43	58.75
Form 2	184	24,438	46.0	38.58	53.53
Form 3	192	30,460	56.3	51.13	61.41
Form 4	179	25,854	48.0	41.37	54.63
Form 5	157	29,771	53.0	45.66	60.15

Table 3.9.6 : Prevalence of parental or guardian respect for privacy in the past 30 days among Form 1 to Form 5 students in Johor, 2017

Socio-demographic Characteristics	Never or rarely				
	Unweighted Count	Estimated Population	Prevalence (%)	95% CI	
				Lower	Upper
JOHOR	1,353	213,949	77.6	74.58	80.42
Locality of the school					
Urban	986	154,494	78.8	75.30	81.90
Rural	367	59,455	74.8	69.95	79.12
Sex					
Male	538	99,033	72.2	69.94	74.37
Female	815	114,917	83.0	78.35	86.84
Form					
Form 1	259	43,670	75.3	69.23	80.58
Form 2	294	39,041	73.2	70.29	76.02
Form 3	273	43,650	80.7	76.54	84.30
Form 4	296	43,133	79.8	74.51	84.27
Form 5	231	44,456	79.1	73.35	83.88

3.10 Sexual Behaviours that contribute to HIV Infection, Other STIs and Unintended Pregnancy

Contributors: Maisarah Omar, Noor Aliza Lodz, Noraida Mohamad Kasim, S Maria Awaluddin, Noor Ani Ahmad, Nik Rubiah Nik Abdul Rashid

3.10.1 Introduction

The WHO reported that 10% to 40% of young unmarried girls aged 13 to 19 years to have had an unintended pregnancy.¹ It was also reported that the highest rate of Sexual Transmitted Illness (STI) worldwide are among young people aged 15 to 24 years.² In Malaysia, Ministry of Health revealed that 12,492 teenage pregnancies were recorded in 2016. Therefore, sexual intercourse among school aged students is a public health concern in Malaysia due to the alarming numbers of the teenage unintended pregnancies and Sexual Transmitted Illness (STI) from this age group.^{2,3}

3.10.2 Objectives

To identify prevalence of:

- i. Current sexual intercourse in the past 30 days among Form 1 to Form 5 students in Malaysia.
- ii. Ever having sexual intercourse among Form 1 to Form 5 students in Malaysia.
- iii. First sexual experience before the age 14 years among those who ever had sex.
- iv. Having at least two sexual partners among those who ever had sex.
- v. Condom use during the last sexual intercourse among those who ever had sex.
- vi. Use of "other birth control methods" during the last sexual intercourse among those who ever had sex.

3.10.3 Variable Definitions

- **Sexual intercourse:** sexual acts of penetration of penis into vagina or anus.
- **Safe sexual practice:** sexual contact that doesn't involve the exchange of semen, vaginal fluids or blood between partners.

3.10.4 Findings

Prevalence of ever-had sex among Form 1 to Form 5 students in Johor was 5.7% (95% CI: 4.66, 7.07). Prevalence of ever-had sex among male students were 6.4% (95% CI: 4.71, 8.55) while female students were at 5.1% (95% CI: 3.83, 6.86). Form 2 students showed the highest prevalence of 7.5% (95% CI:4.43, 12.31) (**Table 3.10.1**). The prevalence of Form 1 to Form 5 students that were currently having sexual intercourse in Johor was 4.2% (95% CI: 3.18, 5.63). Form 2 students showed the highest prevalence of 5.0% (95% CI:3.19, 7.64) compared to other forms (**Table 3.10.2**).

Of those who ever-had sex, 27.9% (95% CI: 21.14, 35.82) of them had sex before the age of 14 years. Form 2 students showed the highest percentage at 37.3% (95% CI:22.67, 54.74) (**Table 3.10.3**). It was noted that 13.9% (95% CI: 7.69, 23.72) of those who were ever-had sex, had at least two sexual partners in which, male students were 15.9% (95% CI: 6.84, 32.92) and females were 11.3% (95% CI: 4.26, 26.82) (**Table 3.10.4**). Only 5.4% (95% CI: 2.80, 10.10) of those who ever-had sex used a condom during their last sexual intercourse while 8.1% (95% CI: 4.39, 14.61) used other birth control methods (**Table 3.10.5 &Table 3.10.6**).

3.10.5 Discussion/ Conclusion

The prevalence of ever had sexual intercourse in Johor AHS 2017 (5.7%) was lower as compared to Malaysia GSHS in 2012 (8.3%)⁴ and national prevalence which was 7.3%. The previous Johor GSHS 2012 observed the prevalence of ever had sexual intercourse was 10.6%.⁴

3.10.6 Recommendations

1. To strengthen sexual health education especially among students and to educate them regarding the complications of unsafe sex.
2. To conduct more studies especially qualitative studies in exploring the sexual orientation of the students, abortion and unintended pregnancy.
3. To strengthen and actively provide services of our adolescent health screening (BSSK) in schools.

3.10.7 References

1. The Lancet's Maternal Survival and Women Deliver Series 2006/2007: 2005 World Health Report
2. Satterwhite CL, Torrone E, Meites E, Dunne EF, Mahajan R, Ocfemia MC, et al. Sexually transmitted infections among US women and men: prevalence and incidence estimates, 2008. *Sex Transm Dis.* 2013;40(3):187-93.
3. Family Health Development Division, Ministry of Health Malaysia, 2016.
4. Institute for Public Health(IPH) 2012.The National Health and Morbidity Survey: Malaysia Global School Based Student Health Survey 2012.Kuala Lumpur: Ministry of Health Malaysia

Table 3.10.1 : Prevalence of ever had sexual intercourse among Form 1 to Form 5 students in Johor, 2017

Socio-demographic characteristics	ever had sexual intercourse				
	Unweighted Count	Estimated Population	Prevalence (%)	95% CI	
				Lower	Upper
JOHOR	97	15,849	5.7	4.66	7.07
Locality of school					
Urban	71	11,361	5.8	4.29	7.77
Rural	26	4,488	5.6	4.28	7.41
Sex					
Male	47	8,728	6.4	4.71	8.55
Female	50	7,121	5.1	3.83	6.86
Form					
Form 1	15	2,851	4.9	2.63	9.02
Form 2	29	3,979	7.5	4.43	12.31
Form 3	19	3,012	5.6	3.80	8.06
Form 4	18	2,817	5.2	3.19	8.42
Form 5	16	3,190	5.7	3.81	8.39

Table 3.10.2 Prevalence of current sexual intercourse in the past 30 days among Form 1 to Form 5 students in Johor, 2017

Socio-demographic characteristics	current sexual intercourse				
	Unweighted Count	Estimated Population	Prevalence (%)	95% CI	
				Lower	Upper
JOHOR	70	11,678	4.2	3.18	5.63
Locality of school					
Urban	54	8,843	4.5	3.39	5.97
Rural	16	2,835	3.6	1.45	8.49
Sex					
Male	37	7,006	5.1	3.37	7.67
Female	33	4,672	3.4	2.35	4.82
Form					
Form 1	10	1,947	3.4	1.54	7.15
Form 2	19	2,645	5.0	3.19	7.64
Form 3	13	2,049	3.8	2.15	6.55
Form 4	16	2,529	4.7	2.87	7.56
Form 5	12	2,508	4.5	2.62	7.49

Table 3.10.3 : Percentage of first sex before the age of 14 years among those who ever had sex among Form 1 to Form 5 students in Johor, 2017

Socio-demographic characteristics	Percentage of first sex before the age 14 years among those who ever had sex				
	Unweighted Count	Estimated Population	Prevalence (%)	95% CI	
				Lower	Upper
JOHOR	28	4,421	27.9	21.14	35.82
Locality of school					
Urban	19	2,927	25.8	21.25	30.86
Rural	9	1,493	33.3	13.52	61.40
Sex					
Male	14	2,446	28.0	16.23	43.90
Female	14	1,975	27.7	17.67	40.69
Form					
Form 1	5	904	31.7	15.42	54.15
Form 2	11	1,485	37.3	22.67	54.74
Form 3	6	964	32.0	8.61	70.12
Form 4	3	449	16.0	4.69	42.26
Form 5	3	619	19.4	6.90	43.83

Table 3.10.4 : Percentage of having at least two sexual partners among students who ever had sex Form 1 to Form 5 students in Johor, 2017

Socio-demographic characteristics	Percentage of having at least two sexual partners among those who ever had sex				
	Unweighted Count	Estimated Population	Prevalence (%)	95% CI	
				Lower	Upper
JOHOR	13	2,198	13.9	7.69	23.72
Locality of school					
Urban	10	1,684	14.8	6.98	28.75
Rural	3	514	11.5	3.40	32.24
Sex					
Male	8	1,392	15.9	6.84	32.92
Female	5	806	11.3	4.26	26.82
Form					
Form 1	1	193	6.8	0.74	41.50
Form 2	6	946	23.8	6.41	58.69
Form 3					
Form 4	2	293	10.4	2.10	38.66
Form 5	4	766	24.0	7.21	56.20

Table 3.10.5 : Percentage of reported condom use during the last sexual intercourse among students who ever had sex Form 1 to Form 5 students in Johor, 2017

Socio-demographic characteristics	condom use				
	Unweighted Count	Estimated Population	Prevalence (%)	95% CI	
				Lower	Upper
JOHOR	5	853	5.4	2.80	10.10
Locality of school					
Urban	4	676	6.0	2.78	12.31
Rural	1	177	3.9	0.64	20.80
Sex					
Male	2	366	4.2	1.12	14.51
Female	3	487	6.8	2.15	19.74
Form					
Form 1	2	349	12.2	2.34	44.82
Form 2	1	173	4.3	0.71	22.30
Form 3					
Form 4					
Form 5	2	332	10.4	2.22	37.21

Table 3.10.6 : Percentage of reported using "other birth control methods" during the last sexual intercourse among students who ever had sex Form 1 to Form 5 students in Johor, 2017

Socio-demographic characteristics	other birth control methods				
	Unweighted Count	Estimated Population	Prevalence (%)	95% CI	
				Lower	Upper
JOHOR	8	1290	8.1	4.39	14.61
Locality of school					
Urban	5	794	7.0	3.29	14.26
Rural	3	496	11.0	2.91	33.92
Sex					
Male	4	642	7.4	2.56	19.38
Female	4	647	9.1	2.61	27.18
Form					
Form 1	1	156	5.5	0.50	39.91
Form 2	4	642	16.1	5.54	38.71
Form 3					
Form 4					
Form 5	3	492	15.4	2.68	54.64

3.11 Tobacco Use

Contributors: Muhammad Fadhli Bin Mohd Yusoff, Jane Ling Miaw Yn, Tee Guat Hiong, Noraryana Binti Hassan, Nizam Bin Baharom, Wan Shakira Rodzlan Hasani, Tania Gayle Robert, Halizah Mat Rifin, Hamizatul Akmal Abd Hamid, Abdul Aiman Abd Ghani, Hasimah Ismail, Thamil Arasu Saminathan, Nur Liana Ab. Majid.

3.11.1 Introduction

Most smokers experimented and started smoking during their school aged days. For example, in Malaysian Global Adults Tobacco Survey (GATS) 2011, 51.8% of the smokers started smoking daily before the age of 18.¹ Therefore, monitoring tobacco use among Malaysian adolescents is an important process in tobacco control. Malaysia is committed with the implementation of the WHO Framework Convention on Tobacco Control to achieve targeted lower smoking prevalence and creating smokefree Malaysian generations.

3.11.2 Objectives

To identify the prevalence of:

- i. Current smokers
- ii. Current cigarette smokers
- iii. Current E-cigarette/vape users
- iv. Current any tobacco product users
- v. Current smoke and smokeless tobacco product use by types
- vi. Having tried a cigarette before the age of 14 years, among ever cigarette smokers
- vii. Having tried E-cigarettes or vape before the age of 14 years, among ever E-cigarette or vape users
- viii. Quit smoking attempt in the past 12 months among those who ever smoked cigarettes
- ix. Quit E-cigarettes or vape attempt in the past 12 months among those who ever used E-cigarettes or vape
- x. Having been exposed to people smoking in their presence for at least one day in the past seven days

- xi. Having parents or guardians who used any form of tobacco products
- xii. Having parents or guardians who used E-cigarettes or vape

3.11.3 Variable Definitions

- **Current smoker:** Used any smoked tobacco products in the past 30 days (Manufactured cigarette, Roll-your-own cigarette, Traditional hand-rolled cigarette, Shisha, Cigar, or Pipe).
- **Current cigarette smoker:** smoked manufactured cigarette, rolled-your-own, or traditional hand roll cigarette in the past 30 days.
- **Current E-cigarette/Vape user:** Used E-cigarette/Vape in the past 30 days.
- **Current any tobacco product user:** Used any tobacco products in the past 30 days (Manufactured cigarette, Roll-your-own cigarette, Traditional hand-rolled cigarette, Shisha, Cigar, and Pipe, E-cigarette/Vape, Chewing tobacco or Snuff)

3.11.4 Findings

The prevalence of current smokers was 10.6% (95% CI: 7.33, 15.20) (**Table 3.11.1**). The prevalence was significantly higher among males (18.4%; 95% CI: 12.17, 26.77) as compared to females (3.0%; 95% CI: 2.09, 4.24). Form 4 students had the highest prevalence of 13.5% (95% CI: 6.68, 25.26) with no significant difference across all forms.

The prevalence of current cigarettes smokers was 9.2% (95% CI: 6.27, 13.32) (**Table 3.11.2**). It was significantly higher in males (16.2%; 95% CI: 10.76, 23.58) than females (2.3%; 95% CI: 1.48, 3.60). Form 4 students reported highest prevalence (11.1%; 95% CI: 5.10, 22.35) compared to the other forms. Among those who ever smoked cigarettes, 73.6% (95%CI: 58.45, 84.71) had first tried a cigarette before the age of 14 years, with no significant difference by gender (**Table 3.11.6**).

Overall, 14.9% (95% CI: 10.68, 20.38) currently use any tobacco products (**Table 3.11.3**). Males (24.1%; 95% CI: 17.22, 32.58) had significantly higher prevalence compared to females (5.8%; 95% CI: 4.10, 8.16). Form 4 students reported highest prevalence (17.9%; 95% CI: 10.22, 29.41) compared to the other forms.

The prevalence of using shisha/hookah in the past 30 days was 5.7% (95% CI: 3.47, 9.13) with significantly more males [9.7% (95% CI: 5.46,16.64)] than females [1.7% (95% CI: 0.95, 2.95)] (**Table 3.11.5**). The prevalence of traditional rolled cigarette use was 2.2% (95% CI: 1.53, 3.17), with no significant difference between gender (**Table 3.11.5**). The prevalence of rolled your own was 2.0% (95% CI: 1.34,2.88) and with no significant difference between gender (**Table 3.11.5**). About 1.8% (95% CI: 1.15, 2.78) of the students used cigar (**Table 3.11.5**). The prevalence of using pipe was 1.6% (95% CI: 0.96, 2.79) (**Table 3.11.5**).

Among those who ever smoked cigarettes, 74.6% (95% CI: 59.31, 85.54) had tried to stop smoking in the past 12 months (**Table 3.11.8**). Males reported significantly higher prevalence (77.5%; 95% CI: 62.54, 87.66) than females (27.6%; 95% CI: 9.46, 58.08). Form 4 students had the highest prevalence (90.7%; 95% CI: 76.00, 96.76) compared to the other forms.

A total of 46.5% (95% CI: 40.79, 52.36) reported having been exposed to people who smoked in their presence in the past seven days (**Table 3.11.10**). Males reported significantly higher prevalence (53.6%; 95% CI: 45.78, 61.17) than females (39.6%; 95% CI: 34.01, 45.39). Form 4 students reported highest prevalence (57.6%; 95% CI: 46.86, 67.60) compared to other forms.

Overall, 42.8% (95% CI: 38.30, 47.32) of students reported having parents or guardian who used any form of tobacco products with no significant difference between student's gender (**Table 3.11.11**). Form 3 students had the highest prevalence (44.9%; 95% CI: 38.96, 50.94) compared to other forms.

The prevalence of current E-cigarette use was 6.6% (95% CI: 4.70, 9.21) (**Table 3.11.4**). Males reported significantly higher prevalence 12.2% (95% CI: 8.99, 16.39) compared to females at 1.1% (95% CI: 0.53, 2.12). Form 5 students had the highest prevalence (8.5%; 95% CI: 4.42, 15.74) compared to the other forms. Among those who ever smoked e-cigarette/vape, 46.7% (95%CI: 32.87, 61.13) had first tried e-cigarette/vape before the age of 14 years with no significant difference by gender (**Table 3.11.7**). Among those who ever used e-cigarette/vape, 68.7% (95% CI: 55.92, 79.16) had tried to stop using e-cigarette/ vape in the past 12 months (**Table 3.11.9**). The prevalence was significantly higher in males (74.0%; 95% CI: 60.78, 83.95) than in females (31.8%; 95% CI: 13.58, 58.12). Form 3 students had the highest prevalence

(80.4%; 95% CI: 52.53, 93.80) compared to other forms. Overall, 10.5% (95% CI: 8.58, 12.77) of students reported having parents or guardians who used e-cigarette/vape (**Table 3.11.12**). No significant difference was reported between the genders. Form 2 students had the highest prevalence (13.0%; 95% CI: 8.95, 18.56) compared to other forms.

The prevalence of **chewing tobacco** was 1.0% (95% CI: 0.58, 1.66), with no significant difference between gender (**Table 3.11.5**). The prevalence of **snuff use** was 4.5% (95% CI: 3.13, 6.30) with no significant difference between gender (**Table 3.11.5**).

3.11.5 Discussion/Conclusion

The prevalence of cigarette smoking in Johor was not significantly different compared to the national prevalence. Similarly, the prevalence of having tried a cigarette before the age of 14 among ever cigarette smokers and the prevalence of having been exposed to people who smoke in their presence in the past seven days showed no significant difference compared to the national prevalence.

3.11.6 Recommendations

All screening, prevention and intervention programmes among adolescents must be strengthened and delivered in synergy by all governmental and non-governmental agencies. Interventions should also be targeted to higher risk groups such as states with high prevalence, males and Form 1 students. The rise of prevalence of smoking among females should also be a concern for additional interventions to halt this increment. Overall national tobacco control programmes have to be amplified to achieve denormalization of public smoking, and to help reduce exposure to cigarette smoke among our youth. These would help to achieve the medium and long-term targets as stipulated in the National Strategic Plan for Tobacco Control.

3.11.7 References

1. Institute for Public Health (IPH). Report of the Global Adult Tobacco Survey (GATS) Malaysia, 2011, Ministry of Health Malaysia, 2012.

Table 3.11.1: Prevalence of current smokers among Form 1 to Form 5 students in Johor, 2017

Socio-Demographic Characteristics	Current Smokers				
	Unweighted Count	Estimated Population	Prevalence (%)	95% CI	
				Lower	Upper
JOHOR	163	29,327	10.6	7.33	15.20
Locality of school					
Urban	95	16,749	8.5	5.41	13.21
Rural	68	12,578	15.8	9.83	24.49
Sex					
Male	134	25,193	18.4	12.17	26.77
Female	29	4,134	3.0	2.09	4.24
Form					
Form 1	16	3,001	5.2	2.36	10.99
Form 2	43	6,188	11.6	6.40	20.14
Form 3	35	6,396	11.8	7.22	18.69
Form 4	42	7,273	13.5	6.68	25.26
Form 5	27	6,469	11.5	6.23	20.30

Table 3.11.2: Prevalence of current cigarette smokers among Form 1 to Form 5 students in Johor, 2017

Socio-Demographic Characteristics	Current Cigarette Smokers				
	Unweighted Count	Estimated Population	Prevalence (%)	95% CI	
				Lower	Upper
JOHOR	141	25,386	9.2	6.27	13.32
Locality of school					
Urban	79	13,951	7.1	4.53	10.99
Rural	62	11,435	14.4	8.97	22.28
Sex					
Male	119	22,177	16.2	10.76	23.58
Female	22	3,208	2.3	1.48	3.60
Form					
Form 1	14	2,563	4.4	1.85	10.19
Form 2	39	5,616	10.5	5.60	18.96
Form 3	30	5,471	10.1	5.90	16.72
Form 4	34	5,978	11.1	5.10	22.35
Form 5	24	5,757	10.2	5.44	18.46

Table 3.11.3: Prevalence of current any tobacco products use among Form 1 to Form 5 students in Johor, 2017

Socio-Demographic	Unweighted Count	Estimated Population	Prevalence (%)	95% CI	
				Lower	Upper
JOHOR	231	41,053	14.9	10.68	20.38
Locality of school					
Urban	142	25,046	12.8	8.34	19.04
Rural	89	16,007	20.1	13.02	29.82
Sex					
Male	175	33,011	24.1	17.22	32.58
Female	56	8,042	5.8	4.10	8.16
Form					
Form 1	31	5,805	10.0	5.73	16.92
Form 2	59	8,328	15.6	9.32	25.01
Form 3	48	8,628	15.9	9.62	25.19
Form 4	57	9,662	17.9	10.22	29.41
Form 5	36	8,629	15.4	8.52	26.09

Table 3.11.4: Prevalence of current E-cigarettes/Vape use among Form 1 to Form 5 students in Johor, 2017

Socio-Demographic Characteristics	Current e-Cigarette Users				
	Unweighted Count	Estimated Population	Prevalence (%)	95% CI	
				Lower	Upper
JOHOR	99	18,217	6.6	4.70	9.21
Locality of school					
Urban	63	11,436	5.8	3.62	9.25
Rural	36	6,781	8.5	6.05	11.91
Sex					
Male	88	16,752	12.2	8.99	16.39
Female	11	1,465	1.1	0.53	2.12
Form					
Form 1	8	1,499	2.6	0.97	6.72
Form 2	25	3,588	6.7	3.11	13.96
Form 3	23	4,201	7.7	4.18	13.93
Form 4	24	4,148	7.7	3.96	14.36
Form 5	19	4,781	8.5	4.42	15.74

Table 3.11.5: Prevalence of current smoke and smokeless tobacco product use by types (except manufactured cigarette and E-cigarette/Vape) among Form 1 to Form 5 students in Johor, 2017

	Total					Male				
	Unweighted Count	Estimated Population	Prevalence (%)	95% CI		Unweighted Count	Estimated Population	Prevalence (%)	95% CI	
				Lower	Upper				Lower	Upper
Shisha Use	84	15,586	5.7	3.47	9.13	67	13,265	9.7	5.46	16.64
Traditional Rolled Cigarette Use	38	6,064	2.2	1.53	3.17	23	3,904	2.9	1.66	4.84
Roll-Your-Own" With Cigarette Paper Use"	33	5,418	2.0	1.34	2.88	20	3,622	2.6	1.57	4.43
Cigar Use	28	4,932	1.8	1.15	2.78	16	3,210	2.3	1.31	4.18
Pipe Smoking Use	27	4,524	1.6	0.96	2.79	14	2,700	2.0	0.95	4.07
Chewing Tobacco Use	17	2,712	1.0	0.58	1.66	6	1,149	0.8	0.38	1.83
Snuff Use	73	12,239	4.5	3.13	6.30	41	7,585	5.6	3.90	7.84

Table 3.11.5: Prevalence of current smoke and smokeless tobacco product use by types (except manufactured cigarette and E-cigarette/Vape) among Form 1 to Form 5 students in Johor, 2017 (Cont.)

	Female				
	Unweighted Count	Estimated Population	Prevalence (%)	95% CI	
				Lower	Upper
Shisha Use	17	2,321	1.7	0.95	2.95
Traditional Rolled Cigarette Use	15	2,160	1.6	0.94	2.59
Roll-Your-Own" With Cigarette Paper Use"	13	1,796	1.3	0.63	2.65
Cigar Use	12	1,723	1.2	0.59	2.60
Pipe Smoking Use	13	1,824	1.3	0.67	2.60
Chewing Tobacco Use	11	1,563	1.1	0.53	2.40
Snuff Use	32	4,654	3.4	1.82	6.15

Table 3.11.6: Prevalence of having tried a cigarette before the age of 14 years among ever smokers among Form 1 to Form 5 students in Johor, 2017

	Total					Male				
	Unweighted Count	Estimated Population	Prevalence (%)	95% CI		Unweighted Count	Estimated Population	Prevalence (%)	95% CI	
				Lower	Upper				Lower	Upper
Yes	140	25,745	73.6	58.45	84.71	134	24,894	74.0	58.15	85.37
No	45	9,220	26.4	15.29	41.55	42	8,741	26.0	14.63	41.85

Table 3.11.6: Prevalence of having tried a cigarette before the age of 14 years among ever smokers among Form 1 to Form 5 students in Johor, 2017 (Cont.)

	Female				
	Unweighted Count	Estimated Population	Prevalence (%)	95% CI	
				Lower	Upper
Yes	6	852	64.0	18.84	93.16
No	3	479	36.0	6.84	81.16

Table 3.11.7: Prevalence of having tried a E-cigarette/Vape before the age of 14 years among ever E-cigarette/Vape users among Form 1 to Form 5 students in Johor, 2017

	Total					Male				
	Unweighted Count	Estimated Population	Prevalence (%)	95% CI		Unweighted Count	Estimated Population	Prevalence (%)	95% CI	
				Lower	Upper				Lower	Upper
Yes	85	14,917	46.7	32.87	61.13	81	14,401	47.5	33.34	62.05
No	86	16,997	53.3	38.87	67.13	78	15,924	52.5	37.95	66.66

Table 3.11.7: Prevalence of having tried a E-cigarette/Vape before the age of 14 years among ever E-cigarette/Vape users among Form 1 to Form 5 students in Johor, 2017 (Cont.)

	Female				
	Unweighted Count	Estimated Population	Prevalence (%)	95% CI	
				Lower	Upper
Yes	4	516	32.5	15.42	55.90
No	8	1,073	67.5	44.10	84.58

Table 3.11.8: Prevalence of quit smoking attempt in the past 12 month among those who ever smoked among Form 1 to Form 5 students in Johor, 2017

Socio-Demographic Characteristics	Quit Smoking Attempts In The Past 12 Months Among Those Who Ever Smoked Cigarettes				
	Unweighted Count	Estimated Population	Prevalence (%)	95% CI	
				Lower	Upper
JOHOR	116	21,600	74.6	59.31	85.54
Locality of school					
Urban	67	12,436	68.9	50.08	83.08
Rural	49	9,163	83.9	73.47	90.80
Sex					
Male	113	21,136	77.5	62.54	87.66
Female	3	464	27.6	9.46	58.08
Form					
Form 1	14	2,671	89.6	68.62	97.15
Form 2	27	3,901	63.7	31.07	87.27
Form 3	22	4,067	72.5	58.95	82.88
Form 4	34	6,120	90.7	76.00	96.76
Form 5	19	4,840	64.6	35.86	85.59

Table 3.11.9: Prevalence of quit E-cigarette/ Vape attempt in the past 12 month among those who ever used E-Cigarette/Vape among Form 1 to Form 5 students in Johor, 2017

Socio-Demographic Characteristics	Stop Using e-Cigarettes In The Past 12 Months Among Those Who Ever Smoked e-Cigarettes				
	Unweighted Count	Estimated Population	Prevalence (%)	95% CI	
				Lower	Upper
JOHOR	97	17,864	68.7	55.92	79.16
Locality of school					
Urban	54	9,815	61.4	47.77	73.38
Rural	43	8,050	80.4	72.46	86.54
Sex					
Male	89	16,823	74.0	60.78	83.95
Female	8	1,041	31.8	13.58	58.12
Form					
Form 1	13	2,505	69.6	39.76	88.80
Form 2	22	3,086	54.4	33.90	73.57
Form 3	19	3,598	80.4	52.53	93.80
Form 4	30	5,296	79.0	60.62	90.21
Form 5	13	3,380	60.8	43.01	76.19

Table 3.11.10: Prevalence of being expose to people smoking in their presence for at least one day in the past 7 days among Form 1 to Form 5 students in Johor, 2017

Socio-Demographic Characteristics	Having Been Exposed To People Smoking In Their Presence For At Least One Days In The Past 7 Days				
	Unweighted Count	Estimated Population	Prevalence (%)	95% CI	
				Lower	Upper
JOHOR	780	128,215	46.5	40.79	52.36
Locality of school					
Urban	521	84,833	43.3	37.47	49.25
Rural	259	43,382	54.6	45.38	63.49
Sex					
Male	394	73,456	53.6	45.78	61.17
Female	386	54,759	39.6	34.01	45.39
Form					
Form 1	113	19,550	33.7	27.86	40.15
Form 2	157	21,366	40.1	29.85	51.27
Form 3	166	27,150	50.2	44.25	56.17
Form 4	201	31,104	57.6	46.86	67.60
Form 5	143	29,044	51.7	44.62	58.67

Table 3.11.11: Prevalence of having father or mother or guardian who used any form of tobacco product among Form 1 to Form 5 students in Johor, 2017

Socio-Demographic Characteristics	Either Father Or Mother Or Guardian				
	Unweighted Count	Estimated Population	Prevalence (%)	95% CI	
				Lower	Upper
JOHOR	704	112,294	42.8	38.30	47.32
Locality of school					
Urban	479	75,500	40.4	35.58	45.46
Rural	225	36,793	48.5	42.63	54.39
Sex					
Male	284	52,272	40.4	35.83	45.08
Female	420	60,021	45.1	38.86	51.43
Form					
Form 1	142	23,955	42.6	38.50	46.89
Form 2	148	19,767	39.8	32.27	47.86
Form 3	149	23,466	44.9	38.96	50.94
Form 4	152	22,589	43.9	36.18	52.00
Form 5	113	22,516	42.4	33.77	51.47

Table 3.11.12: Prevalence of having father or mother or guardian who used E-cigarette/ Vape among Form 1 to Form 5 students in Johor, 2017

Socio-Demographic Characteristics	Either Father Or Mother Or Guardian				
	Unweighted Count	Estimated Population	Prevalence (%)	95% CI	
				Lower	Upper
JOHOR	172	26,613	10.5	8.58	12.77
Locality of school					
Urban	114	17,329	9.5	7.37	12.25
Rural	58	9,284	12.9	10.72	15.47
Sex					
Male	59	10,326	8.2	5.90	11.24
Female	113	16,287	12.8	9.40	17.15
Form					
Form 1	39	6,616	12.3	9.23	16.24
Form 2	46	6,118	13.0	8.95	18.56
Form 3	37	5,732	11.2	9.13	13.59
Form 4	34	5,169	10.6	8.01	13.78
Form 5	16	2,977	5.7	3.03	10.33

3.12 Violence and Unintentional Injury

Contributors: Rajini Sooryanarayana, Shubash Shander Ganapathy, Azriman Rosman, Mohd Hazrin Bin Hasim@Hashim, Fazila Haryati Ahmad, Mohamad Fuad Mohamad Anuar, Tan Lee Ann, Thamil Arasu a/l Saminathan, Muhammad Fadhli Mohd Yusoff, S Maria Awaluddin

3.12.1 Introduction

The World Health Assembly through WHA 49.25 had declared violence to be a public health problem globally. Malaysia is committed towards resolving this issue, covering various aspects of interpersonal violence.¹ The prevalence of bullying varies between 9% to 25% in school-going adolescents in Western countries², and 10.8% to 17.6% locally³, differences largely explained by linguistic and cultural factors.

Injuries are inevitable among anyone, including adolescents. Injuries however, may be due to intentional or unintentional causes. Unintentional injuries are caused by various factors and this leads to health problems and disability in an otherwise healthy population. Conflicts happen due to a wide spectrum ranging from physical fights to bullying. These may affect the mental health, well-being and healthy development of the adolescent.

This survey aims to examine the prevalence and distribution of unintentional injuries, physical attacks, physical fights, bullying, as well as physical and verbal abuse experienced by adolescents in Form 1 to Form 5 in Malaysia.

3.12.2 Objectives

To describe the prevalence of:

- i. Having been physically attacked at least once in the past 12 months
- ii. Involvement in a physical fight at least once in the past 12 months
- iii. Having been bullied at least once in the past 30 days
- iv. Physical abuse at home at least once in the past 30 days
- v. Verbal abuse at home at least once in the past 30 days
- vi. Having had a serious injury at least once in the past 12 months

3.12.3 Variable Definitions

- **Physical attack:** when one or more persons hurt another person with or without a weapon such as sticks or knives in the past 12 months. It is NOT a physical attack when two individuals or students of about the same strength or power choose to fight each other.
- **Physical fight:** when two individuals or students of about the same strength or power choose to fight each other in the past 12 months.
- **Bullying:** when a student or group of students say or do bad and unpleasant things to another student, such as teasing a lot in an unpleasant way or leaving out of things on purpose in the past 30 days. It is NOT bullying when two students of about the same strength or power argue or fight or when teasing is done in a friendly and fun way.
- **Physical abuse at home:** when someone is hit so hard that it left a mark OR caused an injury in the past 30 days
- **Verbal abuse at home:** when someone has had hurtful or insulting things said to them in the past 30 days
- **Unintentional injury:** a serious injury which makes the student miss at least one full day of usual activity (such as school, sports or a job) OR requires treatment by a doctor or a medical personnel in the past 12 months

3.12.4 Findings

Among the adolescents in this survey, 26.1% (95% CI: 22.15, 30.47) of them had been physically attacked, significantly higher in males [33.8% (95% CI: 28.44, 39.60)] compared to females [18.5% (95% CI: 14.66, 23.03)]. There was no significant difference found between urban [24.7% (95% CI: 19.78, 30.29)] and rural [29.7% (95% CI: 25.33, 34.38)] areas. Adolescents from Form 1 had the highest prevalence of being physically attacked; 34.7% (95% CI: 26.34, 44.09) while adolescents from Form 5 had the lowest prevalence at 18.8% (95% CI: 13.86, 25.03) (**Table 3.12.1**).

Similarly, 21.3% (95% CI: 17.62, 25.62) of adolescents claimed to have been involved in physical fight(s), with significantly more males [27.8% (95% CI: 23.34, 32.80)] than females [14.9% (95% CI: 11.62, 18.96)] involved. There was no significant difference observed between urban and rural areas, with a reported prevalence of 20.8% (95% CI: 16.22, 26.27) and 22.7% (95% CI: 17.18, 29.37) respectively. Adolescents from Form 1 had the highest prevalence at 31.7% (95% CI: 24.86, 39.47) (**Table 3.12.1**).

With regards to bullying, 13.9% (95% CI: 10.89, 17.54) of adolescents reported to have been bullied, and this was higher among males [17.0% (95% CI: 13.62, 21.08)] compared to females [10.8% (95% CI: 7.81, 14.72)]. There was no significant difference found between urban [12.8% (95% CI: 9.10, 17.59)] and rural [16.7% (95% CI: 13.35, 20.65)] areas. Adolescents from Form 2 had the highest prevalence of have been bullied; 21.5% (95% CI: 13.14, 33.18) (**Table 3.12.2**). The two most common forms of bullying were 'made fun of with sexual jokes, comments or gestures' [17.6% (95% CI: 10.81, 27.33)] and 'being made fun of because of how body or face looks' [16.2% (95% CI: 11.99, 21.64)]. Bullying in the form of 'being made fun of because of religion' was the least common at 2.5% (95% CI: 1.01, 6.15) (**Table 3.12.3**).

Physical abuse at home was reported by 8.6% (95% CI: 5.59, 13.02) of the adolescents. Urban areas reported a prevalence of 8.7% (95% CI: 5.11, 14.32) with rural at 8.5% (95% CI: 4.02, 16.92). There was no significant difference between males and females with a prevalence of 7.7% (95% CI: 4.77, 12.04) and 9.6% (95% CI: 5.79, 15.38) respectively. Form 5 adolescents reported the lowest prevalence at 4.5% (95% CI: 1.84, 10.65) (**Table 3.12.4**).

It was reported that 45.2% (95% CI: 38.78, 51.71) of adolescents were abused verbally at home. The prevalence of verbal abuse was higher among females [51.4% (95% CI: 41.98, 60.79)] compared to males [38.8% (95% CI: 32.79, 45.27)]. There was no significant difference found between urban [42.1% (95% CI: 34.49, 50.02)] and rural [52.8% (95% CI: 45.89, 59.64)] areas. Adolescents from Form 4 had the highest prevalence of verbal abuse at home with a reported prevalence of 48.5% (95% CI: 38.97, 58.16) (**Table 3.12.4**).

It was found that 26.2% of adolescents in Johor (95% CI: 22.20, 30.70) had been seriously injured in the past 12 months. There was no significant difference found between urban [25.8% (95% CI: 20.59, 31.86)] and rural [27.2% (95% CI: 22.70, 32.24)] localities. Prevalence of serious injury was higher among males [30.6% (95% CI: 25.20, 36.53)] as compared to females [21.9% (95% CI: 18.03, 26.38)]. Adolescents from Form 2 had the highest prevalence of serious injury at 30.6% (95% CI: 23.29, 38.95) while adolescents from Form 5 had the lowest prevalence at 21.1% (95% CI: 15.62, 28.08) (**Table 3.12.5**). Among those who had been seriously injured, the two most common injuries were 'a broken bone/dislocated joint' [20.4% (95% CI: 15.45, 26.54)] and 'a cut or stab wound' [19.9% (95% CI: 15.89, 24.57)] (**Table 3.12.6**). The two most common causes of serious injury were falls [37.9% (95% CI: 31.84, 44.31)] and motor vehicle accidents [15.2% (95% CI: 11.07, 20.60)] (**Table 3.12.7**).

3.12.5 Discussion / Conclusion

The prevalence of being physically attacked among adolescents in Johor was higher than the national prevalence (26.1% vs 25.3%) but lower than the 29.1% reported in Johor GSHS 2012⁴. Those involved in physical fights were reported to be lower than the national prevalence (21.3% vs 24.9%) as well as lower than the 28.1% reported in Johor GSHS 2012⁴.

The prevalence of bullying among school going adolescents in Johor was lower than the reported national prevalence (13.9% vs 16.2%) and lower than the 18.5% reported in Johor GSHS 2012⁴. The prevalence of physical abuse at home among adolescents was lower in Johor compared to national prevalence (8.6% vs 11.8%), as well as lower than the 13.3% reported in Johor GSHS 2012⁴. However, the prevalence of verbal abuse was found to be higher than the national prevalence (45.2% vs 43.2%), as well as higher than the 41.2% reported in Johor

in 2012⁴. The prevalence of serious injury among school going adolescents in Johor was lower than the national prevalence found in this study (26.2% vs. 29.9%) as well as the 35.5% reported in Johor GSHS 2012⁴.

Studies have shown that violence related behaviour is associated with other negative factors such as smoking, drug abuse, truancy and physical and mental health problems.⁵ Thus, despite the successes achieved by our current programs, the relatively high prevalence found in this study reminds us that we should continue our efforts in addressing this issue.

3.12.6 Recommendations

1. Multiple factors are associated with the perpetuation and the victimization of bullying, violent behaviour and unintentional injuries among school-going adolescents. Although this study identified the prevalence and types of violent behaviour experienced, further studies may be conducted in these areas to better understand this problem. These should include risk factors such as health and sociocultural factors, help seeking behaviour and short and long term effects of engaging or being exposed to these situations.
2. Written policies and laws by themselves alone are not adequate to tackle the problem of physical attacks, physical fights, bullying, physical and verbal abuse experienced by school-going adolescents. Strategies should be developed with the involvement of parents, family, and the community. Support services for both perpetrator and victim at community and school levels should be developed. Increased efforts need to be directed towards male adolescents in the younger age group and within the Indian community. Helping them with counselling rather than punitive measures, besides early detection of underlying factors would aid better prevention strategies. Innovative measures including use of social media, need to be adapted in line with the problems faced by the current generation such as cyberbullying and online forms of abuse. An integrated, multifactorial and multisectoral approach from the family level to school environment is crucial in addressing this problem.

3. Unintentional injuries should be addressed by stepping up safety awareness programs. This should be targeted towards road and traffic safety, starting from the younger age group. This would help reduce the burden of avoidable and unintentional injuries. Measures to create a safe environment within the school, home and other areas should be implemented by all parties.

3.12.7 References

1. World Health Organization. Fact Sheet: Violence prevention in the South-East Asia Region. 2015.
2. Menesini E, Salmivalli C. Bullying in schools: the state of knowledge and effective interventions. *Psychology, health & medicine*. 2017;22(sup1):240-53.
3. Sittichai R, Smith PK. Bullying in south-east Asian countries: A review. *Aggression and Violent Behavior*. 2015;23:22-35.
4. Institute for Public Health (IPH). 2012. National Health and Morbidity Survey, NHMS 2012: Johor Global School-based Student Health Survey 2012. Kuala Lumpur Ministry of Health Malaysia.
5. Lee L, Chen PC, Lee K, Kaur J. Violence-related behaviours among Malaysian adolescents: a cross sectional survey among secondary school students in Negeri Sembilan. *Annals-Academy of Medicine Singapore*. 2007;36(3):169.

Table 3.12.1 : Prevalence of involvement in violence at least once in the past 12 months in Form 1 to Form 5 students in Johor, 2017

Socio-demographic Characteristics	Having Been Physically Attacked At Least Once					Involvement In Physical Fight At Least Once				
	Unweighted Count	Estimated Population	Prevalence (%)	95% CI		Unweighted Count	Estimated Population	Prevalence (%)	95% CI	
				Lower	Upper				Lower	Upper
JOHOR	438	71,790	26.1	22.15	30.47	357	58,798	21.3	17.62	25.62
Locality of school										
Urban	296	48,261	24.7	19.78	30.29	245	40,757	20.8	16.22	26.27
Rural	142	23,528	29.7	25.33	34.38	112	18,040	22.7	17.18	29.37
Sex										
Male	258	46,260	33.8	28.44	39.60	211	38,158	27.8	23.34	32.80
Female	180	25,529	18.5	14.66	23.03	146	20,640	14.9	11.62	18.96
Form										
Form 1	113	20,012	34.7	26.34	44.09	103	18,300	31.7	24.86	39.47
Form 2	130	17,634	33.2	29.51	37.05	109	14,872	27.9	21.70	35.08
Form 3	63	10,803	19.9	13.92	27.68	59	9,975	18.4	12.25	26.70
Form 4	79	12,814	23.7	15.72	34.12	49	7,663	14.2	8.84	21.97
Form 5	53	10,527	18.8	13.86	25.03	37	7,988	14.2	8.48	22.85

Table 3.12.2 : Prevalence of being bullied at least once in the past 30 days among Form 1 to Form 5 students in Johor, 2017

Socio-demographic Characteristics	Having Been Bullied At Least Once				
	Unweighted Count	Estimated Population	Prevalence (%)	95% CI	
				Lower	Upper
JOHOR	238	38,243	13.9	10.89	17.54
Locality of school					
Urban	157	24,986	12.8	9.10	17.59
Rural	81	13,256	16.7	13.35	20.65
Sex					
Male	133	23,295	17.0	13.62	21.08
Female	105	14,947	10.8	7.81	14.72
Form					
Form 1	51	9,147	15.8	11.06	22.16
Form 2	81	11,438	21.5	13.14	33.18
Form 3	46	7,688	14.2	10.16	19.45
Form 4	45	6,808	12.6	9.40	16.69
Form 5	15	3,162	5.6	3.39	9.20

Table 3.12.3: Most common ways of being bullied in the past 30 days among students Form 1 to Form 5 in Johor, 2017

	Unweighted Percentage		95% CI	
	Count	(%)	Lower	Upper
Hit, kicked, pushed, shoved around or locked indoors	14	6.3	3.04	12.73
Made fun of because of race, nationality or colour	18	7.6	3.90	14.34
Made fun of because of religion	5	2.5	1.01	6.15
Made fun of with sexual jokes, comments or gestures	41	17.6	10.81	27.33
Left out of activities on purpose or completely ignored	16	6.2	3.12	12.00
Made fun of because of how body or face looks	38	16.2	11.99	21.64

Table 3.12.4: Prevalence of being abused at least once in the past 30 day among Form 1 to Form 5 students in Johor, 2017

Socio-demographic Characteristics	Physical Abuse At Home At Least Once					Verbal Abuse At Home At Least Once				
	Unweighted Count	Estimated Population	Prevalence (%)	95% CI		Unweighted Count	Estimated Population	Prevalence (%)	95% CI	
				Lower	Upper				Lower	Upper
JOHOR	152	23,731	8.6	5.59	13.02	784	124,016	45.2	38.78	51.71
Locality of school										
Urban	108	17,007	8.7	5.11	14.32	523	82,179	42.1	34.49	50.02
Rural	44	6,724	8.5	4.02	16.92	261	41,837	52.8	45.89	59.64
Sex										
Male	60	10,492	7.7	4.77	12.04	291	53,154	38.8	32.79	45.27
Female	92	13,239	9.6	5.79	15.38	493	70,862	51.4	41.98	60.79
Form										
Form 1	44	7,655	13.2	7.97	21.11	155	26,028	45.2	35.49	55.23
Form 2	54	7,497	14.1	7.50	24.82	172	23,092	43.8	36.46	51.36
Form 3	17	2,535	4.7	2.67	8.07	152	24,267	44.8	37.18	52.60
Form 4	24	3,507	6.5	3.64	11.31	176	26,097	48.5	38.97	58.16
Form 5	13	2,538	4.5	1.84	10.65	129	24,532	43.6	34.33	53.44

Table 3.12.5: Prevalence of having had serious injury at least once in the past 12 months Form 1 to Form 5 students in Johor, 2017

Socio-demographic Characteristics	Having Had Serious Injury At Least Once				
	Unweighted Count	Estimated Population	Prevalence (%)	95% CI	
				Lower	Upper
JOHOR	443	72,305	26.2	22.20	30.70
Locality of school					
Urban	314	50,683	25.8	20.59	31.86
Rural	129	21,623	27.2	22.70	32.24
Sex					
Male	228	41,930	30.6	25.20	36.53
Female	215	30,375	21.9	18.03	26.38
Form					
Form 1	95	17,063	29.4	23.23	36.52
Form 2	119	16,286	30.6	23.29	38.95
Form 3	82	13,331	24.6	18.63	31.71
Form 4	90	13,717	25.4	21.14	30.16
Form 5	57	11,907	21.2	15.62	28.08

Table 3.12.6: Types of the most serious injury sustained in the past 12 months among students who were injured among Form 1 to Form 5 students in Johor, 2017

	Unweighted Percentage		95% CI	
	Count	(%)	Lower	Upper
Broken bone / dislocated Joint	90	20.4	15.45	26.54
A cut or stab wound	85	19.9	15.89	24.57
Concussion / head or neck injury, knocked out or could not breathe	25	5.4	3.32	8.56
Bad burn	3	0.7	0.25	2.02
Poisoned	3	0.6	0.20	1.69

Table 3.12.7: Major cause of the most serious injury sustained in the past 12 months among students who were injured among Form 1 to Form 5 students in Johor, 2017

	Unweighted Percentage		95% CI	
	Count	(%)	Lower	Upper
In a motor vehicle accident or hit by a motor vehicle	64	15.2	11.07	20.60
Fell	169	37.9	31.84	44.31
Something fell or hit him/her	28	5.9	3.78	9.02
Attacked or abused or fighting with someone	9	2.3	1.05	4.87
In a fire or too near a flame or something hot	4	1.0	0.32	2.95
Inhaled or swallowed something bad	6	1.4	0.70	2.99

APPENDICES

Appendix 1: Members of Steering Committee NHMS 2015-2018

1. Director General of Health
2. Deputy Director General of Health (Public Health)
3. Deputy Director General of Health (Medical)
4. Deputy Director General of Health (Research & Tech. Support)
5. Director, Oral Health Division
6. Director, Pharmaceutical Services Division
7. Director, Food Safety and Quality Programme Division
8. Director, Medical Development Division
9. Director, Planning Division
10. Director, Health Education Division
11. Director, Disease Control Division
12. Director, Family Health Development Division
13. Director, Nutrition Division
14. Representative of State Directors
15. Director, Institute for Public Health
16. Dean Faculty of Medicine, University of Malaya
17. Dean Faculty of Medicine, National University of Malaysia
18. Principle Investigator, NHMS

Appendix 2: Term of Reference for NHMS 2015-2018 Steering Committee

1. To approve the objectives and scopes of NHMS 2015-2018.
2. To facilitate inter and intra-sectorial collaboration.
3. To monitor the implementation of the NHMS 2015-2018.
4. To facilitate the utilisation of the NHMS 2015-2018 findings.

Appendix 3: List of members of Central Coordinating Committee, NHMS 2017

1. Dr. Hj Tahir bin Aris, Director of Institute for Public Health
2. Dr. Muhammad Fadhli bin Mohd Yusoff, Coordinator of NHMS 2015-2018
3. Dr. S Maria binti Awaluddin, Principal Investigator of Adolescent Health Survey
4. Pn. Ruhaya binti Salleh, Principal Investigator of Adolescent Nutrition Survey
5. En. Mohamad Aznuddin bin Abd Razak, Principal Investigator of Healthy Mind Screening using DASS
6. Dr. Mohd Azahadi bin Omar, Head Data Processing and Data Management
7. Dr. Noor Ani binti Ahmad, Central Field Supervisor of Perlis & Kedah
8. Dr. Nor Asiah binti Mohamad, Central Field Supervisor of Johor, Melaka & Negeri Sembilan
9. Dr. Rajini a/p Sooryanarayana, Data Processing & Quality
- 10.Pn. Tee Guat Hiong, Central Field Supervisor of Sarawak
- 11.Dr. Nur Liana binti Ab. Majid, Central Field Supervisor of WP Kuala Lumpur, WP Putrajaya & Selangor
- 12.Cik Hasimah binti Ismail, Central Field Supervisor of Pahang, Kelantan & Terengganu
- 13.Pn. Norzawati binti Yoep, Central Field Supervisor of Perak & Kedah
- 14.Pn. Norazizah binti Ibrahim Wong, Data Processing & Quality
- 15.En. Mohd Hazrin bin Hasim @ Hashim, Central Field Supervisor of WP Labuan & Sabah
- 16.Pn. Lalitha a/p Palanivello, Person in charge for 24Hour Diet Recall
- 17.Pn. Siti Nor'Ain binti Hashim, Head of ICT Support
- 18.En. Lim Kuang Kuay, Logistic Support
- 19.Pn. Hamizatul Akmal binti Abd. Hamid, Project Manager
- 20.Pn. Wan Shakira binti Rodzlan Hasani, Project Manager
- 21.Pn. Cheong Siew Man, Person in-charge for Food Frequency Questionnaire
- 22.Pn. Nazirah Bt Alias, Data Processing & Quality
- 23.Dr. Fazila Haryati Ahmad, Data Processing &Quality

Appendix 4: Terms of Reference for NHMS 2017 Central Coordinating Team

No	Team	Duties	Officers
1	Project Management and Finance	Work closely with recruitment group for employment of RA Prepare Questionnaires manual, Data collection manual Meeting with Liason Officers Planning for data collection training Prepare security cards/name tags for research team Arrangement for advanced payment for team managers, nurses and drivers Process claims of MOH staff Prepare tickets for travelling Monitor the expenditure/budget	Dr. Muhammad Fadhli bin Mohd Yusoff Dr. S. Maria Binti Awaluddin Pn. Hamizatul Akmal binti Abd Hamid Pn. Wan Shakira binti Rodzlan Hasani Cik Nur Hazwani Binti Mohd Hasri
2	Survey Research Centre	Calculate the sample size Determine the sample distribution by state	Dr. Muhammad Fadhli bin Mohd Yusoff Pn. Norazizah binti Ibrahim Wong Pn. Wan Shakira binti Rodzlan Hasani
3	ICT Unit	Maintenance of the scanning machine Daily back up for databases	Pn. Siti Nor'ain Binti Hashim En. Sulaiman Bin Harun En. Yusmirol Bin Yusop En. Andy Bin Mustaming

4	Central Field Supervisors	<p><u>Before Data Collection</u></p> <p>Central Field Supervisors are expected to prepare for the initiation of data collection. The preparation tasks include: Conduct meeting with State Education Office, School Principals, Teacher in-charged for the selected schools.</p> <p>To ensure adequate logistic support for the data collection and liaise with the District Education Office, District Health Office and other relevant departments to ensure that:</p> <ul style="list-style-type: none"> • Human resources are available: Field Supervisors, Team leaders, Research Assistants and drivers. • Manage transport: Vehicles • Manage survey instruments and relevant form • Manage lodging for data collectors <p><u>During Data Collection</u></p> <p>Gather feedback from the field on the data collection status and problems related to logistics.</p> <p>Visit the field to help data collectors solve the problem if necessary.</p> <p>To ensure all data collection monitoring forms have been received on time.</p> <p>To ensure bundle from field received by the Operation Centre by hand and by post (Sabah, Sarawak, WP Labuan)</p> <p>Updating the monitoring board for state achievement and attending CCT meeting.</p>	<p>Dr Nor Asiah Binti Muhamad</p> <p>Dr Nur Liana Binti Ab Majid</p> <p>Pn. Norzawati Binti Yeop</p> <p>Dr. Noor Ani Binti Ahmad</p> <p>Pn. Hasimah Binti Ismail</p> <p>Pn. Tee Guat Hiong</p> <p>En. Mohd Hazrin Bin Hasim @ Hashim</p>
5	Data Processing and management	<p>Setting up data processing facility</p> <p>Development of directory of variables database</p>	<p>Dr. Mohd Azahadi bin Omar</p>

		<p>Development of quality control (QC) manual for data processing</p> <p>Specify data structure for data processing and data output requirement</p> <p>Responsible for data entry and data cleaning</p> <p>Monitoring and evaluation of QC performance for data processing</p>	<p>Dr. Rajini a/p sooryanarayana</p> <p>Dr. Fazila Haryati Binti Ahmad</p> <p>Pn. Nazirah Binti Alias</p>
6	Operation Centre	<p>Arrange date and place of meeting</p> <p>Prepare and circulate briefing materials</p> <p>Prepare and circulate minutes of CCT meeting</p> <p>Prepare letters of appointment to state liaison officers, nurses, scouts and data collectors</p> <p>Prepare advertisement material for recruitment of data collectors, team leaders, and interviewers,</p> <p>Prepare letters of notifications for data collections</p> <p>Prepare manuals for field Supervisors and data collectors</p> <p>Develop a system/format and monitor the distribution of materials/equipment for field work</p> <p>Arrange transport/drivers for distribution and collection of materials</p>	<p>Pn. Hamizatul Akmal binti Abd Hamid</p> <p>En. Azli bin Baharudin</p> <p>Cik Nur Hazwani Binti Mohd Hasri</p> <p>Pn. Siti Noafika Binti Anwar</p> <p>En. Muhammad Suhaimi Bin Mohamad Idrus</p> <p>Cik Shahibul Bariah binti Mat Ghani</p> <p>Pn. Nur Fadzilla binti Mohd Radzi</p> <p>En. Muhammad Zuhdi Bin Khiruddin</p> <p>Cik Nurbaiti Binti Asmawi</p>

Appendix 5: List of Research Team Members, NHMS 2017

Alcohol Use

1. Dr. Muhammad Fadhli Mohd Yusoff
2. Dr. Tania Gayle Robert
3. Dr. Halizah Mat Rifin
4. Dr. Norli Abdul Jabbar
5. Dr. Rozanim Kamaruddin
6. Dr. Jane Ling Miaw Yn
7. Ms. Hasimah Ismail
8. Ms. Hamizatul Akmal Binti Abd Hamid
9. Mr. Mohd Hatta Mutalip
10. Ms. Wan Shakira Rodzlan Hasani

Dietary Behaviours

1. Ms. Rashidah Ambak
2. Ms. Ruhaya Salleh
3. Ms. Norlida Zulkafly
4. Dr. S. Maria Awaluddin
5. Ms. Rusidah Selamat
6. Ms. Syafinaz Mohd Sallehuddin
7. Mr. Mohd Hasnan Ahmad
8. Ms. Cheong Siew Man

Drug Use

1. Dr. Muhammad Fadhli Mohd Yusof
2. Dr. Rushidi Ramly
3. Dr. Norli Abdul Jabbar

4. Dr. Jane Ling Miaw Yn
5. Dr. Halizah Mat Rifin
6. Dr. Tania Gayle Robert
7. Dr. Thamil Arasu Saminathan
8. Dr. Nur Liana Ab. Majid
9. Ms. Hasimah Ismail
10. Ms. Hamizatul Akmal Abd Hamid
11. Ms. Wan Shakira Rodzlan Hasani

Hygiene (Including Oral Health)

1. Dr. Yaw Siew Lian
2. Dr. Nurrul Ashikin Abdullah
3. Dr. Natifah Che Salleh
4. Ms. Norazizah Ibrahim Wong
5. Mr. Mohamad Fuad Mohamad Anuar
6. Mr. Sayan a/l Pan

Internet Use and Addiction

1. Dr S. Maria Awaluddin
2. Prof Madya Dr. Normala Ibrahim
3. Ms. Chan Ying Ying
4. Dr. Rimah Melati Abd. Ghani
5. Dr. Amal Shamsuddin
6. Prof Madya Dr. Wan Salwina Wan Ismail
7. Dr. Norharlina Bahar
8. Dr. Saidatul Norbaya Binti Buang
9. Dr. Nik Rubiah Nik Abd. Rashid

Mental Health Problems

1. Dr. Noor Ani Ahmad
2. Prof Dr. Sherina Mohd Sidik
3. Dr. Fazly Azry Abdul Aziz
4. Ms. Noraida Mohamad Kasim
5. Mr. Mohammad Aznuddin Abd Razak
6. Ms. Muslimah Yusof

Physical Activity

1. Mr. Lim Kuang Kuay
2. Dr. Hj. Mohd Azahadi Omar
3. Ms. Chan Ying Ying
4. Dr. Chandrika A/p Jeevananthan
5. Mr. Azli Baharudin @Shaharuddin
6. Ms. Nazirah Alias

Protective Factors

1. Pn Norzawati binti Yoep
2. Ms. Faizah Paiwai
3. Dr. Noor Aliza Lodz
4. Dr. S. Maria Awaluddin
5. Ms. Azna Ahmad
6. Dr. Nik Rubiah Nik Abd Rashid

Sexual Behaviour that contribute to HIV infection, other STI and unintended pregnancy

1. Dr. Maisarah Omar
2. Dr. Noor Aliza Lodz
3. Dr. S.Maria Awaluddin
4. Dr. Noor Ani Ahmad

5. Dr. Nik Rubiah Nik Abdul Rashid
6. Ms. Noraida Mohamad Kasim

Tobacco Use

1. Dr. Noraryana Hassan
2. Dr. Nizam Baharom
3. Dr. Muhammad Fadhli Mohd Yusoff,
4. Dr. Abdul Aiman Abd Ghani
5. Dr. Nur Liana Ab. Majid
6. Dr. Thamil Arasu Saminathan
7. Dr. Jane Ling Miaw Yn
8. Dr. Tania Gayle Robert
9. Dr. Halizah Mat Rifin
10. Ms. Tee Guat Hiong
11. Ms. Hasimah Ismail
12. Ms. Hamizatul Akmal Abd Hamid
13. Ms. Wan Shakira Rodzlan Hasani

Violence and Unintentional Injury

1. Dr. Rajini Sooryanarayana
2. Dr. Shubash Shander Ganapathy
3. Dr. Muhammad Fadhli Mohd Yusoff
4. Dr. S Maria Awaluddin
5. Dr. Thamil Arasu a/l Saminathan
6. Dr. Azriman Rosman
7. Dr. Fazila Haryati Ahmad
8. Mr. Mohamad Fuad Mohamad Anuar
9. Mr. Mohd Hazrin Bin Hasim@Hashim

Appendix 6: List of Liaison Officer and Data Collection Teams

JOHOR

Liaison Officer

Mr. Yahya Bin Ahmad

Field Supervisor

Ms. Fatimah Binti Othman

Nutritionist

1. Ms. Nik Normasitah Binti Nik Ismail
2. Mr. Mohd Razif Bin Mohd Jamain
3. Ms. Siti Dinie Syazwani Binti Azlam
4. Ms. Lee Pei Shan
5. Mr. Mohd Zaid Bin Ramlan

Drivers

1. Mr. Osman Bin Mohd
2. Mr. Mohd Safar Bin Sarbani

Research Assistants

1. Mr. Mohd Nazrulshah Bin Salim
2. Ms. Sharifah Solihah Binti Syed Hashim
3. Mr. Khairulhanafi Bin Muhammad Khair
4. Ms. Nur Hamizah Binti Md Sha'eb
5. Ms. Siti Aisyah Binti Ibrahim
6. Ms. Nor Hamizah Binti Atan
7. Ms. Siti Nur Nadiah Binti Khozaini
8. Ms. Nurul Siddiqah Binti Whakiddin

Appendix 7: Questionnaire & OMR

iku
INSTITUTE FOR PUBLIC HEALTH

**KEMENTERIAN
PENDIDIKAN
MALAYSIA**

**TINJAUAN KEBANGSAAN
KESIHATAN DAN MORBIDITI** **NHMS**
NATIONAL HEALTH AND **2017**
MORBIDITY SURVEY

**TINJAUAN KESIHATAN REMAJA
BORANG SOAL SELIDIK**

**SILA KEMBALIKAN
BORANG SOAL SELIDIK INI
SELEPAS MENJAWAB**

**JANGAN CONTENG
BORANG SOAL SELIDIK INI**

PENGENALAN

Kementerian Kesihatan Malaysia dengan kerjasama Kementerian Pendidikan Malaysia sedang menjalankan tinjauan yang berkaitan kesihatan remaja. Tinjauan ini adalah berkenaan dengan kesihatan anda dan tindakan yang anda ambil yang memberi kesan kepada kesihatan anda.

Terima kasih kerana bersetuju untuk menyertai tinjauan ini. Tiada jawapan yang **BETUL** atau **SALAH**. Markah peperiksaan anda tidak akan terjejas sekiranya anda menjawab atau tidak menjawab soalan dalam tinjauan ini. Setiap jawapan yang anda berikan boleh membantu memantapkan program kesihatan untuk remaja. Oleh itu, sila jawab dengan ikhlas dan tepat.

Segala maklumat individu yang diberikan adalah RAHSIA kerana TIADA MAKLUMAT PENGENALAN DIRI DIAMBIL dan TIDAK AKAN DIDEBAHKAN. Tiada siapa akan tahu jawapan anda. Sila jawab berdasarkan apa yang anda tahu.

INTRODUCTION

The Ministry of Health Malaysia, with the cooperation of the Ministry of Education Malaysia is conducting a survey on adolescent health. This survey is about your health and the things that you do that may affect your health.

*Thank you for agreeing to participate in this survey. There are no **CORRECT** or **WRONG** answers. None of your grades or marks will be affected whether or not you answer the questions. Each answer that you provide will help in the improvisation of health programs for adolescents. Therefore, please answer as honestly and accurately as possible.*

*All individual information given will be kept **SECRET** because **NO IDENTIFIERS ARE TAKEN** nor **WILL BE EXPOSED**. No one will know your answer. Answer the questions based on the best of your knowledge.*

PANDUAN MENGISI BORANG (SILA BACA DENGAN TELITI)

GUIDE ON FILLING THE QUESTIONNAIRE (PLEASE READ CAREFULLY)

a. **JANGAN** tulis **NAMA ANDA** pada kertas soalan mahupun kertas jawapan.

DO NOT write **YOUR NAME** on the questionnaire or the answer sheet.

b. Sila **BACA PERNYATAAN** untuk soalan yang memberikan pernyataan atau definisi sebelum menjawab soalan.

Please **READ STATEMENT** for questions with a preceding statement or definition before answering.

c. Sila **HITAMKAN** jawapan yang bersamaan jawapan anda pada kertas jawapan yang disediakan. Hanya hitamkan **SATU JAWAPAN** bagi setiap soalan. Sila gunakan pensel 2B yang disediakan. Sekiranya telah selesai, ikut arahan pegawai yang melakukan survei di sekolah/kelas anda.

Please **FILL IN THE CIRCLES** on your answer sheet that matches your answer. There can be **ONLY ONE ANSWER** for each question. Use only the provided 2B pencil. When you are done, follow the instructions of the person conducting the survey in your school/class

BERIKUT ADALAH CONTOH BAGAIMANA MENGISI JAWAPAN

HERE IS AN EXAMPLE OF HOW TO FILL IN THE CIRCLE:

ISIKAN BULATAN SEPERTI INI
 FILL IN THE CIRCLES LIKE THIS



BUKAN SEPERTI INI
 NOT LIKE THIS



ATAU
 OR



TINJAUAN/SURVEY

1. Adakah ikan tinggal dalam air?

Do fish live in the water?

A. Ya

Yes

B. Tidak

No

Borang jawapan
 Answer sheet



BAHAGIAN 1
PART 1

1. Berapakah umur anda?
 How old are you?
 - a. 11 tahun atau ke bawah
11 years old or younger
 - b. 12 tahun
12 years old
 - c. 13 tahun
13 years old
 - d. 14 tahun
14 years old
 - e. 15 tahun
15 years old
 - f. 16 tahun
16 years old
 - g. 17 tahun
17 years old
 - h. 18 tahun atau ke atas
18 years old or older

2. Apakah jantina anda?
 What is your sex?
 - a. Lelaki
Male
 - b. Perempuan
Female

3. Anda belajar di tingkatan/kelas apa?
 In what form/class are you?
 - a. Kelas peralihan
Remove class
 - b. Tingkatan 1
Form 1
 - c. Tingkatan 2
Form 2
 - d. Tingkatan 3
Form 3
 - e. Tingkatan 4
Form 4
 - f. Tingkatan 5
Form 5

4. Apakah etnik anda?
 What is your ethnicity?
 - a. Melayu
Malay
 - b. Cina
Chinese
 - c. India
Indian
 - d. Bumiputera Sabah
Bumiputera Sabah
 - e. Bumiputera Sarawak
Bumiputera Sarawak
 - f. Lain-lain etnik
Some other ethnic

5. Apakah status perkahwinan ibu bapa anda?
 What is the marital status of your parents?
 - a. Berkahwin dan tinggal bersama
Married and living together
 - b. Berkahwin tetapi tidak tinggal bersama atas dasar bekerja di tempat lain
Married but living apart due to working in other place
 - c. Berceraai
Divorced
 - d. Balu (ayah atau ibu telah meninggal)
Widower (my mother or father has died)
 - e. Berpisah (ibu bapa tidak tinggal serumah)
Separated (my parent do not live together)
 - f. Tidak tahu
I do not know

BAHAGIAN 2
PART 2

6. Dalam tempoh 30 hari yang lepas, berapa kerap anda kelaparan kerana tidak cukup makanan di rumah?

During the past 30 days, how often did you go hungry because there was not enough food in your home?

- a. Tidak pernah
Never
- b. Jarang-jarang
Rarely
- c. Kadang-kadang
Sometimes
- d. Kebanyakan masa
Most of the time
- e. Sentiasa
Always

Enam soalan yang berikutnya adalah berkenaan apa yang anda mungkin makan atau minum.

The next six questions ask about what you might eat and drink.

7. Dalam tempoh 30 hari yang lepas, berapa kali dalam sehari anda **biasanya** makan buah?

*During the past 30 days, how many times per day did you **usually** eat fruits?*

- a. Saya tidak makan buah dalam 30 hari yang lepas
I did not eat fruits during the past 30 days
- b. Kurang dari 1 kali sehari
Less than 1 time per day
- c. 1 kali sehari
1 time per day
- d. 2 kali sehari
2 times per day
- e. 3 kali sehari
3 times per day
- f. 4 kali sehari
4 times per day
- g. 5 kali atau lebih sehari
5 or more times per day

8. Dalam tempoh 30 hari yang lepas, berapa kali dalam sehari anda **biasanya** makan sayur?

*During the past 30 days, how many times per day did you **usually** eat vegetables?*

- a. Saya tidak makan sayur dalam 30 hari yang lepas
I did not eat vegetables during the past 30 days
- b. Kurang dari 1 kali sehari
Less than 1 time per day
- c. 1 kali sehari
1 time per day
- d. 2 kali sehari
2 times per day
- e. 3 kali sehari
3 times per day
- f. 4 kali sehari
4 times per day
- g. 5 kali atau lebih sehari
5 or more times per day

9. Dalam tempoh 30 hari yang lepas, berapa kali dalam sehari anda **biasanya** minum air berkarbonat seperti Coca Cola, Sprite, Pepsi dan lain-lain? (Air berkarbonat diet tidak termasuk dalam kumpulan ini)

*During the past 30 days, how many times per day did you **usually** drink carbonated soft drinks such as Coca Cola, Sprite, and Pepsi? (Do not include diet soft drinks)*

- a. Saya tidak minum air berkarbonat dalam 30 hari yang lepas
I did not drink carbonated soft drink during the past 30 days
- b. Kurang dari 1 kali sehari
Less than 1 time per day
- c. 1 kali sehari
1 time per day
- d. 2 kali sehari
2 times per day
- e. 3 kali sehari
3 times per day
- f. 4 kali sehari
4 times per day
- g. 5 kali atau lebih sehari
5 or more times per day

10. Dalam tempoh 30 hari yang lepas, berapa kali dalam sehari anda **biasanya** minum air kosong seperti air mineral, air masak atau air paip?

*During the past 30 days, how many times per day did you **usually** drink plain water such as mineral water, boiled water, or tap water?*

- a. Saya tidak minum air kosong dalam 30 hari yang lepas
I did not drink plain water during the past 30 days
- b. Kurang dari 1 kali sehari
Less than 1 time per day
- c. 1 kali sehari
1 time per day
- d. 2 kali sehari
2 times per day
- e. 3 kali sehari
3 times per day
- f. 4 kali sehari
4 times per day
- g. 5 kali atau lebih sehari
5 or more times per day

11. Dalam tempoh 30 hari yang lepas, berapa kali dalam sehari anda **biasanya** minum susu atau makan produk tenusu seperti susu, keju, yogurt dan lain-lain? (Ini tidak termasuk susu pekat manis)

*During the past 30 days, how many times per day did you **usually** drink milk or eat milk products, such as milk, cheese, and yogurt? (This does not include sweetened condensed milk)*

- a. Saya tidak minum susu atau makan produk tenusu dalam 30 hari yang lepas

I did not drink milk or eat milks products during the past 30 days

- b. Kurang dari 1 kali sehari
Less than 1 time per day
- c. 1 kali sehari
1 time per day
- d. 2 kali sehari
2 times per day
- e. 3 kali sehari
3 times per day
- f. 4 kali sehari
4 times per day
- g. 5 kali atau lebih sehari
5 or more times per day

12. Dalam tempoh 7 hari yang lepas, berapa hari anda makan makanan segera dari restoran **makanan segera** seperti McDonald, KFC, dan Pizza Hut?

*During the past 7 days, how many days did you eat food from a **fast food** restaurant, such as McDonalds, KFC and Pizza Hut?*

- a. 0 hari
0 day
- b. 1 hari
1 day
- c. 2 hari
2 days
- d. 3 hari
3 days
- e. 4 hari
4 days
- f. 5 hari
5 days
- g. 6 hari
6 days
- h. 7 hari
7 days

BAHAGIAN 3
PART 3

6 soalan seterusnya adalah berkenaan amalan pembersihan gigi, lawatan pergigian anda dan juga bagaimana gigi anda memberi kesan kepada aktiviti-aktiviti anda.

The 6 next question are about the cleaning of your teeth, your dental visits and also how your teeth affect your activities.

13. Dalam tempoh 30 hari yang lepas, berapa kali dalam sehari anda membersihkan atau memberus gigi anda?
During the past 30 days, how many times per day did you usually clean or brush your teeth?
- Saya tidak membersihkan atau memberus gigi dalam 30 hari yang lepas
I did not clean or brush my teeth during the past 30 days
 - Kurang dari 1 kali sehari
Less than 1 time per day
 - 1 kali sehari
1 time per day
 - 2 kali sehari
2 times per day
 - 3 kali sehari
3 times per day
 - 4 kali atau lebih dalam sehari
4 or more times per day
14. Adakah anda menggunakan ubat gigi berflorida?
Do you use toothpaste that contain fluoride?
- Ya
Yes
 - Tidak
No
 - Tidak tahu
I do not know
15. Adakah anda menggunakan flos/benang gigi untuk membersihkan gigi anda?
Do you use dental floss to clean your teeth?
- Ya
Yes
 - Tidak
No
16. Bilakah kali terakhir anda berjumpa doktor gigi atau jururawat pergigian untuk pemeriksaan, pembersihan gigi atau rawatan pergigian yang lain?
When was the last time you saw a dentist or dental nurse for a check-up, teeth cleaning, or other dental treatment?
- Dalam tempoh 12 bulan yang lepas
During the past 12 months
 - Di antara 12 hingga 24 bulan yang lepas
Between 12 and 24 months ago
 - Lebih daripada 24 bulan yang lepas
More than 24 months ago
 - Tidak pernah
Never
 - Tidak tahu
I do not know
17. Dalam tempoh 12 bulan yang lepas, adakah sakit gigi menyebabkan anda tidak hadir ke kelas atau sekolah?
During the past 12 months, did a toothache cause you to miss classes or school?
- Ya
Yes
 - Tidak
No
18. Adakah anda mengelak untuk senyum atau ketawa kerana risau dengan rupa gigi anda?
Do you avoid smiling or laughing because how your teeth look?
- Ya
Yes
 - Tidak
No

**3 soalan seterusnya adalah berkenaan amalan membasuh tangan.
 The next 3 questions are about your hand washing practices.**

19. Dalam tempoh 30 hari yang lepas, berapa kerap anda menggunakan sabun semasa membasuh tangan anda?

During the past 30 days, how often did you use soap when washing your hands?

- a. Tidak pernah
Never
- b. Jarang-jarang
Rarely
- c. Kadang-kadang
Sometimes
- d. Kebanyakan masa
Most of the time
- e. Setiap kali
Always

20. Dalam tempoh 30 hari yang lepas, berapa kerap anda membasuh tangan sebelum makan?

During the past 30 days, how often did you wash your hands before eating?

- a. Tidak pernah
Never
- b. Jarang-jarang
Rarely
- c. Kadang-kadang
Sometimes
- d. Kebanyakan masa
Most of the time
- e. Setiap kali
Always

21. Dalam tempoh 30 hari yang lepas, berapa kerap anda membasuh tangan selepas menggunakan tandas?

During the past 30 days, how often did you wash your hands after using the toilet?

- a. Tidak pernah
Never
- b. Jarang-jarang
Rarely
- c. Kadang-kadang
Sometimes
- d. Kebanyakan masa
Most of the time
- e. Setiap kali
Always

**BAHAGIAN 4
 PART 4**

SILA BACA PERNYATAAN DI BAWAH:

Soalan berikutnya adalah berkenaan serangan fizikal. Serangan fizikal berlaku apabila seseorang atau lebih ramai orang menyerang individu lain dengan anggota badan atau senjata seperti kayu dan pisau. Serangan fizikal tidak diambil kira jika dua individu atau pelajar yang sama saiz atau kekuatan bersetuju untuk bergaduh secara fizikal.

PLEASE READ THE STATEMENT BELOW:
The next question asks about physical attacks. A physical attack occurs when one or more people hurt another person with/without a weapon such as sticks and knife. It is not a physical attack when two individuals or students of about the same strength or power choose to fight each other.

22. Dalam tempoh 12 bulan yang lepas, berapa kali anda telah diserang secara fizikal?

During the past 12 months, how many times were you physically attacked?

- a. 0 kali
0 times
- b. 1 kali
1 time
- c. 2 atau 3 kali
2 or 3 times
- d. 4 atau 5 kali
4 or 5 times
- e. 6 atau 7 kali
6 or 7 times
- f. 8 atau 9 kali
8 or 9 times
- g. 10 atau 11 kali
10 or 11 times
- h. 12 kali atau lebih
12 or more times

SILA BACA PERNYATAAN DI BAWAH:

Soalan seterusnya adalah berkaitan pergaduhan fizikal. Pergaduhan fizikal berlaku apabila dua individu atau pelajar yang sama saiz atau kekuatan bersetuju untuk bergaduh secara fizikal.

PLEASE READ THE STATEMENT BELOW:
The next question asks about physical fights. A physical fight occurs when two individuals or students of about the same strength or power choose to fight each other.

23. Dalam tempoh 12 bulan yang lepas, berapa kali anda terlibat dalam pergaduhan secara fizikal?

During the past 12 months, how many times were you in physical fight?

- a. 0 kali
0 times
- b. 1 kali
1 time
- c. 2 atau 3 kali
2 or 3 times
- d. 4 atau 5 kali
4 or 5 times
- e. 6 atau 7 kali
6 or 7 times
- f. 8 atau 9 kali
8 or 9 times
- g. 10 atau 11 kali
10 or 11 times
- h. 12 kali atau lebih
12 or more times

SILA BACA PERNYATAAN DI BAWAH:

3 soalan seterusnya adalah berkenaan kecederaan serius yang pernah anda alami. Kecederaan serius berlaku apabila anda tidak hadir sekurang-kurangnya satu hari aktiviti yang biasa dilakukan (termasuk sekolah, sukan atau kerja) atau kecederaan yang memerlukan rawatan daripada doktor atau anggota kesihatan.

PLEASE READ THE STATEMENT BELOW:

The next 3 questions ask you about serious injuries that happened to you. An injury is serious when it makes you miss at least one full day of usual activities (such as school, sports or a job) or requires treatment by a doctor or medical personnel.

24. Dalam tempoh 12 bulan yang lepas, berapa kali anda mengalami kecederaan serius?

During the past 12 months, how many times were you seriously injured?

- a. 0 kali
0 times
- b. 1 kali
1 time
- c. 2 atau 3 kali
2 or 3 times
- d. 4 atau 5 kali
4 or 5 times
- e. 6 atau 7 kali
6 or 7 times
- f. 8 atau 9 kali
8 or 9 times
- g. 10 atau 11 kali
10 or 11 times
- h. 12 kali atau lebih
12 or more times

25. Dalam tempoh 12 bulan yang lepas, apakah kecederaan yang **paling serius** yang anda pernah alami?

*During the past 12 months, what was the **most serious** injury that happened to you?*

- a. Saya tidak mengalami kecederaan dalam tempoh 12 bulan yang lepas
I was not seriously injured during the past 12 months
- b. Patah tulang atau sendi terkehel/terkeluar
I had a broken bone or a dislocated joint
- c. Luka atau tikaman
I had a cut or stab wound
- d. Gegaran (konkusi) kepala atau kecederaan leher, pengsan atau tidak boleh bernafas
I had a concussion or other head or neck injury, was knocked out or could not breathe
- e. Kebakaran kulit yang serius
I had a bad burn
- f. Diracun atau mengambil ubat berlebihan
I was poisoned or took too much of a drug
- g. Sesuatu yang lain berlaku kepada saya
Something else happened to me

26. Dalam tempoh 12 bulan yang lepas, apakah penyebab utama terhadap kecederaan serius yang anda alami?

During the past 12 months, what was the major cause of the most serious injury that happened to you?

- a. Saya tidak mengalami kecederaan dalam 12 bulan yang lepas
I was not seriously injured during the past 12 months
- b. Saya terlibat dalam kemalangan melibatkan kenderaan
I was in a motor vehicle accident or hit by a motor vehicle
- c. Saya terjatuh
I fell
- d. Sesuatu telah jatuh atau terkena saya
Something fell on me or hit me
- e. Saya telah diserang atau didera atau bergaduh dengan orang lain
I was attacked or abused or was fighting with someone
- f. Saya terlibat dalam kebakaran atau berada terlalu dekat dengan api atau sesuatu yang panas
I was in the fire or too near a flame or something hot
- g. Saya sedut atau telan sesuatu yang membahayakan saya
I inhaled or swallowed something bad for me
- h. Sesuatu yang lain menyebabkan kecederaan saya
Something else caused my injury

SILA BACA PERNYATAAN BI BAWAH:

2 soalan seterusnya adalah berkenaan buli. Buli berlaku apabila seseorang atau sekumpulan pelajar mengata atau melakukan sesuatu yang tidak menyenangkan pelajar lain. Seseorang juga boleh dikatakan dibuli apabila dia diejek secara berterusan atau dipulaukan dengan sengaja. Buli tidak diambil kira apabila dua pelajar yang sama saiz atau kekuatan bergaduh atau ejekan dilakukan secara bergurau atau berseronok bagi kedua-dua pihak.

**PLEASE READ THE STATEMENT BELOW:
 The next 2 questions ask about bullying. Bullying occurs when a student or group of students say or do bad and unpleasant things to another student. It is also bullying when a student is teased a lot in an unpleasant way or when a student is left out of things on purpose. It is not bullying when two students of about the same strength or power argue or fight or when teasing is done in a friendly and fun way.**

27. Dalam tempoh 30 hari yang lepas, berapa hari anda telah dibuli?

During the past 30 days, on how many days were you bullied?

- a. 0 hari
0 days
- b. 1 atau 2 hari
1 or 2 days
- c. 3 hingga 5 hari
3 to 5 days
- d. 6 hingga 9 hari
6 to 9 days
- e. 10 hingga 19 hari
10 to 19 days
- f. 20 hingga 29 hari
20 to 29 days
- g. Kesemua 30 hari
All 30 days

28. Dalam 30 hari yang lepas, apakah perlakuan buli **paling kerap** anda alami?
*During the past 30 days, how were you bullied **most often**?*
- Saya tidak dibuli dalam 30 hari yang lepas
I was not bullied during the past 30 days
 - Saya telah dipukul, ditendang, ditolak atau dikunci dalam suatu tempat
I was hit, kicked, punched, shoved around, or locked indoors
 - Saya telah diejek kerana bangsa, kerakyatan atau warna kulit saya
I was made fun of because of my race, nationality or colour
 - Saya telah diejek kerana agama saya
I was made fun of because of my religion
 - Saya telah diejek dengan ejekan seksual seperti secara komen, perilaku atau gurauan
I was made fun of with sexual jokes, comments or gestures
 - Saya telah dipulau dari apa-apa aktiviti secara sengaja atau langsung tidak dipedulikan
I was left out of activities on purpose or completely ignored
 - Saya diejek kerana bentuk badan atau paras rupa saya
I was made fun of because of how my body or face looks
 - Saya telah dibuli dengan cara lain
I was bullied in some other way

SILA BACA PERNYATAAN BI BAWAH:
 2 soalan berikutnya adalah berkenaan deraan fizikal dan lisan di rumah. Apabila seseorang memukul atau mengatakan ayat yang menyakitkan hati ia diambil kira sebagai deraan fizikal atau lisan.

PLEASE READ THE STATEMENT BELOW:
 The next 2 questions ask about physical and verbal abuse at home. When someone hits you or says hurtful or insulting things to you it is called physical abuse or verbal abuse.

29. Dalam tempoh 30 hari yang lepas, adakah anda telah dipukul di rumah sehingga meninggalkan kesan atau mengalami kecederaan?
During the past 30 days, how many times did someone at home hit you so hard that they left a mark or caused an injury?
- 0 kali
0 times
 - 1 kali
1 time
 - 2 atau 3 kali
2 or 3 times
 - 4 atau 5 kali
4 or 5 times
 - 6 atau 7 kali
6 or 7 times
 - 8 atau 9 kali
8 or 9 times
 - 10 atau 11 kali
10 or 11 times
 - 12 kali atau lebih
12 or more times

30. Dalam tempoh 30 hari yang lepas, berapa kali seseorang di rumah menyatakan sesuatu yang menyakitkan hati atau menghina anda?

During the past 30 days, how many times has someone at home said hurtful or insulting things to you?

- a. 0 kali
0 times
- b. 1 kali
1 time
- c. 2 atau 3 kali
2 or 3 times
- d. 4 atau 5 kali
4 or 5 times
- e. 6 atau 7 kali
6 or 7 times
- f. 8 atau 9 kali
8 or 9 times
- g. 10 atau 11 kali
10 or 11 times
- h. 12 kali atau lebih
12 or more times

BAHAGIAN 5
PART 5

6 soalan seterusnya adalah berkenaan perasaan dan persahabatan anda.
The next 6 questions ask about your feelings and friendships.

31. Dalam tempoh 12 bulan yang lepas, berapa kerap anda merasa kesunyian?
During the past 12 months, how often have you felt lonely?

- a. Tidak pernah
Never
- b. Jarang-jarang
Rarely
- c. Kadang-kadang
Sometimes
- d. Kebanyakan masa
Most of the time
- e. Setiap masa
Always

32. Dalam tempoh 12 bulan yang lepas, berapa kerap anda merasa terlalu risau tentang sesuatu perkara sehingga anda tidak dapat tidur di waktu malam?

During the past 12 months, how often have you been so worried about something that you could not sleep at night?

- a. Tidak pernah
Never
- b. Jarang-jarang
Rarely
- c. Kadang-kadang
Sometimes
- d. Kebanyakan masa
Most of the time
- e. Setiap masa
Always

33. Dalam tempoh 12 bulan yang lepas, pernahkah anda terfikir secara serius untuk membunuh diri?

During the past 12 months, did you ever seriously consider attempting suicide?

- a. Ya
Yes
- b. Tidak
No

34. Dalam tempoh 12 bulan yang lepas, adakah anda membuat perancangan untuk membunuh diri?

During the past 12 months, did you make a plan about how you would attempt suicide?

- a. Ya
Yes
- b. Tidak
No

35. Dalam tempoh 12 bulan yang lepas, berapa kali anda telah cuba untuk membunuh diri?

During the past 12 months, how many times did you actually attempt suicide?

- a. 0 kali
0 times
- b. 1 kali
1 time
- c. 2 atau 3 kali
2 or 3 times
- d. 4 hingga 5 kali
4 to 5 times
- e. 6 kali atau lebih
6 or more times

36. Berapa ramai kawan rapat yang anda ada?
How many close friends do you have?

- a. 0 kawan
0 friends
- b. 1 kawan
1 friend
- c. 2 kawan
2 friends
- d. 3 atau lebih
3 or more

BAHAGIAN 6
PART 6

Rokok dan produk tembakau yang lain.
Cigarettes and other cigarette products.

37. Berapakah umur anda ketika kali pertama menghisap rokok?

How old were you when you first tried a cigarette?

a. Saya tidak pernah merokok
I have never smoked cigarettes

b. tahun/ *years old*

38. Dalam tempoh 30 hari yang lepas, berapa hari anda merokok?

During the past 30 days, on how many days did you smoke cigarettes?

a. 0 hari

0 days

b. 1 atau 2 hari

1 or 2 days

c. 3 hingga 5 hari

3 to 5 days

d. 6 hingga 9 hari

6 to 9 days

e. 10 hingga 19 hari

10 to 19 days

f. 20 hingga 29 hari

20 to 29 days

g. Kesemua 30 hari

All 30 days

39. Dalam tempoh 30 hari yang lepas, adakah anda menggunakan mana-mana produk seperti di bawah?

During the past 30 days, did you use any of the products listed below?

39.1 Shisha/*Shisha*

a. Ya/Yes

b. Tidak/No

39.2 Rokok Daun/*Traditional hand-rolled cigarette*

a. Ya/Yes

b. Tidak/No

39.3 Rokok gulung sendiri dengan kertas rokok/*“Roll-your-own” with cigarette paper*

a. Ya/Yes

b. Tidak/No

39.4 Cerut/ *Cigar*

a. Ya/Yes

b. Tidak/No

39.5 Hisap paip/*Pipe smoking*

a. Ya/Yes

b. Tidak/No

39.6 Rokok elektronik atau vape/*E-cigarettes or vape*

a. Ya/Yes

b. Tidak/No

39.7 Mengunyah tembakau (Sentil atau songel)/*Chewing tobacco*

a. Ya/Yes

b. Tidak/No

39.8 Menghidu tembakau/*Snuff*

a. Ya/Yes

b. Tidak/No

40. Dalam tempoh 12 bulan yang lepas, adakah anda cuba untuk berhenti merokok?
During the past 12 months, have you ever tried to stop smoking cigarettes?

- a. Saya tidak pernah merokok
I have never smoked cigarettes
- b. Saya tidak pernah merokok dalam tempoh 12 bulan yang lepas
I did not smoke cigarettes during the past 12 months
- c. Ya
Yes
- d. Tidak
No

41. Dalam tempoh 7 hari yang lepas, berapa hari anda bersama dengan perokok yang sedang merokok?
During the past 7 days, on how many days have people smoked in your presence?

- a. 0 hari
0 days
- b. 1 atau 2 hari
1 or 2 days
- c. 3 atau 4 hari
3 or 4 days
- d. 5 atau 6 hari
5 or 6 days
- e. Kesemua 7 hari
All 7 days

42. Siapa antara bapa, ibu atau penjaga yang menggunakan produk tembakau termasuk rokok?
Which of your parents or guardians use any form of tobacco including cigarettes?

- a. Kedua-duanya tidak merokok
Neither
- b. Ayah atau penjaga lelaki
My father or male guardian
- c. Ibu atau penjaga perempuan
My mother or female guardian
- d. Kedua-duanya
Both
- e. Tidak tahu
I do not know

43. Berapakah umur anda ketika mula-mula menggunakan rokok elektronik/vape?
How old were you when you first tried using e-cigarettes/vape?

- a. Saya tidak pernah gunakan rokok elektronik/vape
I have never tried using e-cigarettes/vape
- b. tahun/ years old

44. Dalam tempoh 30 hari yang lepas, berapa hari anda telah menggunakan rokok elektronik/vape?
During the past 30 days, on how many days did you use e-cigarettes/vape?

- a. 0 hari
0 days
- b. 1 atau 2 hari
1 or 2 days
- c. 3 hingga 5 hari
3 to 5 days
- d. 6 hingga 9 hari
6 to 9 days
- e. 10 hingga 19 hari
10 to 19 days
- f. 20 hingga 29 hari
20 to 29 days
- g. Kesemua 30 hari
All 30 days

45. Dalam tempoh 12 bulan yang lepas, adakah anda cuba untuk berhenti menggunakan rokok elektronik/vape?

During the past 12 months, did you ever try to stop using e-cigarettes/vape?

a. Saya tidak pernah menggunakan rokok elektronik/ vape

I have never used e-cigarettes/vape

b. Saya tidak pernah menggunakan rokok elektronik/ vape dalam tempoh 12 bulan yang lepas

I don't use e-cigarettes/vape in the past 12 months

c. Ya

Yes

d. Tidak

No

46. Siapakah antara bapa, ibu atau penjaga anda menggunakan rokok elektronik/vape?

Which of your parents or guardians use e-cigarettes/vape?

a. Kedua-duanya tidak merokok

Neither

b. Ayah atau penjaga lelaki

My father or male guardian

c. Ibu atau penjaga perempuan

My mother or female guardian

d. Kedua-duanya

Both

e. Tidak tahu

I do not know

**BAHAGIAN 7
 PART 7**

SILA BACA PERNYATAAN DI BAWAH:

6 soalan berikutnya adalah berkenaan meminum minuman beralkohol. Satu “minuman merujuk kepada satu gelas wain, tuak, lihing, bahar, ijuk atau todi; sebotol bir, segelas kecil arak, langkau, montoku; atau minuman campuran. Minuman beralkohol tidak termasuk beberapa hirup wain untuk tujuan keagamaan.

PLEASE READ THE STATEMENT BELOW:

The next 6 questions ask about drinking alcohol. A “drink’ is a glass of wine, tuak, lihing, bahar, ijuk or toddy; a bottle of beer, a small glass of liquor’ or mixed drink. Drinking alcohol does not include drinking a few sip of wine for religious purposes.

47. Berapakah umur anda ketika kali pertama anda minum minuman beralkohol?

How old were you when you had your first drink of alcohol?

- a. Saya tidak pernah minum minuman beralkohol
I have never had a drink of alcohol
- b. 7 tahun atau ke bawah
7 years old or younger
- c. 8 atau 9 tahun
8 or 9 years old
- d. 10 atau 11 tahun
10 or 11 years old
- e. 12 atau 13 tahun
12 or 13 years old
- f. 14 atau 15 tahun
14 or 15 years old
- g. 16 tahun atau ke atas
16 years old or older

48. Dalam tempoh 30 hari yang lepas, berapa hari anda mengambil sekurang-kurangnya satu minuman mengandungi alkohol?

During the past 30 days, on how many days did you have at least one drink containing alcohol?

- a. 0 hari
0 days
- b. 1 atau 2 hari
1 or 2 days
- c. 3 hingga 5 hari
3 to 5 days
- d. 6 hingga 9 hari
6 to 9 days
- e. 10 hingga 19 hari
10 to 19 days
- f. 20 hingga 29 hari
20 to 29 days
- g. Kesemua 30 hari
All 30 days

49. Dalam tempoh 30 hari yang lepas, pada hari anda minum minuman alkohol; berapa banyak minuman yang anda **biasa** ambil dalam sehari?

During the past 30 days, on the day you drank alcohol, how many drinks did you usually drink per day?

- Saya tidak minum minuman beralkohol dalam 30 hari yang lepas
I did not drink alcohol during the past 30 days
- Kurang dari 1 minuman
Less than one drink
- 1 minuman
1 drink
- 2 minuman
2 drinks
- 3 minuman
3 drinks
- 4 minuman
4 drinks
- 5 minuman atau lebih
5 or more drinks

50. Dalam tempoh 30 hari yang lepas, biasanya bagaimana anda mendapatkan minuman beralkohol? **SILA PILIH SATU JAWAPAN SAHAJA**

During the past 30 days, how did you usually get the alcohol you drank? SELECT ONLY ONE RESPONSE

- Saya tidak minum minuman beralkohol dalam 30 hari yang lepas
I did not drink alcohol during the past 30 days
- Saya beli dari kedai atau gerai
I brought it in a store, shop or from a street vendor
- Saya beri duit kepada orang lain untuk membeli
I gave someone else money to buy it for me
- Saya dapat daripada kawan
I got it from my friend
- Saya dapat daripada keluarga saya
I got it from my family
- Saya curi atau ambil tanpa kebenaran
I stole it or got it without permission
- Saya perolehi dari cara lain
I got it some other way

SILA BACA PERNYATAAN DI BAWAH:
 Terhuyung-hayang semasa berjalan, tidak mampu bercakap dengan betul, dan muntah adalah tanda seseorang itu terlalu mabuk.

PLEASE READ THE STATEMENT BELOW?
 Staggering when walking, not being able to speak right, and throwing up are some signs of being really drunk.

51. Sepanjang hidup anda berapa kali anda minum minuman beralkohol berlebihan sehingga betul-betul mabuk?

During your life, how many times did you drink so much alcohol that you were really drunk?

- 0 kali
0 times
- 1 atau 2 kali
1 or 2 times
- 3 hingga 9 kali
3 or 9 times
- 10 kali atau lebih
10 or more times

52. Sepanjang hidup anda, berapa kali anda mendapat masalah dengan ahli keluarga atau kawan, tidak ke sekolah, atau bertumbuk, akibat daripada minum minuman beralkohol?

During your life, how many times have you got into trouble with your family or friends, missed school, or got into fights, as a result of drinking alcohol?

- 0 kali
0 times
- 1 atau 2 kali
1 or 2 times
- 3 hingga 9 kali
3 or 9 times
- 10 kali atau lebih
10 or more times

BAHAGIAN 8
PART 8

SILA BACA PERNYATAAN DI BAWAH:

4 soalan berikutnya adalah berkenaan penggunaan dadah termasuk heroin, morfin, gam, amfitamin, ecstasy, syabu, ice dan ganja. Ini tidak termasuk ubat-ubatan preskripsi.

PLEASE READ THE STATEMENT BELOW:

The next 4 questions ask about drug use. This include heroin, morphine, glue, amphetamine, ecstasy, methamphetamine, ice and marijuana . This does not include prescribed medicine.

53. Berapakah umur anda ketika **pertama kali** anda menggunakan dadah?

*How old were you when you **first** used drugs?*

- a. Saya tidak pernah menggunakan dadah
I have never used drugs
- b. 7 tahun atau ke bawah
7 years old or younger
- c. 8 atau 9 tahun
8 or 9 years old
- d. 10 atau 11 tahun
10 or 11 years old
- e. 12 atau 13 tahun
12 or 13 years old
- f. 14 atau 15 tahun
14 or 15 years old
- g. 16 tahun atau ke atas
16 years old or older

54. Sepanjang hidup anda, berapa kali anda telah menggunakan dadah?

During your life, how many times have you used drugs?

- a. 0 kali
0 times
- b. 1 atau 2 kali
1 or 2 times
- c. 3 hingga 9 kali
3 or 9 times
- d. 10 hingga 19 kali
10 to 19 times
- e. 20 kali atau lebih
20 or more times

55. Dalam tempoh 30 hari yang lepas, berapa kali anda menggunakan dadah?

During the past 30 days, how many times have you used drugs?

- a. 0 kali
0 times
- b. 1 atau 2 kali
1 or 2 times
- c. 3 hingga 9 kali
3 or 9 times
- d. 10 hingga 19 kali
10 to 19 times
- e. 20 kali atau lebih
20 or more times

56. Dalam tempoh 30 hari yang lepas, bagaimana biasanya anda mendapatkan dadah yang anda gunakan? **SILA PILIH SATU JAWAPAN SAHAJA**

*During the past 30 days, how did you usually get the drugs used? **SELECT ONLY ONE RESPONSE***

- a. Saya tidak menggunakan dadah dalam 30 hari yang lepas
I did not use drugs during the past 30 days
- b. Saya beli dari orang lain
I bought them from someone
- c. Saya beri duit kepada orang lain untuk membeli
I gave someone else money to buy it for me
- d. Saya mencuri atau mengambil tanpa kebenaran
I stole it or got it without permission
- e. Saya mendapatkannya daripada kawan saya
I got it from my friend
- f. Saya mendapatkannya daripada keluarga saya
I got it from my family
- g. Saya memperolehi dari cara lain
I got it some other ways

57. Sepanjang hidup anda, berapa kali anda telah menggunakan ganja?

During your life, how many times have you used marijuana?

- a. 0 kali
0 times
- b. 1 atau 2 kali
1 or 2 times
- c. 3 hingga 9 kali
3 or 9 times
- d. 10 hingga 19 kali
10 to 19 times
- e. 20 kali atau lebih
20 or more times

58. Dalam tempoh 30 hari yang lepas, berapa kali anda menggunakan ganja?

During the past 30 days, how many times have you used marijuana?

- a. 0 kali
0 times
- b. 1 atau 2 kali
1 or 2 times
- c. 3 hingga 9 kali
3 or 9 times
- d. 10 hingga 19 kali
10 to 19 times
- e. 20 kali atau lebih
20 or more times

59. Sepanjang hidup anda, berapa kali anda telah menggunakan amfetamin atau metamfetamin (meth, syabu, ice, chase the dragon)?

During your life, how many times have you used amphetamines or metamphetamines (meth, syabu, ice, chase the dragon)?

- a. 0 kali
0 times
- b. 1 atau 2 kali
1 or 2 times
- c. 3 hingga 9 kali
3 or 9 times
- d. 10 hingga 19 kali
10 to 19 times
- e. 20 kali atau lebih
20 or more times

BAHAGIAN 9
PART 9

SILA BACA PERNYATAAN DI BAWAH:

5 soalan berikutnya adalah berkenaan hubungan seksual. Hubungan seksual adalah perlakuan seks yang melibatkan memasukkan zakar ke dalam faraj atau dubur.

PLEASE READ THE STATEMENT BELOW:

The next 5 questions ask about sexual intercourse. Sexual intercourse is defined as sexual acts of penetration of penis into vagina or anus.

60. Pernahkah anda melakukan hubungan seksual/persetubuhan dalam 30 hari yang lepas ?

Have you ever had sexual intercourse in the past 30 days?

- a. Ya
Yes
- b. No
Tidak

61. Berapa umur anda ketika kali pertama melakukan hubungan seksual/persetubuhan?

How old were you when you had sexual intercourse for the first time?

- a. Saya tidak pernah melakukan hubungan seksual/persetubuhan
I have never had sexual intercourse
- b. 11 tahun atau ke bawah
11 years old or younger
- c. 12 tahun
12 years old
- d. 13 tahun
13 years old
- e. 14 tahun
14 years old
- f. 15 tahun
15 years old
- g. 16 tahun atau ke atas
16 years old or older

62. Sepanjang hidup anda, berapa ramai orang yang telah anda lakukan hubungan seksual/persetubuhan?

During your life, with how many people have you had sexual intercourse?

- a. Saya tidak pernah melakukan hubungan seksual/persetubuhan
I have never had sexual intercourse
- b. 1 orang
1 person
- c. 2 orang
2 people
- d. 3 orang
3 people
- e. 4 orang
4 people
- f. 5 orang
5 people
- g. 6 orang atau lebih
6 or more people

63. Kali terakhir anda melakukan hubungan seksual/persetubuhan; adakah anda atau pasangan anda menggunakan kondom?

The last time you had sexual intercourse; did you or your partner use a condom?

- a. Saya tidak pernah melakukan hubungan seksual/persetubuhan
I have never had sexual intercourse
- b. Ya
Yes
- c. Tidak
No

64. Kali terakhir anda melakukan hubungan seksual/persetubuhan, adakah anda atau pasangan anda menggunakan kaedah pencegahan kehamilan lain seperti teknik pancutan luar, masa selamat, pil pencegah kehamilan, ataupun kaedah lain?

The last time you had sexual intercourse, did you or your partner use any other method of birth control, such as withdrawal, safe time, birth control pills, or any other method to prevent pregnancy?

- a. Saya tidak pernah melakukan hubungan seksual/persetubuhan
I have never had sexual intercourse
- b. Ya
Yes
- c. Tidak
No
- d. Tidak tahu
I do not know

BAHAGIAN 10
PART 10

SILA BACA PERNYATAAN DI BAWAH:

3 soalan seterusnya adalah berkenaan aktiviti fizikal. Aktiviti fizikal adalah apa-apa aktiviti yang meningkatkan kadar denyutan jantung dan menyebabkan kita bernafas dengan kuat. Aktiviti fizikal boleh dilakukan dalam sukan, bermain dengan kawan, atau berjalan ke sekolah. Contoh aktiviti fizikal termasuklah berlari, berjalan pantas, berbasikal, menari, dan bola sepak.

PLEASE READ THE STATEMENT BELOW:

The next 3 questions ask about physical activity. Physical activity is any activity that increases your heart rate and makes you breathe hard. Physical activity can be done in sports, playing with friends, or walking to school. Some examples of physical activity are running, fast walking, biking, dancing, and football.

65. Dalam tempoh 7 hari yang lepas, berapa hari anda melakukan aktiviti fizikal untuk sekurang-kurangnya 60 minit setiap hari? **JUMLAHKAN MASA ANDA MELAKUKAN APA-APA AKTIVITI FIZIKAL SETIAP HARI**

*During the past 7 days, on how many days were you physically active for a total of at least 60 minutes per day? **ADD UP ALL THE TIME YOU SPENT IN ANY KIND OF PHYSICAL ACTIVITY EACH DAY***

- a. 0 hari
0 days
- b. 1 hari
1 day
- c. 2 hari
2 days
- d. 3 hari
3 days
- e. 4 hari
4 days
- f. 5 hari
5 days
- g. 6 hari
6 days
- h. 7 hari
7 days

66. Dalam tempoh 7 hari yang lepas, berapa hari anda berjalan kaki atau berbasikal ke sekolah atau balik ke rumah?

During the past 7 days, on how many days did you walk or ride a bicycle to or from school?

- a. 0 hari
0 days
- b. 1 hari
1 day
- c. 2 hari
2 days
- d. 3 hari
3 days
- e. 4 hari
4 days
- f. 5 hari
5 days
- g. 6 hari
6 days
- h. 7 hari
7 days

SILA BACA PERNYATAAN DIBAWAH:

Soalan berikutnya adalah berkenaan masa yang anda habiskan dengan duduk semasa tidak di sekolah atau semasa membuat kerja rumah.

PLEASE READ THE STATEMENT BELOW:

The next question asks about the time you spend mostly sitting when you are not in school or doing homework.

67. Biasanya berapa masa yang anda habiskan untuk duduk sama ada untuk menonton televisyen, bermain permainan komputer, berbual dengan kawan atau apa-apa aktiviti yang memerlukan anda duduk?

How much time do you spend during a typical or usual day sitting and watching television, playing computer games, talking with friends, or doing other sitting activities?

- a. Kurang dari 1 jam sehari
Less than 1 hour per day
- b. 1 hingga 2 jam sehari
1 to 2 hours per day
- c. 3 hingga 4 jam sehari
3 to 4 hours per day
- d. 5 hingga 6 jam sehari
5 to 6 hours per day
- e. 7 hingga 8 jam sehari
7 to 8 hours per day
- f. Lebih dari 8 jam sehari
More than 8 hours per day

BAHAGIAN 11
PART 11

6 soalan seterusnya adalah berkenaan pengalaman anda di sekolah dan di rumah.
The next 6 questions ask about your experiences at school and at home.

68. Dalam tempoh 30 hari yang lepas, berapa hari anda tidak hadir ke kelas atau sekolah tanpa kebenaran?

During the past 30 days, on how many days did you miss classes or school without permission?

- a. 0 hari
0 days
- b. 1 atau 2 hari
1 or 2 days
- c. 3 hingga 5 hari
3 to 5 days
- d. 6 hingga 9 hari
6 to 9 days
- e. 10 hari atau lebih
10 or more days

69. Dalam tempoh 30 hari yang lepas, berapa kerap rakan sekolah anda bersikap baik hati dan suka membantu?

During the past 30 days, how often were most of the students in your school kind and helpful?

- a. Tidak pernah
Never
- b. Jarang-jarang
Rarely
- c. Kadang-kadang
Sometimes
- d. Kebanyakan masa
Most of the time
- e. Sentiasa
Always

70. Dalam tempoh 30 hari yang lepas, berapa kerap ibu bapa atau penjaga anda menyemak kerja sekolah anda?

During the past 30 days, how often did your parents or guardians check to see if your homework was done?

- a. Tidak pernah
Never
- b. Jarang-jarang
Rarely
- c. Kadang-kadang
Sometimes
- d. Kebanyakan masa
Most of the time
- e. Sentiasa
Always

71. Dalam tempoh 30 hari yang lepas, berapa kerap ibu bapa atau penjaga anda cuba memahami masalah dan kebimbangan anda?

During the past 30 days, how often did your parents or guardians try to understand your problems and worries?

- a. Tidak pernah
Never
- b. Jarang-jarang
Rarely
- c. Kadang-kadang
Sometimes
- d. Kebanyakan masa
Most of the time
- e. Sentiasa
Always

72. Dalam tempoh 30 hari yang lepas, berapa kerap ibu bapa atau penjaga anda benar-benar tahu apa yang anda lakukan pada masa lapang?

During the past 30 days how often did your parents or guardians really know what you were doing with your free time?

- a. Tidak pernah
Never
- b. Jarang-jarang
Rarely
- c. Kadang-kadang
Sometimes
- d. Kebanyakan masa
Most of the time
- e. Sentiasa
Always

73. Dalam tempoh 30 hari yang lepas, berapa kerap ibu bapa atau penjaga anda memeriksa barangan anda tanpa kebenaran?

During the past 30 days, how often did your parents or guardians go through your things without your approval?

- a. Tidak pernah
Never
- b. Jarang-jarang
Rarely
- c. Kadang-kadang
Sometimes
- d. Kebanyakan masa
Most of the time
- e. Sentiasa
Always

BAHAGIAN 12
PART 12

Penggunaan internet di kalangan remaja.
The use of internet among adolescents.

74. Adakah anda melayari internet dalam 30 hari yang lepas?
Did you surf the internet in the past 30 days?

- a. Ya
Yes
- b. Tidak (**Terus ke BAHAGIAN 13, soalan No. 78**)
No (Go to PART 13, question No. 78)

75. Apakah alat yang anda gunakan bagi melayari internet?

Which devices that you use for surfing internet?

75.1 Telefon pintar/ *smartphone*

- a. Ya/Yes
- b. Tidak/No

75.2 Komputer, Komputer Riba/
Computer, Laptop, Notebook

- a. Ya/Yes
- b. Tidak/No

75.3 Papan Elektronik Mudah Alih/
Tablet, Ipad

- a. Ya/Yes
- b. Tidak/No

76. Pernahkah anda belajar menggunakan internet dengan bijak dari.....?

Have you ever learned how to use the internet wisely from.....?

76.1 Ibubapa/penjaga/ *parents/guardian*

- a. Ya/Yes
- b. Tidak/No

76.2 Guru/ *teacher*

- a. Ya/Yes
- b. Tidak/No

76.3 Kawan/ *friend*

- a. Ya/Yes
- b. Tidak/No

76.4 Lain-lain sumber (cth: melalui pembelajaran sendiri, kempen/iklan)
Other sources (eg: through self-learning, campaign/advertisement)

- a. Ya/Yes
- b. Tidak/No

77. Ujian saringan penggunaan internet.
Internet usage screening test.

Arahan: Sila nilaikan setiap soalan dengan menggunakan pilihan “a” sehingga “e” daripada skala dibawah untuk menilai ketepatan fakta tersebut bagi diri anda dalam tempoh **30 HARI YANG LALU**. Sila bulatkan jawapan anda untuk semua soalan.

		Tidak pernah	Kadang-kadang	Agak Kerap	Sederhana Kerap	Sangat Kerap
77.1	Berapa kerapkah anda mendapati yang anda melayari internet lebih lama dari masa yang dirancang? <i>How often do you find that you stay on-line longer than you intended?</i>	a	b	c	d	e
77.2	Berapa kerapkah anda mengabaikan tugas pelajaran/ kerja rumah anda untuk menghabiskan lebih banyak masa melayari internet? <i>How often do you neglect household chores to spend more time on-line?</i>	a	b	c	d	e
77.3	Berapa kerapkah anda memilih keseronokan internet berbanding hubungan rapat yang terjalin dengan kawan/pasangan anda? <i>How often do you prefer the excitement of the Internet to intimacy with your partner?</i>	a	b	c	d	e
77.4	Berapa kerapkah anda memulakan hubungan baru dengan pengguna internet yang lain? <i>How often do you form new relationships with fellow on-line users?</i>	a	b	c	d	e
77.5	Berapa kerapkah orang-orang didalam hidup anda mengadu kepada anda mengenai jumlah masa yang anda gunakan untuk melayari internet? <i>How often do others in your life complain to you about the amount of time you spend on-line?</i>	a	b	c	d	e
77.6	Berapa kerapkah pelajaran kamu terjejas (cth. ponteng kelas, menangguhkan kerja, tidak menyiapkan tugas tepat pada masa, dll) disebabkan oleh masa yang anda habiskan untuk melayari internet? <i>How often do your grades or school work suffers because of the amount of time you spend on-line?</i>	a	b	c	d	e
77.7	Berapa kerapkah anda memeriksa email anda sebelum anda melakukan perkara lain? <i>How often do you check your email before something else that you need to do?</i>	a	b	c	d	e
77.8	Berapa kerapkah pencapaian akademik atau produktiviti anda terjejas disebabkan oleh penggunaan internet? <i>How often does your job performance or productivity suffer because of the Internet?</i>	a	b	c	d	e
77.9	Berapa kerapkah anda cuba mempertahankan diri atau berahsia apabila seseorang bertanya mengenai aktiviti anda (apa yang anda lakukan) di internet?	a	b	c	d	e

TINJAUAN KESIHATAN REMAJA | NHMS 2017
ADOLESCENT HEALTH SURVEY

	<i>How often do you become defensive or secretive when anyone asks you what you do on-line?</i>					
77.10	Berapa kerapkah anda cuba melupakan pemikiran yang mengganggu mengenai kehidupan anda dengan pemikiran yang boleh menenangkan di internet? <i>How often do you block out disturbing thoughts about your life with soothing thoughts of the Internet?</i>	a	b	c	d	e
77.11	Berapa kerapkah anda mendapati diri anda tertunggu-tunggu atau mengharapkan untuk segera menggunakan internet lagi apabila anda tidak menggunakannya? <i>How often do you find yourself anticipating when you will go on-line again?</i>	a	b	c	d	e
77.12	Berapa kerapkah anda merasa bimbang apabila memikirkan kehidupan tanpa internet akan menjadi bosan, kosong dan tidak menyeronokkan? <i>How often do you fear that life without the Internet would be boring, empty, and joyless?</i>	a	b	c	d	e
77.13	Berapa kerapkah anda marah, menjerit atau menunjukkan reaksi tidak gembira kepada seseorang yang mengganggu anda ketika anda melayari internet? <i>How often do you snap, yell, or act annoyed if someone bothers you while you are on-line?</i>	a	b	c	d	e
77.14	Berapa kerapkah anda kurang tidur disebabkan melayari internet hingga lewat malam? <i>How often do you lose sleep due to late-night log-ins?</i>	a	b	c	d	e
77.15	Berapa kerapkah anda asyik terfikir tentang internet atau berkhayal melayarinya walaupun anda tidak menggunakan internet pada masa itu? <i>How often do you feel preoccupied with the Internet when off-line, or fantasize about being on-line?</i>	a	b	c	d	e
77.16	Berapa kerapkah anda berkata kepada diri sendiri "sekejap lagi/ beberapa minit lagi" apabila anda melayari internet? <i>How often do you find yourself saying "just a few more minutes" when on-line?</i>	a	b	c	d	e
77.17	Berapa kerapkah anda cuba untuk mengurangkan masa anda melayari internet tetapi tidak berjaya? <i>How often do you try to cut down the amount of time you spend on-line and fail?</i>	a	b	c	d	e
77.18	Berapa kerapkah anda cuba menyorokkan daripada orang lain mengenai jumlah masa yang anda gunakan untuk melayari internet? <i>How often do you try to hide how long you've been on-line?</i>	a	b	c	d	e
77.19	Berapa kerapkah anda memilih untuk menghabiskan lebih banyak masa melayari internet berbanding keluar bersosial dengan rakan-rakan anda?	a	b	c	d	e

TINJAUAN KESIHATAN REMAJA | NHMS 2017
ADOLESCENT HEALTH SURVEY

	<i>How often do you choose to spend more time on-line over going out with others?</i>					
77.20	<p>Berapa kerapkah anda merasa sedih, berperasaan tidak menentu dan gementar apabila tidak melayari internet, di mana semua perasaan ini akan hilang sebaik saja anda menggunakan internet semula?</p> <p><i>How often do you feel depressed, moody or nervous when you are off-line, which goes away once you are back on-line?</i></p>	a	b	c	d	e

BAHAGIAN 13
PART 13

78. Saringan Minda Sihat
Healthy Mind Screening

Sila baca setiap kenyataan di bawah dan **HITAMKAN** jawapan anda pada KERTAS JAWAPAN berdasarkan jawapan **a, b, c,** atau **d** bagi menggambarkan keadaan anda sepanjang minggu yang lalu. Tiada jawapan betul atau salah. Jangan mengambil masa yang terlalu lama untuk menjawab mana-mana kenyataan.

*Please read each statement and **SHADE** numbers **a, b, c,** or **d** which indicates how much the statement applied to you over the past week. There are no right or wrong answers. Do not spend too much time on any statement.*

Skala pemarkahan adalah seperti berikut:
The rating scale is as follow:

- a. **Tidak langsung** menggambarkan keadaan saya
Did not apply to me at all
- b. **Sedikit atau jarang-jarang** menggambarkan keadaan saya
*Applied to me to some **degree** or **some of the time***
- c. **Banyak atau kerap kali** menggambarkan keadaan saya
*Applied to me to a **considerable degree** or a **good part of time***
- d. **Sangat banyak atau sangat kerap** menggambarkan keadaan saya
*Applied to me **very much,** or **most of the time***

78.1	Saya dapati diri saya sukar ditenteramkan <i>I found it hard to wind down</i>	a	b	c	d
78.2	Saya sedar mulut saya terasa kering <i>I was aware of dryness of my mouth</i>	a	b	c	d
78.3	Saya tidak dapat mengalami perasaan positif sama sekali <i>I couldn't seem to experience any positive feeling at all</i>	a	b	c	d
78.4	Saya mengalami kesukaran bernafas (contohnya pernafasan yang laju, tercungap-cungap walaupun tidak melakukan senaman fizikal) <i>I experienced breathing difficulty (e.g. excessively rapid breathing, breathlessness in the absence of physical exertion)</i>	a	b	c	d
78.5	Saya sukar untuk mendapatkan semangat bagi melakukan sesuatu perkara <i>I found it difficult to work up the initiative to do things</i>	a	b	c	d

78.6	Saya cenderung untuk bertindak keterlaluan dalam sesuatu keadaan <i>I tended to over-react to situations</i>	a	b	c	d
78.7	Saya rasa mengeletar (contohnya pada tangan) <i>I experience trembling (e.g. in the hands)</i>	a	b	c	d
78.8	Saya rasa saya menggunakan banyak tenaga dalam keadaan cemas <i>I felt that I was using a lot of nervous energy</i>	a	b	c	d
78.9	Saya bimbang keadaan di mana saya mungkin menjadi panik dan melakukan perkara yang membodohkan diri sendiri <i>I was worried about situation in which I might panic and make a fool of myself</i>	a	b	c	d
78.10	Saya rasa saya tidak mempunyai apa-apa untuk diharapkan <i>I felt that I had nothing to look forward to</i>	a	b	c	d
78.11	Saya mendapati diri saya semakin gelisah <i>I found myself getting agitated</i>	a	b	c	d
78.12	Saya rasa sukar untuk relaks <i>I found it difficult to relax</i>	a	b	c	d
78.13	Saya rasa sedih dan murung <i>I felt down-hearted and blue</i>	a	b	c	d
78.14	Saya tidakdapat menahan sabar dengan perkara yang menghalang saya meneruskan apa yang saya lakukan <i>I was intolerant of anything that kept me from getting on with what I was doing</i>	a	b	c	d
78.15	Saya rasa hampir-hampir menjadi panik/cemas <i>I felt I was close to panic</i>	a	b	c	d
78.16	Saya tidak bersemangat dengan apa jua yang saya lakukan <i>I was unable to become enthusiastic about anything</i>	a	b	c	d
78.17	Saya tidak begitu berharga sebagai seorang individu <i>I felt I wasn't worth much as a person</i>	a	b	c	d

78.18	Saya rasa yang saya mudah tersentuh <i>I felt that I was rather touchy</i>	a	b	c	d
78.19	Saya sedar tindakbalas jantung saya walaupun tidak melakukan aktiviti fizikal (contohnya kadar denyutan jantung bertambah, atau denyutan jantung berkurangan) <i>I was aware of the action of my heart in the absence of physical exertion (e.g. sense of heart rate increase, heart missing a beat)</i>	a	b	c	d
78.20	Saya berasa takut tanpa sebab yang munasabah <i>I felt scared without any good reason</i>	a	b	c	d
78.21	Saya rasa hidup ini tidak bermakna <i>I felt that life was meaningless</i>	a	b	c	d



INSTITUT KESIHATAN UMUM
KEMENTERIAN KESIHATAN MALAYSIA
Jalan Bangsar
50590 Kuala Lumpur

TINJAUAN KEBANGSAAN KESIHATAN DAN MORBIDITI 2017
KERTAS JAWAPAN TINJAUAN KESIHATAN REMAJA

ID PELAJAR

NEGERI	STRATA	KATEGORI SEKOLAH	KOD SEKOLAH	KELAS	PELAJAR
0 1 2 3 4 5 6 7 8 9	1 2	1 2	0 1 2 3 4 5 6 7 8 9	0 1 2 3 4 5 6 7 8 9	0 1 2 3 4 5 6 7 8 9

1 (A B C D E F G H)	34 (A B)	60 (A B)
2 (A B)	35 (A B C D E)	61 (A B C D E F G)
3 (A B C D E F)	36 (A B C D)	62 (A B C D E F G)
4 (A B C D E F)	37 (A B)	63 (A B C)
5 (A B C D E F)	38 (A B C D E F G)	64 (A B C D)
6 (A B C D E)	39.1 (A B)	65 (A B C D E F G H)
7 (A B C D E F G)	39.2 (A B)	66 (A B C D E F G H)
8 (A B C D E F G)	39.3 (A B)	67 (A B C D E F)
9 (A B C D E F G)	39.4 (A B)	68 (A B C D E)
10 (A B C D E F G)	39.5 (A B)	69 (A B C D E)
11 (A B C D E F G)	39.6 (A B)	70 (A B C D E)
12 (A B C D E F G H)	39.7 (A B)	71 (A B C D E)
13 (A B C D E F)	39.8 (A B)	72 (A B C D E)
14 (A B C)	40 (A B C D)	73 (A B C D E)
15 (A B)	41 (A B C D E)	74 (A B)
16 (A B C D E)	42 (A B C D E)	75.1 (A B)
17 (A B)	43 (A B)	75.2 (A B)
18 (A B)	44 (A B C D E F G)	75.3 (A B)
19 (A B C D E)	45 (A B C D)	76.1 (A B)
20 (A B C D E)	46 (A B C D E)	76.2 (A B)
21 (A B C D E)	47 (A B C D E F G)	76.3 (A B)
22 (A B C D E F G H)	48 (A B C D E F G)	76.4 (A B)
23 (A B C D E F G H)	49 (A B C D E F G)	77.1 (A B C D E)
24 (A B C D E F G H)	50 (A B C D E F G)	77.2 (A B C D E)
25 (A B C D E F G)	51 (A B C D)	77.3 (A B C D E)
26 (A B C D E F G H)	52 (A B C D)	77.4 (A B C D E)
27 (A B C D E F G)	53 (A B C D E F G)	77.5 (A B C D E)
28 (A B C D E F G H)	54 (A B C D E)	77.6 (A B C D E)
29 (A B C D E F G H)	55 (A B C D E)	77.7 (A B C D E)
30 (A B C D E F G H)	56 (A B C D E F G)	77.8 (A B C D E)
31 (A B C D E)	57 (A B C D E)	77.9 (A B C D E)
32 (A B C D E)	58 (A B C D E)	77.10 (A B C D E)
33 (A B)	59 (A B C D E)	77.11 (A B C D E)
		77.12 (A B C D E)
		77.13 (A B C D E)
		77.14 (A B C D E)
		77.15 (A B C D E)
		77.16 (A B C D E)
		77.17 (A B C D E)
		77.18 (A B C D E)
		77.19 (A B C D E)
		77.20 (A B C D E)





Ministry of Health Malaysia



MINISTRY OF
EDUCATION
MALAYSIA

ISBN 978-983-2387-58-9



9 789832 387589