

Methods

This was a nationwide population-based cross-sectional study targeting adults aged 18 or older in Malaysia (10). Two-stage stratified random sampling was performed to sample the respondents. Data collection was conducted from July to September 2023 and involved structured interviews, physical examinations, and venous blood taking. The **response rate was 71.4%, with 1,013 respondents** eligible for analysis. Analyses were conducted using Complex Samples procedures.

The definition of MetS followed the **Harmonised Joint Interim Statement 2009 criteria** that required at least three out of the following five metabolic abnormalities: (i) Waist circumference ≥ 90 cm for males and ≥ 80 cm for females; (ii) Systolic BP ≥ 130 and/or diastolic BP ≥ 85 mmHg or on treatment for hypertension; (iii) Fasting plasma glucose ≥ 5.6 mmol/L or previously diagnosed diabetes; (iv) Triglycerides ≥ 1.7 mmol/L or on treatment for triglycerides; (v) HDL-C < 1.0 mmol/L for males or < 1.3 mmol/L for females or on treatment for HDL-C (1).

References:

1. Alberti KG, Eckel RH, Grundy SM, Zimmet PZ, Cleeman JI, Donato KA, et al. Harmonizing the metabolic syndrome: a joint interim statement of the international diabetes federation task force on epidemiology and prevention; national heart, lung, and blood institute; American heart association; world heart federation; international atherosclerosis society; and international association for the study of obesity. *Circulation*. 2009;120(16):1640-5.
2. Mottillo S, Filion KB, Genest J, Joseph L, Pilote L, Poirier P, et al. The metabolic syndrome and cardiovascular risk: a systematic review and meta-analysis. *J Am Coll Cardiol*. 2010;56(14):1113-32.
3. Esposito K, Chiodini P, Colao A, Lenzi A, Giugliano D. Metabolic syndrome and risk of cancer: a systematic review and meta-analysis. *Diabetes Care*. 2012;35(11):2402-11.
4. Shin J-A, Lee J-H, Lim S-Y, Ha H-S, Kwon H-S, Park Y-M, et al. Metabolic syndrome as a predictor of type 2 diabetes, and its clinical interpretations and usefulness. *J Diabetes Investig*. 2013;4(4):334-43.
5. Noubiap JJ, Nansseu JR, Lontchi-Yimagou E, Nkeck JR, Nyaga UF, Ngouo AT, Tounouga DN, Tianyi FL, Foka AJ, Ndoadoumgue AL, Bigna JJ. Geographic distribution of metabolic syndrome and its components in the general adult population: A meta-analysis of global data from 28 million individuals. *Diabetes research and clinical practice*. 2022 Jun 1;188:109924.
6. Misra A, Khurana L. Obesity and the metabolic syndrome in developing countries. *The Journal of Clinical Endocrinology & Metabolism*. 2008 Nov 1;93(11_supplement_1):s9-30.
7. Oda E. Metabolic syndrome: its history, mechanisms, and limitations. *Acta diabetologica*. 2012 Apr;49:89-95.
8. Gao B, Zhang L, Zhao M. Underweight but metabolically abnormal phenotype: Metabolic features and its association with cardiovascular disease. *Eur J Intern Med*. 2016;29:46-51.
9. Shi TH. The influence of metabolic syndrome in predicting mortality risk among US adults: importance of metabolic syndrome even in adults with normal weight. *Prev Chronic Dis*. 2020;17.
10. Wan KS, Rifin HM, Yusoff MFM, Ratnam KKY, Chan WK, Mohamad M, et al. Prevalence of metabolic syndrome and metabolic dysfunction-associated fatty liver disease in Malaysia 2023: study protocol for a community-based nationwide cross-sectional survey. *BMJ open*. 2023;13(10):e074432.

This document was prepared by:

Kim Sui Wan, Halizah Mat Rifin, Kishwen Kanna Yoga Ratnam, Muhammad Fadhli Mohd Yusoff, Shubash Shander Ganapathy, Nur Rabia'tula Dawiyah Rahim, Norafiqah Shahirah Ayob, Noor Ani Ahmad

Conflict of interest:

There is no conflict of interest

Acknowledgement:


The authors would like to thank the Director-General of Health Malaysia for permission to publish this document

This document should be cited as:

Wan KS, Mat Rifin H, Yoga Ratnam KK, Mohd Yusoff MF, Ganapathy SS, Rahim NRD, Ayob NS, Ahmad NA. Research highlight. The metabolic syndrome epidemic in Malaysia "A ticking time bomb". Prevalence of metabolic syndrome and metabolic dysfunction-associated fatty liver disease in Malaysia 2023. Institute for Public Health (IPH); 2024. Available from: <https://iku.gov.my/mets>

This document is based on:

The prevalence of metabolic syndrome and metabolic dysfunction-associated fatty liver in Malaysia 2023



Ministry of Health Malaysia
Institute for Public Health

PREVALENCE OF METABOLIC SYNDROME AND METABOLIC DYSFUNCTION-ASSOCIATED FATTY LIVER DISEASE IN MALAYSIA 2023

For further information, please visit www.iku.gov.my/mets or contact Dr Wan Kim Sui at kimsui@moh.gov.my



This research highlight is based on the study on **The prevalence of metabolic syndrome and metabolic dysfunction-associated fatty liver in Malaysia 2023**

Who is this publication for?

- Disease Control Division
- Family Health Development Division
- Medical Development Division
- Health Education Division
- Nutrition Division
- Public health practitioners
- Clinicians

Purpose of this document

To highlight the high prevalence of metabolic syndrome in Malaysia and propose policy and clinical recommendations


Disclaimer

The views, interpretation, implications, conclusions and recommendations are those of the authors alone and do not necessarily represent the opinions of the investigators participating in the project nor the views or policy of the Ministry of Health, Malaysia.

Project reference number:

NMRR-22-0284-GUT

Funded by:



MINISTRY OF HEALTH MALAYSIA

The Metabolic Syndrome Epidemic in Malaysia

"A ticking time bomb"

Issue

Metabolic syndrome (**MetS**) is a **cluster of cardiometabolic dysfunctions** characterised by high fasting plasma glucose, waist circumference, blood pressure, triglycerides, and low HDL cholesterol (1).

People with the condition have higher risks for adverse **cardiovascular** outcomes and **all-cause mortality** (2).


MetS is highly predictive of new-onset **type 2 diabetes** and is associated with increased risks of many **cancers**, including liver, colorectal, pancreatic, and breast cancers (3, 4).

According to a meta-analysis, the prevalence of MetS using the JIS criteria was **31.4% globally, 34.1% in upper-middle-income countries**, and **29.5% in the Western Pacific region** (5).

MetS is a disorder with a **high socioeconomic impact** on global health (6).

Despite the clinical and public health ramifications of MetS, there is a knowledge gap on the latest prevalence of MetS in Malaysia.

Key Messages



or a staggering 8.5 million adults in Malaysia have MetS!

- MetS is more common among **older age categories**, those with **lower education levels**, and the **B40 income group**.
- MetS is **not uncommon among underweight and normal-weight** individuals in Malaysia and should not be overlooked. They are also at increased risk of cardiovascular disease, kidney disease, and mortality.
- **Urgent decisive actions** are needed to manage the epidemic in Malaysia

Background


In the late twentieth century, insulin resistance was shown to be fundamental to metabolic syndrome (MetS), which means a constellation of obesity-related metabolic derangements predisposing to type 2 diabetes and cardiovascular disease (7).

Many diagnostic criteria for MetS have been established in recent decades. The prominent ones include the first definition by the World Health Organization (WHO) in 1998, followed by the United States National Cholesterol Education Program Adult Treatment Panel III (NCEP-ATP III) in 2001, and the International Diabetes Federation (IDF) in 2005 (8).


In 2009, the Joint Interim Statement (JIS) was introduced to harmonise the differences between various diagnostic definitions and is being widely adopted by studies worldwide (1).

Key Findings


Prevalence of individual metabolic risk factors among adults in Malaysia




Prevalence of multiple metabolic risk factors among adults in Malaysia




Prevalence of metabolic syndrome (≥3 risk factors)



Prevalence of MetS by sociodemographic factors and body mass index



Clinical characteristics of individuals with MetS



They are **young** in their economically productive age and have **many comorbidities** with **unfavourable clinical profiles**

Key Considerations

Policymakers & programme managers

- **Acknowledge the issue and raise priority of MetS** in the national health agenda-setting. For example, MetS can be considered to be included in the next National Strategic Plan for Non-Communicable Diseases.
- **Allocate sufficient resources** to operationalise the plan, including disease surveillance and clinical management.
- **Expand population screening programmes** such as the National Health Screening Initiative (NHSI) to include MetS screening, especially among high-risk subpopulations such as older age categories, the B40 group and those with lower education levels. Lipid panels can be checked using finger prick capillary blood for point-of-care testing.
- **Health promotion** via traditional and social media to increase awareness and knowledge of MetS among the general population is critical. The information should include **metabolically obese normal weight (MONW)**, which has an increased risk of cardiovascular disease, kidney disease, and mortality (8, 9).

Healthcare practitioners

- **Increase MetS management knowledge** among healthcare personnel, which includes primary care providers and specialists from relevant disciplines.
- The development of **Clinical Practice Guidelines** on MetS can help bring experts from different backgrounds to harmonise the management of patients.
- Patient management should focus on a **multidisciplinary approach** with a clear specialist referral pathway.
- Healthcare personnel, especially those in primary care, should **actively screen, diagnose, and manage MetS** to prevent complications.

Summary of Action Points

- Raise the priority of MetS in the national health agenda-setting
- Include MetS screening in existing population screening programmes
- Health promotion to increase awareness and knowledge among the general population
- Increase knowledge about MetS management among healthcare personnel